

BARRIER II WOUND CARE WITH PAIN RELIEF- povidone-iodine, lidocaine, and alcohol solution

Aurora Pharmaceutical, Inc.

Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click here.

BARRIER II®

Wound Care Spray With Pain Relief

INDICATIONS

For topical use on animals as an aid in preventing infection and reducing pain. Kills up to 99% of surface germs in 15 seconds or less that can potentially cause an infection.

ACTIVE INGREDIENTS	PURPOSE
Povidone-Iodine (2% available iodine)	Antiseptic
Lidocaine 2% w/v	Pain Relief
Ethyl Alcohol - less than 80%	Antiseptic
Isopropyl Alcohol - less than 5%	Antiseptic

DIRECTIONS

Hold sprayer about 4-6 inches from area to be treated. Spray 1 or more times - allow to dry. Repeat as needed. Forms a highly visible coating that is durable.

WARNINGS

Do not use in the eyes. Do not use if allergic or sensitive to the active ingredients. Prolonged exposure may cause irritation. If swallowed, get medical help or contact a Poison Control Center right away. Avoid inhalation.

Store upright at 15°-25° C (59°-77° F). Brief excursions up to 40° C (104° F) are permitted.

**FLAMMABLE — USE WITH CAUTION
FOR EXTERNAL USE ONLY
KEEP OUT OF REACH OF CHILDREN
WET HAIR OR FUR IS FLAMMABLE**

Limited Quantity



REORDER

NO: 19004

MANUFACTURED BY:

Aurora Pharmaceutical, Inc.

NORTHFIELD, MN 55057

1-888-215-1256

www.aurorapharmaceutical.com

IN 50-1121 06/2023

MANUFACTURED IN THE USA

PRINCIPAL DISPLAY PANEL - 473 ml Bottle Label

NDC 51072-041-16

BARRIER II®

Wound Care Spray

With Pain Relief

2% Available Iodine with Lidocaine

Does Not Contain Bitrex

(Patent pending)

For Veterinary Use Only

16 fl. oz. (473 ml)

AURORA PHARMACEUTICAL®

INDICATIONS: For topical use on animals as an aid in preventing infection and reducing pain. Kills up to 99% of surface germs in 15 seconds or less that can potentially cause an infection.

ACTIVE INGREDIENTS
Povidone-Iodine (2% available iodine)
Lidocaine 2% w/v
Ethyl Alcohol – less than 80%
Isopropyl Alcohol – less than 5%

DIRECTIONS: Hold sprayer about 4–6 inches from area to be treated. Spray 1 or more times — allow to dry. Repeat as needed. Forms a highly visible coating that is durable.

WARNINGS: Do not use in the eyes. Do not use if allergic or sensitive to the active ingredients. Prolonged exposure may cause irritation. If swallowed, get medical help or contact a Poison Control Center right away. Avoid inhalation.

Store upright at 15°–25° C (59°–77° F). Brief excursions up to 40° C (104° F) are permitted.

FLAMMABLE — USE WITH CAUTION FOR EXTERNAL USE ONLY
KEEP OUT OF REACH OF CHILDREN
WET HAIR OR FUR IS FLAMMABLE
Limited Quantity

TAKE TIME  OBSERVE LABEL DIRECTIONS

REORDER NO: 19004
MANUFACTURED BY:
Aurora Pharmaceutical, Inc.
NORTHFIELD, MN 55057
1-888-215-1256
www.aurorapharmaceutical.com
IN 50-1121 06/2023

NDC 51072-041-16

BARRIER II®

Wound Care Spray
With Pain Relief

2% Available Iodine With Lidocaine
Does Not Contain Bitrex
(Patent pending)

For Veterinary Use Only
16 fl. oz. (473 mL)



INDICACIONES: Para uso tópico en animales para ayudar a prevenir la infección y a reducir dolor. Mata hasta el 99% de los gérmenes superficiales en 15 segundos o menos, que potencialmente pueden causar una infección.

INGREDIENTES ACTIVOS	PROPÓSITO
Yodopovidona (Yodo al 2%)	Antiséptico
Lidocaína 2% p/v	Anestésico local
Alcohol etílico – menos del 80%	Antiséptico
Alcohol isopropílico – menos del 5%	Antiséptico

INSTRUCCIONES DE USO: Mantener rociador a unas 4–6 pulgadas de la zona a tratar. Rocíar una o más veces, dejar a secar. Repetir según sea necesario. Forma una capa que es altamente visible y duradera.

ADVERTENCIAS: No utilice en los ojos. No utilice en caso de alergia o sensibilidad a los ingredientes activos. La exposición prolongada puede causar irritación. En caso de ingestión, busque ayuda médica o contacte al Centro de Control de Envenenamiento (Poison Control Center) inmediatamente. Evite la inhalación.

Almacenar verticalmente a 15°–25° C (59°–77° F). Excursiones breves hasta 40° C (104° F) son admisibles.

INFLAMABLE – USAR CON CUIDADO
SÓLO PARA USO EXTERNO
MANTENER FUERA DEL ALCANCE DE LOS NIÑOS
PELO O PELAJE MOJADO ES INFLAMABLE
Cantidad Limitada

MANUFACTURED IN THE USA 

BARRIER II WOUND CARE WITH PAIN RELIEF

povidone-iodine, lidocaine, and alcohol solution

Product Information

Product Type	OTC ANIMAL DRUG	Item Code (Source)	NDC:51072-041
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
Povidone-Iodine (UNII: 85H0HZU99M) (Iodine - UNII:9679TC07X4)	Iodine	2 g in 100 mL
Lidocaine (UNII: 98PI200987) (Lidocaine - UNII:98PI200987)	Lidocaine	2 g in 100 mL
ALCOHOL (UNII: 3K9958V90M) (ALCOHOL - UNII:3K9958V90M)	ALCOHOL	80 mL in 100 mL
ISOPROPYL ALCOHOL (UNII: ND2M416302) (ISOPROPYL ALCOHOL - UNII:ND2M416302)	ISOPROPYL ALCOHOL	5 mL in 100 mL

Product Characteristics

Color	BROWN	Score	
Shape		Size	
Flavor		Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:51072-041-16	473 mL in 1 BOTTLE		
2	NDC:51072-041-01	3790 mL in 1 BOTTLE		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
UNAPPROVED DRUG OTHER		03/28/2011	

Labeler - Aurora Pharmaceutical, Inc. (832848639)

Establishment

Name	Address	ID/FEI	Business Operations
Aurora Pharmaceutical, Inc.		832848639	MANUFACTURE