

**DAYTIME NIGHTTIME COLD AND FLU- acetaminophen, dextromethorphan hydrobromide, phenylephrine hydrochloride doxylaminesuccinate**  
**Gobrand, Inc. (Goodnow)**

-----  
**Drug Facts**

**Active ingredients (in each 15 mL) DAYTIME**

Acetaminophen 325 mg

Dextromethorphan HBr 10 mg

Phenylephrine HCl 5 mg

**Active ingredients for (in each 30 mL) NIGHTTIME**

Acetaminophen 650 mg

Dextromethorphan HBr 30 mg

Doxylamine Succinate 12.5 mg

**Purposes for Day Time**

Pain reliever/fever reducer

Cough suppressant

Nasal decongestant

**Purpose for Night Time**

Pain reliever/fever reducer

Cough suppressant

Antihistamine

**Uses**

**DAYTIME**

- temporarily relieves common cold and flu symptoms
  - minor aches and pains
  - headache
  - sore throat
  - nasal congestion
  - fever
  - cough due to minor throat and bronchial irritation

**NIGHTTIME**

- temporarily relieves these common cold/flu symptoms`
  - minor aches and pains
  - headache
  - sore throat
  - runny nose and sneezing
  - fever
  - cough due to minor throat and bronchial irritation

## Warnings

### DAYTIME

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if:

- adult takes more than 4 doses (30 mL each) of acetaminophen in 24 hours, which is the maximum daily amount
- child takes more than 4 doses (15 mL each) in 24 hours
- taken with other drugs containing acetaminophen
- adult has 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash.

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

### NIGHTTIME

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash.

if a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

## **Do not use**

### **DAYTIME NIGHTTIME**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist
- If you are taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your child's prescription drug contains an MAOI, ask a doctor or pharmacist before giving this product.

## **Ask a doctor before use if you have**

### **DAYTIME**

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- a sodium-restricted diet
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema
- cough that occurs with too much phlegm (mucus)

### **NIGHTTIME**

- liver disease
- glaucoma
- cough that occurs with too much phlegm (mucus)
- trouble urinating due to an enlarged prostate gland
- a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis, or emphysema

## **Ask a doctor or pharmacist before use if you are**

### **DAYTIME**

taking the blood thinning drug warfarin.

### **NIGHTTIME**

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

## **When using this product**

### **DAYTIME**

**do not exceed recommended dosage.**

### **NIGHTTIME**

- excitability may occur, especially in children
- marked drowsiness may occur

- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives and tranquilizers may increase drowsiness

**Stop use and ask a doctor if**

**DAYTIME**

- nervousness, dizziness or sleeplessness occur
- pain, nasal congestion, or cough gets worse, or lasts more than 5 days(children) or 7 days (adult)
- fever gets worse, or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with rash or headache that lasts.

These could be signs of a serious condition.

**NIGHTTIME**

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with rash, or headache that lasts.

These could be signs of a serious condition.

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

**DAYTIME NIGHTTIME**

Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center(1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

**DAYTIME**

- **do not take more than directed (see Overdose warning)**
- do not take more than 4 doses in any 24-hour period
- measure only with dosing cup provided. Do not use any other dosing device.
- keep dosing cup with product
- mL= milliliter

adults and children 12 years and over	30 mL every 4 hours
children 6 to under 12 years	15 mL every 4 hours

children 4 to under 6 years	ask a doctor
children under 4 years	do not use

## **NIGHTTIME**

- do not take more than directed (see Overdose warning)
- Do not take more than 4 doses in any 24-hours period
- measure only with dosing cup provided. Do not use any other dosing device.
- mL= milliliter
- keep dosing cup with product
- adults and children 12 years and over: 30 mL every 6 hours
- children under 12 years of age: do not use

## **Other information**

### **DAYTIME**

- **each 15 mL contains:** sodium 12 mg
- store between 20°-25°C (68°-77°). Do not refrigerate

### **NIGHTTIME**

- **each 30 mL contains:** potassium 5 mg
- **each 30 mL contains** sodium 24 mg
- store between 20°-25°C (68°-77°F). Do not refrigerate

## **Inactive ingredients**

### **Day Time**

citric acid, FD&C yellow #6, flavor, glycerin, propylene glycol, purified water, saccharin sodium, sodium benzoate, sodium chloride, sodium citrate, sorbitol, sucralose, xanthan gum

### **Night Time**

acesulfame potassium, alcohol, anhydrous citric acid, FD&C blue #1, FD&C red #40, flavor, high fructose corn syrup, polyethylene glycol, propylene glycol, purified water, saccharin sodium trisodium citrate dihydrate

## **Questions or comments?**

Call **1-877-753-3935** Monday-Friday 9AM-5PM EST

## **Principal Display Panel**

**Compare to active ingredients in Vicks® DayQuil® & NyQuil® Cold & Flu**

### **DAYTIME**

NON-DROWSY

Cold + Flu

Day Relief

Acetaminophen

Dextromethorphan HBr

Phenylephrine HCl

Pain Reliever

Fever Reducer

Cough Suppressant

Nasal Decongestant

Alcohol-free / Antihistamine-free

Multi-Symptom Relief

Headaches & Fever

Sore Throat

Minor Aches & Pains

Nasal Congestion

Cough

**NIGHTTIME**

**CHERRY FLAVOR**

**Cold + Flu**

**Night Relief**

**Acetaminophen**

**Dextromethorphan HBr**

**Doxylamine Succinate**

Pain Reliever

Fever Reducer

Cough Suppressant

Antihistamine

**Alcohol 10%**

Multi-Symptom Relief

Headache & Fever

Sore Throat

Minor Aches & Pains

Sneezing

Running Nose

Cough

FL OZ (mL)

When using other Daytime or Nighttime products, carefully read each label to ensure correct dosing.

†This product is not manufactured or distributed by The Procter & Gamble Company. Vicks®, DayQuil® and NyQuil® are registered trademarks of The Procter & Gamble Company.

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND DOSAGE CUP OR UNDER CAP IS BROKEN OR MISSING.

DISTRIBUTED BY: GOBRANDS, INC.

PHILADELPHIA, PA 19123

Product Label



GOOD NOW Cold + Flu Day Relief Cold + Flu Night Relief Cherry Flavor

DAYTIME NIGHTTIME COLD AND FLU

acetaminophen, dextromethorphan hydrobromide, phenylephrine hydrochloride doxylaminesucinate kit

Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:82501-4672
---------------------	----------------	---------------------------	----------------

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:82501-4672-4	1 in 1 KIT; Type 0: Not a Combination Product	04/29/2022	

## Quantity of Parts

Part #	Package Quantity	Total Product Quantity
Part 1	1 BOTTLE	355 mL
Part 2	1 BOTTLE	355 mL

## Part 1 of 2

### COLD AND FLU RELIEF

acetaminophen, dextromethorphan hydrobromide, phenylephrine hydrochloride liquid

## Product Information

Item Code (Source)	NDC:82501-4661
Route of Administration	ORAL

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg in 15 mL
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg in 15 mL
<b>PHENYLEPHRINE HYDROCHLORIDE</b> (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg in 15 mL

## Inactive Ingredients

Ingredient Name	Strength
<b>ANHYDROUS CITRIC ACID</b> (UNII: XF417D3PSL)	
<b>FD&amp;C YELLOW NO. 6</b> (UNII: H77VEI93A8)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>TRISODIUM CITRATE DIHYDRATE</b> (UNII: B22547B95K)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>SACCHARIN SODIUM</b> (UNII: SB8ZUX40TY)	
<b>SODIUM BENZOATE</b> (UNII: OJ245FE5EU)	
<b>SODIUM CHLORIDE</b> (UNII: 451W47IQ8X)	
<b>SORBITOL</b> (UNII: 506T60A25R)	
<b>SUCRALOSE</b> (UNII: 96K6UQ3ZD4)	
<b>XANTHAN GUM</b> (UNII: TTV12P4NEE)	

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1		355 mL in 1 BOTTLE; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	04/29/2022	

## Part 2 of 2

### NIGHTTIME COLD AND FLU

acetaminophen, dextromethorphan hydrobromide, doxylamine succinate liquid

## Product Information

Item Code (Source)	NDC:82501-3431
Route of Administration	ORAL

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	650 mg in 30 mL
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	30 mg in 30 mL
<b>DOXYLAMINE SUCCINATE</b> (UNII: V9BI9B5YI2) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	12.5 mg in 30 mL

## Inactive Ingredients

Ingredient Name	Strength
<b>ACESULFAME POTASSIUM</b> (UNII: 23OV73Q5G9)	
<b>ANHYDROUS CITRIC ACID</b> (UNII: XF417D3PSL)	
<b>ALCOHOL</b> (UNII: 3K9958V90M)	
<b>FD&amp;C BLUE NO. 1</b> (UNII: H3R47K3TBD)	
<b>FD&amp;C RED NO. 40</b> (UNII: WZB9127XOA)	
<b>HIGH FRUCTOSE CORN SYRUP</b> (UNII: XY6UN3QB6S)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>TRISODIUM CITRATE DIHYDRATE</b> (UNII: B22547B95K)	
<b>SACCHARIN SODIUM</b> (UNII: SB8ZUX40TY)	

## Product Characteristics

Color		Score	
Shape		Size	
Flavor	CHERRY	Imprint Code	

**Contains****Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1		355 mL in 1 BOTTLE; Type 0: Not a Combination Product		

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	04/29/2022	

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	04/29/2022	

**Labeler** - Gobrand, Inc. (Goodnow) (057499049)

Revised: 10/2024

Gobrand, Inc. (Goodnow)