

**PREVIFEM- norgestimate and ethinyl estradiol**  
**RPK Pharmaceuticals, Inc.**

-----  
**HIGHLIGHTS OF PRESCRIBING INFORMATION**

These highlights do not include all the information needed to use PREVIFEM® or TRI-PREVIFEM® safely and effectively. See full prescribing information for PREVIFEM® and TRI-PREVIFEM®.

PREVIFEM® and TRI-PREVIFEM® (norgestimate/ethinyl estradiol tablets), for oral use  
Initial U.S. Approval: 1989

**WARNING: CIGARETTE SMOKING AND SERIOUS CARDIOVASCULAR EVENTS**

*See full prescribing information for complete boxed warning.*

- Previfem® or Tri-Previfem® (norgestimate/ethinyl estradiol tablets) are contraindicated in women over 35 years old who smoke. (4)
- Cigarette smoking increases the risk of serious cardiovascular events from combination oral contraceptives (COC) use. (4)

-----  
**RECENT MAJOR CHANGES**

Contraindications (4) 08/2017  
Warnings and Precautions (5.3) 08/2017

-----  
**INDICATIONS AND USAGE**

Previfem® (norgestimate/ethinyl estradiol tablets) and Tri-Previfem® (norgestimate/ethinyl estradiol tablets) are estrogen/progestin COCs, indicated for use by women to prevent pregnancy. (1.1)

Tri-Previfem® (norgestimate/ethinyl estradiol tablets) is also indicated for the treatment of moderate acne vulgaris in females at least 15 years of age, who have no known contraindications to oral contraceptive therapy and have achieved menarche.

Tri-Previfem® (norgestimate/ethinyl estradiol tablets) should be used for the treatment of acne only if the patient desires an oral contraceptive for birth control. (1.2)

-----  
**DOSAGE AND ADMINISTRATION**

- Take one tablet daily by mouth at the same time every day. (2.2)
- Take tablets in the order directed on the blister pack. (2.2)
- Do not skip or delay tablet intake. (2.2)

-----  
**DOSAGE FORMS AND STRENGTHS**

Previfem® (norgestimate/ethinyl estradiol tablets) consists of 28 round, unscored, film-coated tablets in the following order (3):

- 21 blue tablets each containing 0.25 mg norgestimate and 0.035 mg ethinyl estradiol
- 7 light-green tablets (inert)

Tri-Previfem® (norgestimate/ethinyl estradiol tablets) consists of 28 round, unscored, film-coated tablets in the following order (3):

- 7 white tablets each containing 0.18 mg norgestimate and 0.035 mg ethinyl estradiol
- 7 light blue tablets each containing 0.215 mg norgestimate and 0.035 mg ethinyl estradiol
- 7 blue tablets each containing 0.25 mg norgestimate and 0.035 mg ethinyl estradiol
- 7 light-green tablets (inert)

-----  
**CONTRAINDICATIONS**

- A high risk of arterial or venous thrombotic diseases (4)
- Liver tumors or liver disease (4)
- Undiagnosed abnormal uterine bleeding (4)
- Pregnancy (4)
- Breast cancer or other estrogen- or progestin-sensitive cancer (4)
- Co-administration with Hepatitis C drug combinations containing ombitasvir/paritaprevir/ritonavir, with or without dasabuvir (4)

-----  
**WARNINGS AND PRECAUTIONS**

- **Thromboembolic Disorders and Other Vascular Problems:** Stop Previfem® or Tri-Previfem® if a thrombotic event occurs. Stop at least 4 weeks before and through 2 weeks after major surgery. Start no earlier than 4 weeks after delivery, in women who are not breastfeeding. (5.1)
- **Liver disease:** Discontinue Previfem® or Tri-Previfem® if jaundice occurs. (5.2)
- **High blood pressure:** If used in women with well-controlled hypertension monitor blood pressure and stop Previfem® or Tri-Previfem® if blood pressure rises significantly. (5.4)

- Carbohydrate and lipid metabolic effects: Monitor prediabetic and diabetic women taking Prevfem<sup>®</sup> or Tri-Previfem<sup>®</sup>. Consider an alternate contraceptive method for women with uncontrolled dyslipidemia. (5.6)
- Headache: Evaluate significant change in headaches and discontinue Prevfem<sup>®</sup> or Tri-Previfem<sup>®</sup> if indicated. (5.7)
- Bleeding Irregularities and Amenorrhea: Evaluate irregular bleeding or amenorrhea. (5.8)

----- **ADVERSE REACTIONS** -----

The most common adverse reactions reported during clinical trials (≥2%) were:

Prevfem<sup>®</sup>: headache/migraine, abdominal/gastrointestinal pain, vaginal infection, genital discharge, breast issues (including breast pain, discharge, and enlargement), mood disorders (including depression and mood altered), flatulence, nervousness, rash. (6.1)

Tri-Previfem<sup>®</sup>: headache/migraine, breast issues (including breast pain, enlargement, and discharge), vaginal infection, abdominal/gastrointestinal pain, mood disorders (including mood alteration and depression), genital discharge, changes in weight (including weight increased or decreased). (6.1)

**To report SUSPECTED ADVERSE REACTIONS, contact Par Pharmaceutical at 1-800-828-9393 or FDA at 1-800-FDA-1088 or [www.fda.gov/medwatch](http://www.fda.gov/medwatch).**

----- **DRUG INTERACTIONS** -----

Drugs or herbal products that induce certain enzymes including CYP3A4, may decrease the effectiveness of COCs or increase breakthrough bleeding. Counsel patients to use a back-up or alternative method of contraception when enzyme inducers are used with COCs. (7.1)

----- **USE IN SPECIFIC POPULATIONS** -----

Nursing mothers: Not recommended; can decrease milk production. (8.3)

**See 17 for PATIENT COUNSELING INFORMATION and FDA-approved patient labeling.**

**Revised: 8/2017**

**FULL PRESCRIBING INFORMATION: CONTENTS\***

**WARNING: CIGARETTE SMOKING AND SERIOUS CARDIOVASCULAR EVENTS**

**1 INDICATIONS AND USAGE**

- 1.1 Oral Contraceptive
- 1.2 Acne

**2 DOSAGE AND ADMINISTRATION**

- 2.1 How to Start Prevfem<sup>®</sup> (norgestimate/ethinyl estradiol tablets) or Tri-Previfem<sup>®</sup> (norgestimate/ethinyl estradiol tablets)
- 2.2 How to Take Prevfem<sup>®</sup> or Tri-Previfem<sup>®</sup>
- 2.3 Missed Tablets
- 2.4 Advice in Case of Gastrointestinal Disturbances
- 2.5 Tri-Previfem<sup>®</sup> Use for Acne

**3 DOSAGE FORMS AND STRENGTHS**

**4 CONTRAINDICATIONS**

**5 WARNINGS AND PRECAUTIONS**

- 5.1 Thromboembolic Disorders and Other Vascular Problems
- 5.2 Liver Disease
- 5.3 Risk of Liver Enzyme Elevations with Concomitant Hepatitis C Treatment
- 5.4 High Blood Pressure
- 5.5 Gallbladder Disease
- 5.6 Carbohydrate and Lipid Metabolic Effects
- 5.7 Headache
- 5.8 Bleeding Irregularities and Amenorrhea
- 5.9 COC Use Before or During Early Pregnancy
- 5.10 Depression
- 5.11 Carcinoma of Breast and Cervix
- 5.12 Effect on Binding Globulins
- 5.13 Monitoring
- 5.14 Hereditary Angioedema

5.15 Chloasma

## **6 ADVERSE REACTIONS**

6.1 Clinical Trial Experience

6.2 Postmarketing Experience

## **7 DRUG INTERACTIONS**

7.1 Effects of Other Drugs on Combined Oral Contraceptives

7.2 Effects of Combined Oral Contraceptives on Other Drugs

7.3 Interference with Laboratory Tests

7.4 Concomitant Use with HCV Combination Therapy - Liver Enzyme Elevation

## **8 USE IN SPECIFIC POPULATIONS**

8.1 Pregnancy

8.3 Nursing Mothers

8.4 Pediatric Use

8.5 Geriatric Use

8.6 Hepatic Impairment

8.7 Renal Impairment

## **10 OVERDOSAGE**

## **11 DESCRIPTION**

## **12 CLINICAL PHARMACOLOGY**

12.1 Mechanism of Action

12.2 Pharmacodynamics

12.3 Pharmacokinetics

## **13 NONCLINICAL TOXICOLOGY**

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

## **14 CLINICAL STUDIES**

14.1 Contraception

14.2 Acne

## **16 HOW SUPPLIED/STORAGE AND HANDLING**

16.1 How Supplied

16.2 Storage Conditions

## **17 PATIENT COUNSELING INFORMATION**

\* Sections or subsections omitted from the full prescribing information are not listed.

---

## **FULL PRESCRIBING INFORMATION**

### **WARNING: CIGARETTE SMOKING AND SERIOUS CARDIOVASCULAR EVENTS**

**Cigarette smoking increases the risk of serious cardiovascular events from combination oral contraceptive (COC) use. This risk increases with age, particularly in women over 35 years of age, and with the number of cigarettes smoked. For this reason, COCs are contraindicated in women who are over 35 years of age and smoke [see Contraindications (4)].**

## **1 INDICATIONS AND USAGE**

### **1.1 Oral Contraceptive**

Previfem<sup>®</sup> (norgestimate/ethinyl estradiol tablets) and Tri-Previfem<sup>®</sup> (norgestimate/ethinyl estradiol tablets) are indicated for use by females of reproductive potential to prevent pregnancy [see *Clinical Studies (14)*].

## 1.2 Acne

Tri-Previfem<sup>®</sup> (norgestimate/ethinyl estradiol tablets) is indicated for the treatment of moderate acne vulgaris in females at least 15 years of age, who have no known contraindications to oral contraceptive therapy and have achieved menarche. Previfem<sup>®</sup> (norgestimate/ethinyl estradiol tablets) should be used for the treatment of acne only if the patient desires an oral contraceptive for birth control [see *Clinical Studies (14)*].

## 2 DOSAGE AND ADMINISTRATION

### 2.1 How to Start Previfem<sup>®</sup> (norgestimate/ethinyl estradiol tablets) or Tri-Previfem<sup>®</sup> (norgestimate/ethinyl estradiol tablets)

Previfem<sup>®</sup> (norgestimate/ethinyl estradiol tablets) and Tri-Previfem<sup>®</sup> (norgestimate/ethinyl estradiol tablets) are dispensed in a blister pack tablet dispenser [see *How Supplied/Storage and Handling (16)*].

Previfem<sup>®</sup> (norgestimate/ethinyl estradiol tablets) and Tri-Previfem<sup>®</sup> (norgestimate/ethinyl estradiol tablets) may be started using either a Day 1 start or a Sunday start (see Table 1). For the first cycle of a Sunday Start regimen, an additional method of contraception should be used until after the first 7 consecutive days of administration.

### 2.2 How to Take Previfem<sup>®</sup> or Tri-Previfem<sup>®</sup>

**Table 1: Instructions for Administration of Previfem<sup>®</sup> or Tri-Previfem<sup>®</sup>**

<p><b>Starting COCs in women not currently using hormonal contraception (Day 1 Start or Sunday Start)</b></p> <p><b>Important:</b> Consider the possibility of ovulation and conception prior to initiation of this product.</p> <p><b>Tablet Color:</b></p> <ul style="list-style-type: none"> <li>• Previfem<sup>®</sup> active tablets are blue (Day 1 to Day 21).</li> <li>• Tri-Previfem<sup>®</sup> active tablets are white (Day 1 to Day 7), light blue (Day 8 to Day 14) and blue (Day 15 to Day 21).</li> <li>• Previfem<sup>®</sup> and Tri-Previfem<sup>®</sup> both have light-green inactive tablets (Day 22 to Day 28).</li> </ul>	<p><b>Day 1 Start:</b></p> <ul style="list-style-type: none"> <li>• Take first active tablet without regard to meals on the first day of menses.</li> <li>• Take subsequent active tablets once daily at the same time each day for a total of 21 days.</li> <li>• Take one <b>light-green</b> inactive tablet daily for 7 days and at the same time of day that active tablets were taken.</li> <li>• Begin each subsequent pack on the same day of the week as the first cycle pack (i.e., on the day after taking the last inactive tablet)</li> </ul> <p><b>Sunday Start:</b></p> <ul style="list-style-type: none"> <li>• Take first active tablet without regard to meals on the first Sunday after the onset of menses. <b>Due to the potential risk of becoming pregnant, use additional non-hormonal contraception (such as condoms and spermicide) for the first seven days of the patient's first cycle pack of Previfem<sup>®</sup> or Tri-Previfem<sup>®</sup>.</b></li> <li>• Take subsequent active tablets once daily at the same time each day for a total of 21 days.</li> <li>• Take one light-green inactive tablet daily for the following 7 days and at the same time of day that active tablets were taken.</li> <li>• Begin each subsequent pack on the same day of the week as the first cycle pack (i.e., on the Sunday after taking the last inactive tablet) and additional non-hormonal contraceptive is not needed.</li> </ul>
<p><b>Switching to Previfem<sup>®</sup> or Tri-Previfem<sup>®</sup> from</b></p>	<p>Start on the same day that a new pack of the previous oral contraceptive</p>

another oral contraceptive	would have started.
<b>Switching from another contraceptive method to Prevfem<sup>®</sup> or Tri-Previfem<sup>®</sup></b>	<b>Start Prevfem<sup>®</sup> or Tri-Previfem<sup>®</sup> :</b>
• <b>Transdermal patch</b>	• On the day when next application would have been scheduled
• <b>Vaginal ring</b>	• On the day when next insertion would have been scheduled
• <b>Injection</b>	• On the day when next injection would have been scheduled
• <b>Intrauterine contraceptive</b>	• On the day of removal • If the IUD is not removed on first day of the patient's menstrual cycle, additional non-hormonal contraceptive (such as condoms and spermicide) is needed for the first seven days of the first cycle pack.
• <b>Implant</b>	• On the day of removal
<b>Complete instructions to facilitate patient counseling on proper tablet usage are located in the FDA-Approved Patient Labeling.</b>	

#### Starting Prevfem<sup>®</sup> and Tri-Previfem<sup>®</sup> after Abortion or Miscarriage

##### *First-trimester*

- After a first-trimester abortion or miscarriage, Prevfem<sup>®</sup> or Tri-Previfem<sup>®</sup> may be started immediately. An additional method of contraception is not needed if Prevfem<sup>®</sup> or Tri-Previfem<sup>®</sup> is started immediately.
- If Prevfem<sup>®</sup> or Tri-Previfem<sup>®</sup> is not started within 5 days after termination of the pregnancy, the patient should use additional non-hormonal contraception (such as condoms and spermicide) for the first seven days of her first cycle pack of Prevfem<sup>®</sup> or Tri-Previfem<sup>®</sup>.

##### *Second-trimester*


- Do not start until 4 weeks after a second-trimester abortion or miscarriage, due to the increased risk of thromboembolic disease. Start Prevfem<sup>®</sup> or Tri-Previfem<sup>®</sup>, following the instructions in Table 1 for Day 1 or Sunday start, as desired. If using Sunday start, use additional non-hormonal contraception (such as condoms and spermicide) for the first seven days of the patient's first cycle pack of Prevfem<sup>®</sup> or Tri-Previfem<sup>®</sup> [see *Contraindications (4)*, *Warnings and Precautions (5.1)*, and *FDA-Approved Patient Labeling*].

#### Starting Prevfem<sup>®</sup> or Tri-Previfem<sup>®</sup> after Childbirth

- Do not start until 4 weeks after delivery, due to the increased risk of thromboembolic disease. Start contraceptive therapy with Prevfem<sup>®</sup> or Tri-Previfem<sup>®</sup> following the instructions in Table 1 for women not currently using hormonal contraception.
- Prevfem<sup>®</sup> or Tri-Previfem<sup>®</sup> are not recommended for use in lactating women [see *Use in Specific Populations (8.3)*].
- If the woman has not yet had a period postpartum, consider the possibility of ovulation and conception occurring prior to use of Prevfem<sup>®</sup> or Tri-Previfem<sup>®</sup> [see *Contraindications (4)*, *Warnings and Precautions (5.1)*, *Use in Specific Populations (8.1 and 8.3)*, and *FDA-Approved Patient Labeling*].

**Prevfem<sup>®</sup> and Tri-Previfem<sup>®</sup> come in a blister pack pill dispenser. Read the instructions below for using the blister pack pill dispenser.**

The blister package consists of three parts, the calendar label, the sleeve and the blister pack containing 28 individually sealed pills. Note that the pills are arranged in four numbered rows of 7 pills, with the pre-printed days of the week printed above them. Refer to the sample of the blister pack below:



Previfem<sup>®</sup> consists of 21 blue “active” birth control pills and 7 light green “reminder” pills.

Tri-Previfem<sup>®</sup> consists of 7 white “active” pills, 7 light-blue “active” pills, 7 blue “active” pills and 7 light green “reminder” pills.

There are two ways to start taking birth-control pills, Sunday Start or Day 1 Start.

### How to use Blister Cards for the 28 tablets

1. If Sunday Start, the patient discards the stickers and takes the first active pill on the first Sunday after their menstrual period begins. Due to the potential risk of becoming pregnant, use additional non-hormonal contraception (such as condoms and spermicide) for the first seven days of the patient’s first cycle pack of Previfem<sup>®</sup> or Tri-Previfem<sup>®</sup>.
2. If Day 1 Start, the patient picks the Days of the Week Sticker that starts the first day of their period. When the patient has picked the right sticker, they need to throw away the others and place the sticker on the blister card over the preprinted days of the week and make sure it lines up with the pills.
3. The patient removes the first pill by pushing down on the pill and waits 24 hours to take their next pill. The patient continues to take one pill each day until all the pills have been taken.
4. The pill should be taken at the same time each day.
5. After taking the last pill, the patient starts a new blister pack the very next day, no matter when their next period starts.
6. The patient should take the pills in each new package as before and start with the pill on the first row and take one pill each day, left to right, until the last pill has been taken.

### 2.3 Missed Tablets

**Table 2: Instructions for Missed Previfem<sup>®</sup> or Tri-Previfem<sup>®</sup> Tablets**

<ul style="list-style-type: none"> <li>• If one active tablet is missed in Weeks 1, 2, or 3</li> </ul>	Take the tablet as soon as possible. Continue taking one tablet a day until the pack is finished.
	Take the two missed tablets as soon as possible and the next two active

<ul style="list-style-type: none"> <li>• If two active tablets are missed in Week 1 or Week 2</li> </ul>	<p>tablets the next day. Continue taking one tablet a day until the pack is finished. <b>Additional non-hormonal contraception (such as condoms and spermicide) should be used as back-up if the patient has sex within 7 days after missing tablets.</b></p>
<ul style="list-style-type: none"> <li>• If two active tablets are missed in the third week or three or more active tablets are missed in a row in Weeks 1, 2, or 3</li> </ul>	<p><u>Day 1 start:</u> Throw out the rest of the pack and start a new pack that same day.  <u>Sunday start:</u> Continue taking one tablet a day until Sunday, then throw out the rest of the pack and start a new pack that same day. <b>Additional non-hormonal contraception (such as condoms and spermicide) should be used as back-up if the patient has sex within 7 days after missing tablets.</b></p>

## 2.4 Advice in Case of Gastrointestinal Disturbances

In case of severe vomiting or diarrhea, absorption may not be complete and additional contraceptive measures should be taken. If vomiting or diarrhea occurs within 3 to 4 hours after taking an active tablet, handle this as a missed tablet [see FDA-Approved Patient Labeling].

## 2.5 Tri-Previfem<sup>®</sup> Use for Acne

The timing of initiation of dosing with Tri-Previfem<sup>®</sup> for acne should follow the guidelines for use of Tri-Previfem<sup>®</sup> as an oral contraceptive. Consult the DOSAGE AND ADMINISTRATION section (2.1) for instructions.

## 3 DOSAGE FORMS AND STRENGTHS

Previfem<sup>®</sup>:

Previfem<sup>®</sup> (norgestimate and ethinyl estradiol tablets USP) is packaged in cartons of 6 blister pack tablet dispensers containing 28 tablets as follows:

21 blue tablets containing 0.25 mg of norgestimate and 0.035 mg of ethinyl estradiol which are round, unscored, film-coated tablets debossed with “93” and “748” on each side.

7 light-green, round, film-coated tablets debossed with “93” and “743” containing inert ingredients.

Tri-Previfem<sup>®</sup>:

Tri-Previfem<sup>®</sup> (norgestimate and ethinyl estradiol tablets USP) is packaged in cartons of 6 blister pack tablet dispensers, each blister pack tablet dispenser contains 28 tablets as follows:

Each white tablet contains 0.18 mg norgestimate and 0.035 mg of ethinyl estradiol.

Each light-blue tablet contains 0.215 mg of norgestimate and 0.035 mg ethinyl estradiol.

Each blue tablet contains 0.25 mg of, norgestimate and 0.035 mg of ethinyl estradiol.

Each light-green tablet contains inert ingredients.

The white tablets are round, unscored film-coated, imprinted with “93” on one side and “746” on the other side; the light-blue tablets are round, unscored film-coated, imprinted with “93” on one side and “747” on the other side; the blue tablets are round, unscored film-coated, imprinted with “93” on one side and “748” on the other side; the light-green tablets are round, film-coated, imprinted with “93” on one side and “743” on the other side.

## 4 CONTRAINDICATIONS

Do not prescribe Prevfem<sup>®</sup> or Tri-Previfem<sup>®</sup> to women who are known to have the following conditions:

- A high risk of arterial or venous thrombotic diseases. Examples include women who are known to:
  - Smoke, if over age 35 [*see Boxed Warning and Warnings and Precautions (5.1)*]
  - Have deep vein thrombosis or pulmonary embolism, now or in the past [*see Warnings and Precautions (5.1)*]
  - Have inherited or acquired hypercoagulopathies [*see Warnings and Precautions (5.1)*]
  - Have cerebrovascular disease [*see Warnings and Precautions (5.1)*]
  - Have coronary artery disease [*see Warnings and Precautions (5.1)*]
  - Have thrombogenic valvular or thrombogenic rhythm diseases of the heart (for example, subacute bacterial endocarditis with valvular disease, or atrial fibrillation) [*see Warnings and Precautions (5.1)*]
  - Have uncontrolled hypertension [*see Warnings and Precautions (5.4)*]
  - Have diabetes mellitus with vascular disease [*see Warnings and Precautions (5.6)*]
  - Have headaches with focal neurological symptoms or migraine headaches with aura [*see Warnings and Precautions (5.7)*]
    - Women over age 35 with any migraine headaches [*see Warnings and Precautions (5.7)*]
- Liver tumors, benign or malignant, or liver disease [*see Warnings and Precautions (5.2)*]
- Undiagnosed abnormal uterine bleeding [*see Warnings and Precautions (5.8)*]
- Pregnancy, because there is no reason to use COCs during pregnancy [*see Warnings and Precautions (5.9) and Use in Specific Populations (8.1)*]
- Breast cancer or other estrogen- or progestin-sensitive cancer, now or in the past [*see Warnings and Precautions (5.11)*]
- Use of Hepatitis C drug combinations containing ombitasvir/paritaprevir/ritonavir, with or without dasabuvir, due to the potential for ALT elevations [*see Warnings and Precautions (5.3)*]

## 5 WARNINGS AND PRECAUTIONS

### 5.1 Thromboembolic Disorders and Other Vascular Problems

- Stop Prevfem<sup>®</sup> or Tri-Previfem<sup>®</sup> if an arterial thrombotic event or venous thromboembolic (VTE) event occurs.
- Stop Prevfem<sup>®</sup> or Tri-Previfem<sup>®</sup> if there is unexplained loss of vision, proptosis, diplopia, papilledema, or retinal vascular lesions. Evaluate for retinal vein thrombosis immediately [*see Adverse Reactions (6.2)*].
- If feasible, stop Prevfem<sup>®</sup> or Tri-Previfem<sup>®</sup> at least 4 weeks before and through 2 weeks after major surgery or other surgeries known to have an elevated risk of VTE as well as during and following prolonged immobilization.
- Start Prevfem<sup>®</sup> or Tri-Previfem<sup>®</sup> no earlier than 4 weeks after delivery, in women who are not breastfeeding. The risk of postpartum VTE decreases after the third postpartum week, whereas the risk of ovulation increases after the third postpartum week.
- The use of COCs increases the risk of VTE. However, pregnancy increases the risk of VTE as much or more than the use of COCs. The risk of VTE in women using COCs is 3 to 9 cases per 10,000 woman-years. The risk of VTE is highest during the first year of use of COCs and when restarting hormonal contraception after a break of 4 weeks or longer. The risk of thromboembolic disease due to COCs gradually disappears after use is discontinued.
- Use of COCs also increases the risk of arterial thromboses such as strokes and myocardial infarctions, especially in women with other risk factors for these events. COCs have been shown to increase both the relative and attributable risks of cerebrovascular events (thrombotic and hemorrhagic strokes). This risk increases with age, particularly in women over 35 years of age who smoke.
- Use COCs with caution in women with cardiovascular disease risk factors.



## 5.2 Liver Disease

### Impaired Liver Function

Do not use Prevfem<sup>®</sup> or Tri-Previfem<sup>®</sup> in women with liver disease, such as acute viral hepatitis or severe (decompensated) cirrhosis of liver [see *Contraindications (4)*]. Acute or chronic disturbances of liver function may necessitate the discontinuation of COC use until markers of liver function return to normal and COC causation has been excluded. Discontinue Prevfem<sup>®</sup> or Tri-Previfem<sup>®</sup> if jaundice develops.

### Liver Tumors

Prevfem<sup>®</sup> and Tri-Previfem<sup>®</sup> are contraindicated in women with benign and malignant liver tumors [see *Contraindications (4)*]. Hepatic adenomas are associated with COC use. An estimate of the attributable risk is 3.3 cases/100,000 COC users. Rupture of hepatic adenomas may cause death through intra-abdominal hemorrhage.

Studies have shown an increased risk of developing hepatocellular carcinoma in long-term (>8 years) COC users. However, the risk of liver cancers in COC users is less than one case per million users.

## 5.3 Risk of Liver Enzyme Elevations with Concomitant Hepatitis C Treatment

During clinical trials with the Hepatitis C combination drug regimen that contains ombitasvir/paritaprevir/ritonavir, with or without dasabuvir, ALT elevations greater than 5 times the upper limit of normal (ULN), including some cases greater than 20 times the ULN, were significantly more frequent in women using ethinyl estradiol-containing medications, such as COCs. Discontinue Prevfem<sup>®</sup> or Tri-Previfem<sup>®</sup> prior to starting therapy with the combination drug regimen ombitasvir/paritaprevir/ritonavir, with or without dasabuvir [see *Contraindications (4)*]. Prevfem<sup>®</sup> or Tri-Previfem<sup>®</sup> can be restarted approximately 2 weeks following completion of treatment with the Hepatitis C combination drug regimen.

## 5.4 High Blood Pressure

Prevfem<sup>®</sup> and Tri-Previfem<sup>®</sup> are contraindicated in women with uncontrolled hypertension or hypertension with vascular disease [see *Contraindications (4)*]. For women with well-controlled hypertension, monitor blood pressure and stop Prevfem<sup>®</sup> and Tri-Previfem<sup>®</sup> if blood pressure rises significantly.

An increase in blood pressure has been reported in women taking COCs, and this increase is more likely in older women with extended duration of use. The incidence of hypertension increases with increasing concentrations of progestin.

## 5.5 Gallbladder Disease

Studies suggest a small increased relative risk of developing gallbladder disease among COC users. Use of COCs may worsen existing gallbladder disease. A past history of COC-related cholestasis predicts an increased risk with subsequent COC use. Women with a history of pregnancy-related cholestasis may be at an increased risk for COC related cholestasis.

## 5.6 Carbohydrate and Lipid Metabolic Effects

Carefully monitor prediabetic and diabetic women who take Prevfem<sup>®</sup> or Tri-Previfem<sup>®</sup>. COCs may decrease glucose tolerance.

Consider alternative contraception for women with uncontrolled dyslipidemia. A small proportion of women will have adverse lipid changes while on COCs.

Women with hypertriglyceridemia, or a family history thereof, may be at an increased risk of pancreatitis when using COCs.

## 5.7 Headache

If a woman taking Prevfem<sup>®</sup> or Tri-Previfem<sup>®</sup> develops new headaches that are recurrent, persistent, or severe, evaluate the cause and discontinue Prevfem<sup>®</sup> or Tri-Previfem<sup>®</sup> if indicated.

Consider discontinuation of Prevfem<sup>®</sup> or Tri-Previfem<sup>®</sup> in the case of increased frequency or severity of migraine during COC use (which may be prodromal of a cerebrovascular event).

## 5.8 Bleeding Irregularities and Amenorrhea

### Unscheduled Bleeding and Spotting

Unscheduled (breakthrough or intracyclic) bleeding and spotting sometimes occur in patients on COCs, especially during the first three months of use. If bleeding persists or occurs after previously regular cycles, check for causes such as pregnancy or malignancy. If pathology and pregnancy are excluded, bleeding irregularities may resolve over time or with a change to a different contraceptive product.

In clinical trials of norgestimate and ethinyl estradiol tablets 0.25 mg/0.035 mg and norgestimate and ethinyl estradiol tablets 0.18 mg/0.035 mg, 0.215 mg/0.035 mg, 0.25 mg/0.035 mg, the frequency and duration of breakthrough bleeding and/or spotting was assessed in 1,647 patients (21,275 evaluable cycles) and 4,826 patients (35,546 evaluable cycles), respectively. A total of 100 (7.5%) women discontinued norgestimate and ethinyl estradiol tablets 0.25 mg/0.035 mg and 231 (4.8%) women discontinued norgestimate and ethinyl estradiol tablets 0.18 mg/0.035 mg, 0.215 mg/0.035 mg, 0.25 mg/0.035 mg, at least in part, due to bleeding or spotting. Based on data from the clinical trials, 14-34% of women using norgestimate and ethinyl estradiol tablets 0.25 mg/0.035 mg experienced unscheduled bleeding per cycle in the first year; for norgestimate and ethinyl estradiol tablets 0.18 mg/0.035 mg, 0.215 mg/0.035 mg, 0.25 mg/0.035 mg, the respective numbers were 13-38%. The percent of women who experienced breakthrough/unscheduled bleeding tended to decrease over time.

### Amenorrhea and Oligomenorrhea

Women who use Prevfem<sup>®</sup> or Tri-Previfem<sup>®</sup> may experience amenorrhea. Some women may experience amenorrhea or oligomenorrhea after discontinuation of COCs, especially when such a condition was pre-existent.

If scheduled (withdrawal) bleeding does not occur, consider the possibility of pregnancy. If the patient has not adhered to the prescribed dosing schedule (missed one or more active tablets or started taking them on a day later than she should have), consider the possibility of pregnancy at the time of the first missed period and take appropriate diagnostic measures. If the patient has adhered to the prescribed regimen and misses two consecutive periods, rule out pregnancy.

## 5.9 COC Use Before or During Early Pregnancy

Extensive epidemiological studies have revealed no increased risk of birth defects in women who have used oral contraceptives prior to pregnancy. Studies also do not suggest a teratogenic effect, particularly in so far as cardiac anomalies and limb reduction defects are concerned, when oral contraceptives are taken inadvertently during early pregnancy. Discontinue Prevfem<sup>®</sup> or Tri-Previfem<sup>®</sup> use if pregnancy is confirmed.

Administration of COCs to induce withdrawal bleeding should not be used as a test for pregnancy [*see Use in Specific Populations (8.1)*].

## 5.10 Depression

Carefully observe women with a history of depression and discontinue Prevfem<sup>®</sup> or Tri-Previfem<sup>®</sup> if depression recurs to a serious degree.

## 5.11 Carcinoma of Breast and Cervix

- Prevfem<sup>®</sup> and Tri-Previfem<sup>®</sup> are contraindicated in women who currently have or have had breast cancer because breast cancer may be hormonally sensitive [*see Contraindications (4)*].

There is substantial evidence that COCs do not increase the incidence of breast cancer. Although

some past studies have suggested that COCs might increase the incidence of breast cancer, more recent studies have not confirmed such findings.

- Some studies suggest that COC use has been associated with an increase in the risk of cervical cancer or intraepithelial neoplasia. However, there continues to be controversy about the extent to which such findings may be due to differences in sexual behavior and other factors.

### **5.12 Effect on Binding Globulins**

The estrogen component of COCs may raise the serum concentrations of thyroxine-binding globulin, sex hormone-binding globulin, and cortisol-binding globulin. The dose of replacement thyroid hormone or cortisol therapy may need to be increased.

### **5.13 Monitoring**

A woman who is taking COCs should have a yearly visit with her healthcare provider for a blood pressure check and for other indicated healthcare.

### **5.14 Hereditary Angioedema**

In women with hereditary angioedema, exogenous estrogens may induce or exacerbate symptoms of angioedema.

### **5.15 Chloasma**

Chloasma may occasionally occur, especially in women with a history of chloasma gravidarum. Women with a tendency to chloasma should avoid exposure to the sun or ultraviolet radiation while taking Tri-Previfem<sup>®</sup> or Previfem<sup>®</sup>.

## **6 ADVERSE REACTIONS**

The following serious adverse reactions with the use of COCs are discussed elsewhere in labeling:

- Serious cardiovascular events and stroke [*see Boxed Warning and Warnings and Precautions (5.1)*]
- Vascular events [*see Warnings and Precautions (5.1)*]
- Liver disease [*see Warnings and Precautions (5.2)*]

Adverse reactions commonly reported by COC users are:

- Irregular uterine bleeding
- Nausea
- Breast tenderness
- Headache

### **6.1 Clinical Trial Experience**

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in clinical practice.

Norgestimate and Ethinyl Estradiol Tablets 0.25 mg/0.035 mg:

The safety of norgestimate and ethinyl estradiol tablets 0.25 mg/0.035 mg was evaluated in 1,647 healthy women of child-bearing potential who participated in 3 clinical trials and received at least 1 dose of norgestimate and ethinyl estradiol tablets 0.25 mg/0.035 mg for contraception. Two trials were randomized active-controlled trials and 1 was an uncontrolled open-label trial. In all 3 trials, subjects were followed for up to 24 cycles.

Common Adverse Reactions (≥ 2% of subjects): The most common adverse reactions reported by at least 2% of the 1,647 women were the following in order of decreasing incidence: headache/migraine (32.9%), abdominal/gastrointestinal pain (7.8%), vaginal infection (8.4%), genital discharge (6.8%),

breast issues (including breast pain, discharge, and enlargement) (6.3%), mood disorders (including depression and mood altered) (5.0%), flatulence (3.2%), nervousness (2.9%), and rash (2.6%).

Adverse Reactions Leading to Study Discontinuation: Over the three trials, between 11 to 21% of subjects discontinued the trial due to an adverse reaction. The most common adverse reactions ( $\geq 1\%$ ) leading to discontinuation were: metrorrhagia (6.9%), nausea/vomiting (5.0%), headache (4.1%), mood disorders (including depression and mood altered) (2.4%), premenstrual syndrome (1.7%), hypertension (1.4%), breast pain (1.4%), nervousness (1.3%), amenorrhea (1.1%), dysmenorrhea (1.1%), weight increased (1.1%), and flatulence (1.1%).

Serious Adverse Reactions: breast cancer (1 subject), mood disorders including depression, irritability, and mood swings (1 subject), myocardial infarction (1 subject), and venous thromboembolic events including pulmonary embolism (1 subject) and deep vein thrombosis (DVT) (1 subject).

Norgestimate and Ethinyl Estradiol Tablets 0.18 mg/0.035 mg, 0.215 mg/0.035 mg, 0.25 mg/0.035 mg:

The safety of norgestimate and ethinyl estradiol tablets 0.18 mg/0.035 mg, 0.215 mg/0.035 mg, 0.25 mg/0.035 mg was evaluated in 4,826 healthy women of child-bearing potential who participated in 6 clinical trials and received at least 1 dose of norgestimate and ethinyl estradiol tablets 0.18 mg/0.035 mg, 0.215 mg/0.035 mg, 0.25 mg/0.035 mg for contraception. Two trials were randomized active-controlled trials and 4 were uncontrolled open-label trials. In 3 trials, subjects were followed for up to 24 cycles; in 2 trials, subjects were followed for up to 12 cycles; and in 1 trial, subjects were followed for up to 6 cycles.

Common Adverse Reactions ( $\geq 2\%$  of subjects): The most common adverse reactions reported by at least 2% of the 4,826 women were the following in order of decreasing incidence: headache/migraine (33.6%), breast issues (including breast pain, enlargement, and discharge) (8.0%), vaginal infection (7.1%), abdominal/gastrointestinal pain (5.6%), mood disorders (including mood alteration and depression) (3.8%), genital discharge (3.2%), and changes in weight (including weight fluctuation, increased or decreased) (2.5%).

Adverse Reactions Leading to Study Discontinuation: Over the trials, between 9 to 27% of subjects discontinued the trial due to an adverse reaction. The most common adverse reactions ( $\geq 1\%$ ) leading to discontinuation were: metrorrhagia (4.3%), nausea/vomiting (2.8%), headache/migraine (2.4%), mood disorders (including depression and mood altered) (1.1%), and weight increased (1.1%).

Serious Adverse Reactions: breast cancer (1 subject), carcinoma of the cervix *in situ* (1 subject), hypertension (1 subject), and migraine (2 subjects).

## 6.2 Postmarketing Experience

The following additional adverse drug reactions have been reported from worldwide postmarketing experience with norgestimate/ethinyl estradiol. Because these reactions are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure.

*Infections and Infestations:* Urinary tract infection;

*Neoplasms Benign, Malignant and Unspecified (Incl. Cysts and Polyps):* Breast cancer, benign breast neoplasm, hepatic adenoma, focal nodular hyperplasia, breast cyst;

*Immune System Disorders:* Hypersensitivity; *Metabolism and Nutrition Disorders:* Dyslipidemia; *Psychiatric Disorders:* Anxiety, insomnia;

*Nervous System Disorders:* Syncope, convulsion, paresthesia, dizziness; *Eye Disorders:* Visual impairment, dry eye, contact lens intolerance; *Ear and Labyrinth Disorders:* Vertigo;

*Cardiac Disorders:* Tachycardia, palpitations;

*Vascular Events:* Deep vein thrombosis, pulmonary embolism, retinal vascular thrombosis, hot flush;

*Arterial Events:* Arterial thromboembolism, myocardial infarction, cerebrovascular accident;

*Respiratory, Thoracic and Mediastinal Disorders:* Dyspnea;

*Gastrointestinal Disorders:* Pancreatitis, abdominal distension, diarrhea, constipation;

*Hepatobiliary Disorders:* Hepatitis;

*Skin and Subcutaneous Tissue Disorders:* Angioedema, erythema nodosum, hirsutism, night sweats, hyperhidrosis, photosensitivity reaction, urticaria, pruritus, acne;

*Musculoskeletal, Connective Tissue, and Bone Disorders:* Muscle spasms, pain in extremity, myalgia, back pain;

*Reproductive System and Breast Disorders:* Ovarian cyst, suppressed lactation, vulvovaginal dryness;

*General Disorders and Administration Site Conditions:* Chest pain, asthenic conditions.

## **7 DRUG INTERACTIONS**

Consult the labeling of concurrently used drugs to obtain further information about interactions with hormonal contraceptives or the potential for enzyme alterations.

No drug-drug interaction studies were conducted with norgestimate and ethinyl estradiol tablets.

### **7.1 Effects of Other Drugs on Combined Oral Contraceptives**

Substances decreasing the plasma concentrations of COCs:

Drugs or herbal products that induce certain enzymes, including cytochrome P450 3A4 (CYP3A4), may decrease the plasma concentrations of COCs and potentially diminish the effectiveness of COCs or increase breakthrough bleeding. Some drugs or herbal products that may decrease the effectiveness of hormonal contraceptives include phenytoin, barbiturates, carbamazepine, bosentan, felbamate, griseofulvin, oxcarbazepine, rifampicin, topiramate, rifabutin, rufinamide, aprepitant, and products containing St. John's wort. Interactions between hormonal contraceptives and other drugs may lead to breakthrough bleeding and/or contraceptive failure. Counsel women to use an alternative method of contraception or a back-up method when enzyme inducers are used with COCs, and to continue back-up contraception for 28 days after discontinuing the enzyme inducer to ensure contraceptive reliability.

Colesevelam: Colesevelam, a bile acid sequestrant, given together with a COC, has been shown to significantly decrease the AUC of EE. The drug interaction between the contraceptive and colesevelam was decreased when the two drug products were given 4 hours apart.

Substances increasing the plasma concentrations of COCs:

Co-administration of atorvastatin or rosuvastatin and certain COCs containing ethinyl estradiol (EE) increase AUC values for EE by approximately 20-25%. Ascorbic acid and acetaminophen may increase plasma EE concentrations, possibly by inhibition of conjugation. CYP3A4 inhibitors such as itraconazole, voriconazole, fluconazole, grapefruit juice, or ketoconazole may increase plasma hormone concentrations.

Human immunodeficiency virus (HIV)/Hepatitis C virus (HCV) protease inhibitors and non-nucleoside reverse transcriptase inhibitors:

Significant changes (increase or decrease) in the plasma concentrations of estrogen and/or progestin have been noted in some cases of co-administration with HIV protease inhibitors (decrease [e.g., nelfinavir, ritonavir, darunavir/ritonavir, (fos)amprenavir/ritonavir, lopinavir/ritonavir, and tipranavir/ritonavir] or increase [e.g., indinavir and atazanavir/ritonavir])/HCV protease inhibitors (decrease [e.g., boceprevir and telaprevir]) or with non-nucleoside reverse transcriptase inhibitors (decrease [e.g., nevirapine] or increase [e.g., etravirine]).

### **7.2 Effects of Combined Oral Contraceptives on Other Drugs**

- COCs containing EE may inhibit the metabolism of other compounds (e.g., cyclosporine,

prednisolone, theophylline, tizanidine, and voriconazole) and increase their plasma concentrations.

- COCs have been shown to decrease plasma concentrations of acetaminophen, clofibric acid, morphine, salicylic acid, temazepam and lamotrigine. Significant decrease in plasma concentration of lamotrigine has been shown, likely due to induction of lamotrigine glucuronidation. This may reduce seizure control; therefore, dosage adjustments of lamotrigine may be necessary.

Women on thyroid hormone replacement therapy may need increased doses of thyroid hormone because the serum concentration of thyroid-binding globulin increases with use of COCs.

### **7.3 Interference with Laboratory Tests**

The use of contraceptive steroids may influence the results of certain laboratory tests, such as coagulation factors, lipids, glucose tolerance, and binding proteins.

### **7.4 Concomitant Use with HCV Combination Therapy - Liver Enzyme Elevation**

Do not co-administer Prevfem<sup>®</sup> or Tri-Previfem<sup>®</sup> with HCV drug combinations containing ombitasvir/paritaprevir/ritonavir, with or without dasabuvir, due to potential for ALT elevations [*see Warnings and Precautions (5.3)*].

## **8 USE IN SPECIFIC POPULATIONS**

### **8.1 Pregnancy**

There is little or no increased risk of birth defects in women who inadvertently use COCs during early pregnancy. Epidemiologic studies and meta-analyses have not found an increased risk of genital or non-genital birth defects (including cardiac anomalies and limb reduction defects) following exposure to low dose COCs prior to conception or during early pregnancy.

Do not administer COCs to induce withdrawal bleeding as a test for pregnancy. Do not use COCs during pregnancy to treat threatened or habitual abortion.

### **8.3 Nursing Mothers**

Advise the nursing mother to use other forms of contraception, when possible, until she has weaned her child. COCs can reduce milk production in breastfeeding mothers. This is less likely to occur once breastfeeding is well-established; however, it can occur at any time in some women. Small amounts of oral contraceptive steroids and/or metabolites are present in breast milk.

### **8.4 Pediatric Use**

Safety and efficacy of norgestimate and ethinyl estradiol tablets have been established in women of reproductive age. Efficacy is expected to be the same for post-pubertal adolescents under the age of 18 and for users 18 years and older. Use of this product before menarche is not indicated.

There was no significant difference between norgestimate and ethinyl estradiol tablets 0.18 mg/0.035 mg, 0.215 mg/0.035 mg, 0.25 mg/0.035 mg and placebo in mean change in total lumbar spine (L1-L4) and total hip bone mineral density between baseline and Cycle 13 in 123 adolescent females with anorexia nervosa in a double-blind, placebo-controlled, multicenter, one-year treatment duration clinical trial for the Intent To Treat (ITT) population.

### **8.5 Geriatric Use**

Norgestimate and Ethinyl Estradiol Tablets have not been studied in postmenopausal women and are not indicated in this population.

### **8.6 Hepatic Impairment**

The pharmacokinetics of norgestimate and ethinyl estradiol tablets has not been studied in subjects with hepatic impairment. However, steroid hormones may be poorly metabolized in patients with hepatic impairment. Acute or chronic disturbances of liver function may necessitate the discontinuation of COC use until markers of liver function return to normal and COC causation has been excluded [see *Contraindications (4) and Warnings and Precautions (5.2)*].

## 8.7 Renal Impairment

The pharmacokinetics of norgestimate and ethinyl estradiol tablets have not been studied in women with renal impairment.

## 10 OVERDOSAGE

There have been no reports of serious ill effects from overdosage of oral contraceptives, including ingestion by children. Overdosage may cause withdrawal bleeding in females and nausea.

## 11 DESCRIPTION

Each of the following products is a combination oral contraceptive containing the progestational compound norgestimate and the estrogenic compound ethinyl estradiol. Norgestimate is designated as (18,19-Dinor-17-pregn-4-en-20-yn-3-one,17-(acetyloxy)-13-ethyl-, oxime,(17 $\alpha$ )-(+) ) and ethinyl estradiol is designated as (19-nor-17 $\alpha$ -pregna,1,3,5(10)-trien-20-yne-3,17-diol).

### Previfem<sup>®</sup>

Each active blue tablet contains 0.25 mg of norgestimate and 0.035 mg of ethinyl estradiol. Inactive ingredients include FD&C Blue No. 1 HT Aluminum Lake, hypromellose, lactose monohydrate, magnesium stearate, polyethylene glycol, and pregelatinized starch.

Each light-green placebo tablet contains only inert ingredients, as follows: FD&C Blue No. 2, hypromellose, iron oxide yellow, lactose monohydrate, magnesium stearate, polyethylene glycol, and pregelatinized starch.


### Tri-Previfem<sup>®</sup>

Each active white tablet contains 0.18 mg of norgestimate and 0.035 mg of ethinyl estradiol. Inactive ingredients include hypromellose, lactose monohydrate, magnesium stearate, polyethylene glycol, and pregelatinized starch.

Each active light-blue tablet contains 0.215 mg of norgestimate and 0.035 mg of ethinyl estradiol. Inactive ingredients include FD&C Blue No. 1 Aluminum Lake, hypromellose, lactose monohydrate, magnesium stearate, polyethylene glycol, and pregelatinized starch.

Each active blue tablet contains 0.25 mg of norgestimate and 0.035 mg of ethinyl estradiol. Inactive ingredients include FD&C Blue No. 1 Aluminum Lake, hypromellose, lactose monohydrate, magnesium stearate, polyethylene glycol, and pregelatinized starch.

Each light-green placebo tablet contains only inert ingredients, as follows: FD&C Blue No. 2, hypromellose, iron oxide yellow, lactose monohydrate, magnesium stearate, polyethylene glycol, and pregelatinized starch.



## 12 CLINICAL PHARMACOLOGY

### 12.1 Mechanism of Action

- Oral Contraception  
COCs lower the risk of becoming pregnant primarily by suppressing ovulation. Other possible mechanisms may include cervical mucus changes that inhibit sperm penetration and endometrial changes that reduce the likelihood of implantation.
- Acne  
Acne is a skin condition with a multifactorial etiology, including androgen stimulation of sebum production. While the combination of ethinyl estradiol and norgestimate increases sex hormone-binding globulin (SHBG) and decreases free testosterone, the relationship between these changes and a decrease in the severity of facial acne in otherwise healthy women with this skin condition has not been established.

### 12.2 Pharmacodynamics

No specific pharmacodynamic studies were conducted with norgestimate and ethinyl estradiol tablets.

### 12.3 Pharmacokinetics

#### *Absorption*

Norgestimate (NGM) and EE are rapidly absorbed following oral administration. NGM is rapidly and completely metabolized by first pass (intestinal and/or hepatic) mechanisms to norelgestromin (NGMN) and norgestrel (NG), which are the major active metabolites of norgestimate.

Peak serum concentrations of NGMN and EE are generally reached by 2 hours after administration of norgestimate and ethinyl estradiol tablets. Accumulation following multiple dosing of the 250 mcg NGM / 35 mcg EE dose is approximately 2-fold for NGMN and EE compared with single dose administration. The pharmacokinetics of NGMN is dose-proportional following NGM doses of 180 mcg to 250 mcg. Steady-state concentration of EE is achieved by Day 7 of each dosing cycle. Steady-state concentrations of NGMN and NG are achieved by Day 21. Non-linear accumulation (approximately 8 fold) of NG is observed as a result of high-affinity binding to SHBG, which limits its biological activity (Table 3).

**Table 3: Summary of NGMN, NG and EE pharmacokinetic parameters.**

<b>Mean (SD) Pharmacokinetic Parameters of Norgestimate and Ethinyl Estradiol Tablets 0.18 mg/0.035 mg, 0.215 mg/0.035 mg, 0.25 mg/0.035 mg During a Three Cycle Study</b>						
Analyte	Cycle	Day	C <sub>max</sub>	t <sub>max</sub> (h)	AUC <sub>0-24h</sub>	t <sub>1/2</sub> (h)



NGMN	3	7	1.80 (0.46)	1.42 (0.73)	15.0 (3.88)	NC
		14	2.12 (0.56)	1.21 (0.26)	16.1 (4.97)	NC
		21	2.66 (0.47)	1.29 (0.26)	21.4 (3.46)	22.3 (6.54)
NG	3	7	1.94 (0.82)	3.15 (4.05)	34.8 (16.5)	NC
		14	3.00 (1.04)	2.21 (2.03)	55.2 (23.5)	NC
		21	3.66 (1.15)	2.58 (2.97)	69.3 (23.8)	40.2 (15.4)
EE	3	7	124 (39.5)	1.27 (0.26)	1130 (420)	NC
		14	128 (38.4)	1.32 (0.25)	1130 (324)	NC
		21	126 (34.7)	1.31 (0.56)	1090 (359)	15.9 (4.39)
<b>Mean (SD) Pharmacokinetic Parameters of Norgestimate and Ethinyl Estradiol Tablets 0.25 mg/0.035 mg During a Three Cycle Study</b>						
Analyte	Cycle	Day	C <sub>max</sub>	t <sub>max</sub> (h)	AUC <sub>0-24h</sub>	t <sub>1/2</sub> (h)
NGMN	1	1	1.78 (0.397)	1.19 (0.250)	9.90 (3.25)	18.4 (5.91)
	3	21	2.19 (0.655)	1.43 (0.680)	18.1 (5.53)	24.9 (9.04)
NG	1	1	0.649 (0.49)	1.42 (0.69)	6.22 (2.46)	37.8 (14.0)
	3	21	2.65 (1.11)	1.67 (1.32)	48.2 (20.5)	45.0 (20.4)
EE	1	1	92.2 (24.5)	1.2 (0.26)	629 (138)	10.1 (1.90)
	3	21	147 (41.5)	1.13 (0.23)	1210 (294)	15.0 (2.36)
C <sub>max</sub> = peak serum concentration, t <sub>max</sub> = time to reach peak serum concentration, AUC <sub>0-24h</sub> = area under serum concentration vs time curve from 0 to 24 hours, t <sub>1/2</sub> = elimination half-life, NC = not calculated.						
NGMN and NG: C <sub>max</sub> = ng/mL, AUC <sub>0-24h</sub> = h·ng/mL						
EE: C <sub>max</sub> = pg/mL, AUC <sub>0-24h</sub> = h·pg/mL						

### Food Effect

The effect of food on the pharmacokinetics of norgestimate and ethinyl estradiol tablets has not been studied.

### Distribution

NGMN and NG are highly bound (>97%) to serum proteins. NGMN is bound to albumin and not to SHBG, while NG is bound primarily to SHBG. EE is extensively bound (>97%) to serum albumin and induces an increase in the serum concentrations of SHBG.

### Metabolism

NGM is extensively metabolized by first-pass mechanisms in the gastrointestinal tract and/or liver. NGM's primary active metabolite is NGMN. Subsequent hepatic metabolism of NGMN occurs and metabolites include NG, which is also active, and various hydroxylated and conjugated metabolites. Although NGMN and its metabolites inhibit a variety of P450 enzymes in human liver microsomes, under the recommended dosing regimen, the *in vivo* concentrations of NGMN and its metabolites, even at the peak serum levels, are relatively low compared to the inhibitory constant (K<sub>i</sub>). EE is also metabolized to various hydroxylated products and their glucuronide and sulfate conjugates.

### Excretion

The metabolites of NGMN and EE are eliminated by renal and fecal pathways. Following administration of <sup>14</sup>C-norgestimate, 47% (45-49%) and 37% (16-49%) of the administered radioactivity was eliminated in the urine and feces, respectively. Unchanged NGM was not detected in the urine. In addition to 17-deacetyl norgestimate, a number of metabolites of NGM have been identified in human urine following administration of radiolabeled NGM. These include 18, 19-Dinor-17-pregn-4-en-20-yn-3-one, 17-hydroxy-13-ethyl, (17 $\alpha$ )-(-); 18, 19-Dinor-5 $\beta$  17-pregnan-20-yn, 3 $\alpha$ , 17 $\beta$ -dihydroxy-13-ethyl, (17 $\alpha$ ), various hydroxylated metabolites and conjugates of these metabolites.

## 13 NONCLINICAL TOXICOLOGY

### 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

[See Warnings and Precautions (5.2, 5.11) and Use in Specific Populations (8.1).]

## 14 CLINICAL STUDIES

### 14.1 Contraception

In three US clinical trials with norgestimate and ethinyl estradiol tablets 0.25 mg/0.035 mg, 1,651 women aged 18 to 38 years were studied for up to 24 cycles, proving a total of 24,272 cycles of exposure. The racial demographic was about 73-86% Caucasian, 8-13% African-American, 6-14% Hispanic with the remainder Asian or Other ( $\leq 1\%$ ). There were no exclusions on the basis of weight; the weight range for women treated was 82-303 lbs, with a mean weight of about 135 lbs. The pregnancy rate was approximately 1 pregnancy per 100 women-years.

In four clinical trials with norgestimate and ethinyl estradiol tablets 0.18 mg/0.035 mg, 0.215 mg/0.035 mg, 0.25 mg/0.035 mg, 4,756 women aged 15 to 41 years were studied for 24 cycles, providing a total of 45,244 cycles of exposure. The racial demographic was about 87-90% Caucasian, 6-10% African-American, with the remainder Asian ( $\leq 1\%$ ) or Other (2-5%). There were no exclusions on the basis of weight; the weight range for women treated was 80-310 lbs, with a mean weight of about 132 lbs. The pregnancy rate was approximately 1 pregnancy per 100 women-years.

### 14.2 Acne

Norgestimate and Ethinyl Estradiol Tablets 0.18 mg/0.035 mg, 0.215 mg/0.035 mg, 0.25 mg/0.035 mg was evaluated for the treatment of acne vulgaris in two randomized, double-blind, placebo-controlled, multicenter, six- (28 day) cycle studies. Two hundred twenty- one patients received norgestimate and ethinyl estradiol tablets 0.18 mg/0.035 mg, 0.215 mg/0.035 mg, 0.25 mg/0.035 mg and 234 patients received placebo. Mean age at enrollment for both groups was 28 years. At the end of 6 months, the mean total lesion count changed from 55 to 31 (42% reduction) in patients treated with norgestimate and ethinyl estradiol tablets 0.18 mg/0.035 mg, 0.215 mg/0.035 mg, 0.25 mg/0.035 mg and from 54 to 38 (27% reduction) in patients similarly treated with placebo. Table 4 summarizes the changes in lesion count for each type of lesion. Based on the investigator's global assessment conducted at the final visit, patients treated with norgestimate and ethinyl estradiol tablets 0.18 mg/0.035 mg, 0.215 mg/0.035 mg, 0.25 mg/0.035 mg showed a statistically significant improvement in total lesions compared to those treated with placebo.

**Table 4: Acne Vulgaris Indication. Combined Results: Two Multicenter, Placebo-Controlled Trials. Observed Means at Six Months (LOCF)<sup>1</sup> and at Baseline. Intent-to-Treat Population.**

# of Lesions	norgestimate and ethinyl estradiol tablets, 0.18 mg/0.035 mg, 0.215 mg/0.035 mg, 0.25 mg/0.035 mg (N=221)		Placebo (N=234)		Difference in Counts between norgestimate and ethinyl estradiol tablets, 0.18 mg/0.035 mg, 0.215 mg/0.035 mg, 0.25 mg/0.035 mg and Placebo at 6 Months
	Counts	% Reduction	Counts	% Reduction	
<b>INFLAMMATORY LESIONS</b>					
Baseline Mean	19		19		
Sixth Month Mean	10	48%	13	30%	3 (95% CI: -1.2, 5.1)

NON-  
INFLAMMATORY  
LESIONS

Baseline Mean	36		35		
Sixth Month Mean	22	34%	25	21%	3 (95% CI: -0.2, 7.8)
TOTAL LESIONS					
Baseline Mean	55		54		
Sixth Month Mean	31	42%	38	27%	7 (95% CI: 2.0, 11.9)

<sup>1</sup>LOCF: Last Observation Carried Forward

## 16 HOW SUPPLIED/STORAGE AND HANDLING

### 16.1 How Supplied

#### Previfem<sup>®</sup>

Previfem<sup>®</sup> (norgestimate and ethinyl estradiol tablets USP) is packaged in cartons of 6 blister pack tablet dispensers containing 28 tablets as follows: 21 blue tablets containing 0.25 mg of norgestimate and 0.035 mg of ethinyl estradiol which are round, unscored, film-coated tablets debossed with “93” and “748” on each side and 7 light-green, round, film-coated tablets debossed with “93” and “743” containing inert ingredients.

Blister pack tablet dispenser NDC 0603-7642-01.

Boxes of 6 blister pack tablet dispensers NDC 0603-7642-17.

#### Tri-Previfem<sup>®</sup>

Tri-Previfem<sup>®</sup> (norgestimate and ethinyl estradiol tablets USP) is packaged in cartons of 6 blister pack tablet dispensers, each blister pack tablet dispenser contains 28 tablets as follows:

Each white tablet contains 0.18 mg of norgestimate and 0.035 mg of ethinyl estradiol. Each light-blue tablet contains 0.215 mg of norgestimate and 0.035 mg of ethinyl estradiol. Each blue tablet contains 0.25 mg of norgestimate and 0.035 mg of ethinyl estradiol. Each light-green tablet contains inert ingredients.

The white tablets are round, unscored film-coated, imprinted with “93” on one side and “746” on the other side; the light-blue tablets are round, unscored film-coated, imprinted with “93” on one side and “747” on the other side; the blue tablets are round, unscored film-coated, imprinted with “93” on one side and “748” on the other side; the light-green tablets are round, film-coated, imprinted with “93” on one side and “743” on the other side.

Blister pack tablet dispenser NDC 0603-7663-01.

Boxes of 6 blister pack tablet dispensers NDC 0603-7663-17.

### 16.2 Storage Conditions

- Store at 20° to 25°C (68° to 77°F) [See USP Controlled Room Temperature].
- Protect from light.
- Keep out of reach of children.

## 17 PATIENT COUNSELING INFORMATION

See FDA-approved patient labeling (Patient Information and Instructions for Use).

Counsel patients about the following information:

- Cigarette smoking increases the risk of serious cardiovascular events from COC use, and that women who are over 35 years old and smoke should not use COCs [see *Boxed Warning*].
- Increased risk of VTE compared to non-users of COCs is greatest after initially starting a COC or restarting (following a 4-week or greater pill-free interval) the same or a different COC [see *Warnings and Precautions (5.1)*].
- Prevfem<sup>®</sup> and Tri-Previfem<sup>®</sup> do not protect against HIV infection (AIDS) and other sexually transmitted infections.
- Prevfem<sup>®</sup> and Tri-Previfem<sup>®</sup> are not to be used during pregnancy; if pregnancy occurs during use of Prevfem<sup>®</sup> or Tri-Previfem<sup>®</sup> instruct the patient to stop further use [see *Warnings and Precautions (5.9)*].
- Take one tablet daily by mouth at the same time every day. Instruct patients what to do in the event tablets are missed [see *Dosage and Administration (2.2)*].
- Use a back-up or alternative method of contraception when enzyme inducers are used with Prevfem<sup>®</sup> or Tri-Previfem<sup>®</sup> [see *Drug Interactions (7.1)*].
- COCs may reduce breast milk production; this is less likely to occur if breastfeeding is well established [see *Use in Specific Populations (8.3)*].
- Women who start COCs postpartum, and who have not yet had a period, should use an additional method of contraception until they have taken an active tablet for 7 consecutive days [see *Dosage and Administration (2.2)*].
- Amenorrhea may occur. Consider pregnancy in the event of amenorrhea at the time of the first missed period. Rule out pregnancy in the event of amenorrhea in two or more consecutive cycles [see *Warnings and Precautions (5.8)*].

Manufactured In Canada By:

**Patheon Inc.**

Ontario, Canada L5N 7K9

Distributed by:

**Par Pharmaceutical**

Chestnut Ridge, NY 10977

Revised 08/2017

## **Patient Information**

**PREVIFEM<sup>®</sup>**

**TRI-PREVIFEM<sup>®</sup>**

**(norgestimate and ethinyl estradiol tablets, USP)**

**(nor-JES-ti-mate, ETH-i-nil es-tra-DYE-ol)**

**What is the most important information I should know about Prevfem<sup>®</sup> or Tri-Previfem<sup>®</sup>?**

**Do not use Prevfem<sup>®</sup> or Tri-Previfem<sup>®</sup> if you smoke cigarettes and are over 35 years old.**

Smoking increases your risk of serious cardiovascular side effects from hormonal birth control pills, including death from heart attack, blood clots or stroke. This risk increases with age and the number of cigarettes you smoke.

**What is Prevfem<sup>®</sup> or Tri-Previfem<sup>®</sup>?**

Prevfem<sup>®</sup> or Tri-Previfem<sup>®</sup> is a birth control pill (oral contraceptive) used by women to prevent pregnancy.

Tri-Previfem<sup>®</sup> is also used to treat moderate acne vulgaris in females 15 years of age and older, who have no known history of allergies or problems taking birth control pills, and have started their menstrual cycle (“period”). Tri-Previfem<sup>®</sup> should only be used to treat acne in women who want to take


birth control pills to prevent pregnancy.

### How does Previmem<sup>®</sup> or Tri-Previmem<sup>®</sup> work for contraception?

Your chance of getting pregnant depends on how well you follow the directions for taking your birth control pills. The better you follow the directions, the less chance you have of getting pregnant.

Based on the results of clinical studies, about 1 out of 100 women may get pregnant during the first year they use Previmem<sup>®</sup> or Tri-Previmem<sup>®</sup>.

The following chart shows the chance of getting pregnant for women who use different methods of birth control. Each box on the chart contains a list of birth control methods that are similar in effectiveness. The most effective methods are at the top of the chart. The box on the bottom of the chart shows the chance of getting pregnant for women who do not use birth control and are trying to get pregnant.



**Who should not take Previmem<sup>®</sup> or Tri-Previmem<sup>®</sup>? Do not take Previmem<sup>®</sup> or Tri-Previmem<sup>®</sup> if you:**

- smoke and are over 35 years of age

- had blood clots in your arms, legs, lungs, or eyes
- had a problem with your blood that makes it clot more than normal
- have certain heart valve problems or irregular heart beat that increases your risk of having blood clots
- had a stroke
- had a heart attack
- have high blood pressure that cannot be controlled by medicine
- have diabetes with kidney, eye, nerve, or blood vessel damage
- have certain kinds of severe migraine headaches with aura, numbness, weakness or changes in vision, or any migraine headaches if you are over 35 years of age
- have liver problems, including liver tumors
- take any Hepatitis C drug combination containing ombitasvir/paritaprevir/ritonavir, with or without dasabuvir. This may increase levels of the liver enzyme “alanine aminotransferase” (ALT) in the blood.
- have any unexplained vaginal bleeding
- are pregnant
- had breast cancer or any cancer that is sensitive to female hormones

**If any of these conditions happen while you are taking Prevfem<sup>®</sup> or Tri-Previfem<sup>®</sup>, stop taking Prevfem<sup>®</sup> or Tri-Previfem<sup>®</sup> right away and talk to your healthcare provider. Use non-hormonal contraception when you stop taking Prevfem<sup>®</sup> or Tri-Previfem<sup>®</sup>.**

**What should I tell my healthcare provider before taking Prevfem<sup>®</sup> or Tri-Previfem<sup>®</sup> ?**

**Tell your healthcare provider if you:**

- are pregnant or think you may be pregnant
- are depressed now or have been depressed in the past
- had yellowing of your skin or eyes (jaundice) caused by pregnancy (cholestasis of pregnancy)
- are breastfeeding or plan to breastfeed. Prevfem<sup>®</sup> or Tri-Previfem<sup>®</sup> may decrease the amount of breast milk you make. A small amount of the hormones in Prevfem<sup>®</sup> or Tri-Previfem<sup>®</sup> may pass into your breast milk. Talk to your healthcare provider about the best birth control method for you while breastfeeding.

**Tell your healthcare provider about all the medicines you take**, including prescription and over-the-counter medicines, vitamins and herbal supplements.

Prevfem<sup>®</sup> or Tri-Previfem<sup>®</sup> may affect the way other medicines work, and other medicines may affect how well Prevfem<sup>®</sup> or Tri-Previfem<sup>®</sup> works.

Know the medicines you take. Keep a list of them to show your healthcare provider and pharmacist when you get a new medicine.

**How should I take Prevfem<sup>®</sup> or Tri-Previfem<sup>®</sup> ? Read the Instructions for Use** at the end of this Patient Information.

**What are the possible serious side effects of Prevfem<sup>®</sup> or Tri-Previfem<sup>®</sup> ?**

- **Like pregnancy, Prevfem<sup>®</sup> or Tri-Previfem<sup>®</sup> may cause serious side effects, including blood clots in your lungs, heart attack, or a stroke that may lead to death. Some other examples of serious blood clots include blood clots in the legs or eyes.**

Serious blood clots can happen especially if you smoke, are obese, or are older than 35 years of age. Serious blood clots are more likely to happen when you:

- first start taking birth control pills
- restart the same or different birth control pills after not using them for a month or more

**Call your healthcare provider or go to a hospital emergency room right away if you have:**

- leg pain that will not go away
- sudden severe shortness of breath
- sudden change in vision or blindness
- chest pain
- a sudden, severe headache unlike your usual headaches
- weakness or numbness in your arm or leg
- trouble speaking

**Other serious side effects include:**

- **liver problems, including:**
  - rare liver tumors
  - jaundice (cholestasis), especially if you previously had cholestasis of pregnancy. Call your healthcare provider if you have yellowing of your skin or eyes.
- **high blood pressure.** You should see your healthcare provider for a yearly check of your blood pressure.
- **gallbladder problems**
- **changes in the sugar and fat (cholesterol and triglycerides) levels in your blood**
- **new or worsening headaches including migraine headaches**
- **irregular or unusual vaginal bleeding and spotting between your menstrual periods, especially during the first 3 months of taking Prevfem<sup>®</sup> or Tri-Previfem<sup>®</sup>.**
- **depression**
- **possible cancer in your breast and cervix**
- **swelling of your skin especially around your mouth, eyes, and in your throat (angioedema).** Call your healthcare provider if you have a swollen face, lips, mouth tongue or throat, which may lead to difficulty swallowing or breathing. Your chance of having angioedema is higher if you have a history of angioedema.
- **dark patches of skin around your forehead, nose, cheeks and around your mouth, especially during pregnancy (chloasma).** Women who tend to get chloasma should avoid spending a long time in sunlight, tanning booths, and under sun lamps while taking Prevfem<sup>®</sup> or Tri-Previfem<sup>®</sup>. Use sunscreen if you have to be in the sunlight.

**What are the most common side effects of Prevfem<sup>®</sup> or Tri-Previfem<sup>®</sup>?**

- headache (migraine)
- breast pain or tenderness, enlargement or discharge
- stomach pain, discomfort, and gas
- vaginal infections and discharge
- mood changes, including depression
- nervousness
- changes in weight
- skin rash

These are not all the possible side effects of Prevfem<sup>®</sup> or Tri-Previfem<sup>®</sup>. For more information, ask your healthcare provider or pharmacist.

You may report side effects to the FDA at 1-800-FDA-1088.

**What else should I know about taking Prevfem<sup>®</sup> or Tri-Previfem<sup>®</sup>?**

- If you are scheduled for any lab tests, tell your healthcare provider you are taking Prevfem<sup>®</sup> or Tri-Previfem<sup>®</sup>. Certain blood tests may be affected by PREVIFEM<sup>®</sup> or Tri-Previfem<sup>®</sup>.
- Prevfem<sup>®</sup> or Tri-Previfem<sup>®</sup> does not protect against HIV infection (AIDS) and other sexually transmitted infections.

**How should I store Prevfem<sup>®</sup> or Tri-Previfem<sup>®</sup>?**

- Store Prevfem<sup>®</sup> or Tri-Previfem<sup>®</sup> at room temperature between 68°F to 77°F (20°C to 25°C).
- Keep Prevfem<sup>®</sup> or Tri-Previfem<sup>®</sup> and all medicines out of the reach of children.

- Store away from light.

## **General information about the safe and effective use of Prevfem<sup>®</sup> or Tri-Previfem<sup>®</sup>.**

Medicines are sometimes prescribed for purposes other than those listed in a Patient Information leaflet. Do not use PREVIFEM<sup>®</sup> or Tri-Previfem<sup>®</sup> for a condition for which it was not prescribed. Do not give Prevfem<sup>®</sup> or Tri-Previfem<sup>®</sup> to other people, even if they have the same symptoms that you have.

This Patient Information summarizes the most important information about Prevfem<sup>®</sup> or Tri-Previfem<sup>®</sup>. You can ask your pharmacist or healthcare provider for information about Prevfem<sup>®</sup> or Tri-Previfem<sup>®</sup> that is written for health professionals.

For more information, call Par Pharmaceutical at 1-800-828-9393.

### **Do birth control pills cause cancer?**

Birth control pills do not seem to cause breast cancer. However, if you have breast cancer now, or have had it in the past, do not use birth control pills because some breast cancers are sensitive to hormones.

Women who use birth control pills may have a slightly higher chance of getting cervical cancer. However, this may be due to other reasons such as having more sexual partners.

### **What if I want to become pregnant?**

You may stop taking the pill whenever you wish. Consider a visit with your healthcare provider for a pre-pregnancy checkup before you stop taking the pill.

### **What should I know about my period when taking Prevfem<sup>®</sup> or Tri-Previfem<sup>®</sup>?**

Your periods may be lighter and shorter than usual. Some women may miss a period. Irregular vaginal bleeding or spotting may happen while you are taking Prevfem<sup>®</sup> or Tri-Previfem<sup>®</sup>, especially during the first few months of use. This usually is not a serious problem. It is important to continue taking your pills on a regular schedule to prevent a pregnancy.

### **What are the ingredients in Prevfem<sup>®</sup> or Tri-Previfem<sup>®</sup>?**

Prevfem<sup>®</sup>

**Active ingredients:** Each blue tablet contains norgestimate and ethinyl estradiol.

#### **Inactive ingredients:**

Blue tablets: FD&C Blue No. 1 HT Aluminum Lake, hypromellose, lactose monohydrate, magnesium stearate, polyethylene glycol and pregelatinized starch.

Light-green tablets: FD&C Blue No. 2, hypromellose, iron oxide yellow, lactose monohydrate, magnesium stearate, polyethylene glycol and pregelatinized starch.

Tri-Previfem<sup>®</sup>

**Active ingredients:** Each white, light blue and blue tablet contains norgestimate and ethinyl estradiol.

#### **Inactive ingredients:**

White tablets: hypromellose, lactose monohydrate, magnesium stearate, polyethylene glycol, and pregelatinized starch.

Light-blue tablet: FD&C Blue No. 1 Aluminum Lake, hypromellose, lactose monohydrate, magnesium stearate, polyethylene glycol and pregelatinized starch.

Blue tablet: FD&C Blue No. 1 Aluminum Lake, hypromellose, lactose monohydrate, magnesium stearate, polyethylene glycol and pregelatinized starch.

Light-green tablet: FD&C Blue No. 2, hypromellose, iron oxide yellow, lactose monohydrate, magnesium stearate, polyethylene glycol and pregelatinized starch.



## Instructions For Use

Previfem<sup>®</sup>

Tri-Previfem<sup>®</sup>

(norgestimate and ethinyl estradiol tablets, USP)

(nor-JES-ti-mate, ETH-i-nil es-tra-DYE-ol)

### Important Information about taking Previfem<sup>®</sup> or Tri-Previfem<sup>®</sup>

- Take **1** pill every day at the same time. Take the pills in the order directed on your blister pack pill dispenser.
- Do not skip your pills, even if you do not have sex often. If you miss pills (including starting the pack late) **you could get pregnant**. The more pills you miss, the more likely you are to get pregnant.
- If you have trouble remembering to take Previfem<sup>®</sup> or Tri-Previfem<sup>®</sup>, talk to your healthcare provider. When you first start taking Previfem<sup>®</sup> or Tri-Previfem<sup>®</sup>, spotting or light bleeding in between your periods may occur. Contact your healthcare provider if this does not go away after a few months.
- You may feel sick to your stomach (nauseous), especially during the first few months of taking Previfem<sup>®</sup> or Tri-Previfem<sup>®</sup>. If you feel sick to your stomach, do not stop taking the pill. The problem will usually go away. If your nausea does not go away, call your healthcare provider.
- Missing pills can also cause spotting or light bleeding, even when you take the missed pills later. On the days you take 2 pills to make up for missed pills (see **What should I do if I miss any Previfem<sup>®</sup> or Tri-Previfem<sup>®</sup> pills?** below), you could also feel a little sick to your stomach.
- It is not uncommon to miss a period. However, if you miss a period and have not taken Previfem<sup>®</sup> or Tri-Previfem<sup>®</sup> according to directions, or miss **2** periods in a row, or feel like you may be pregnant, call your healthcare provider. If you have a positive pregnancy test, you should stop taking Previfem<sup>®</sup> or Tri-Previfem<sup>®</sup>.
- If you have vomiting or diarrhea within **3-4** hours of taking your pill, take another pill of the same color from your extra blister pack pill dispenser. If you do not have an extra blister pack pill dispenser, take the next pill in your blister pack pill dispenser. Continue taking all your remaining pills in order. Start the first pill of your next blister pack pill dispenser the day after finishing your current blister pack pill dispenser. This will be 1 day earlier than originally scheduled. Continue on your new schedule.
- If you have vomiting or diarrhea for more than 1 day, your birth control pills may not work as well. Use an additional birth control method, like condoms and a spermicide, until you check with your healthcare provider.
- Stop taking Previfem<sup>®</sup> or Tri-Previfem<sup>®</sup> at least **4** weeks before you have major surgery and do not restart after the surgery without asking your healthcare provider. Be sure to use other forms of contraception (like condoms and spermicide) during this time period.

### Before you start taking Previfem<sup>®</sup> or Tri-Previfem<sup>®</sup>:

- Decide what time of day you want to take your pill. It is important to take it at the same time every day and in the order as directed on your blister pack pill dispenser.
- Have backup contraception (condoms and spermicide) available and if possible, an extra full blister pack of pills as needed.

### When should I start taking Previfem<sup>®</sup> or Tri-Previfem<sup>®</sup>?

#### If you start taking Previfem<sup>®</sup> or Tri-Previfem<sup>®</sup> and you have not used a hormonal birth control method before:

- There are 2 ways to start taking your birth control pills. You can either start on a Sunday (Sunday Start) or on the first day (Day 1) of your natural menstrual period (Day 1 Start). Your healthcare provider should tell you when to start taking your birth control pill.
- If you use the Sunday Start, use non-hormonal backup contraception such as condoms and

If you use the Sunday Start, use non-hormonal back-up contraception such as condoms and spermicide for the first 7 days that you take Prevfem<sup>®</sup> or Tri-Previfem<sup>®</sup>. You do not need back-up contraception if you use the Day 1 Start.

**If you start taking Prevfem<sup>®</sup> or Tri-Previfem<sup>®</sup> and you are switching from another birth control pill:**

- Start your new Prevfem<sup>®</sup> or Tri-Previfem<sup>®</sup> pack on the same day that you would start the next pack of your previous birth control method.
- Do not continue taking the pills from your previous birth control pack.

**If you start taking Prevfem<sup>®</sup> or Tri-Previfem<sup>®</sup> and previously used a vaginal ring or transdermal patch:**

- Start using Prevfem<sup>®</sup> or Tri-Previfem<sup>®</sup> on the day you would have reapplied the next ring or patch.

**If you start taking Prevfem<sup>®</sup> or Tri-Previfem<sup>®</sup> and you are switching from a progestin-only method such as an implant or injection:**

- Start taking Prevfem<sup>®</sup> or Tri-Previfem<sup>®</sup> on the day of removal of your implant or on the day when you would have had your next injection.

**If you start taking Prevfem<sup>®</sup> or Tri-Previfem<sup>®</sup> and you are switching from an intrauterine device or system (IUD or IUS):**

- Start taking Prevfem<sup>®</sup> or Tri-Previfem<sup>®</sup> on the day of removal of your IUD or IUS.
- You do not need back-up contraception if your IUD or IUS is removed on the first day (Day 1) of your period. If your IUD or IUS is removed on any other day, use non-hormonal back-up contraception such as condoms and spermicide for the first 7 days that you take Prevfem<sup>®</sup> or Tri-Previfem<sup>®</sup>.

**Keep a calendar to track your period:**

If this is the first time you are taking birth control pills, read, “**When should I start taking Prevfem<sup>®</sup> or Tri-Previfem<sup>®</sup>?**” above. Follow these instructions for either a **Sunday Start** or a **Day 1 Start**.

**Sunday Start:**

You will use a **Sunday Start** if your healthcare provider told you to take your first pill on a Sunday.

- Take pill **1** on the Sunday **after your period starts**.
- If your period starts on a Sunday, take the first pill that day and refer to Day 1 Start instructions below.
- Take **1** pill every day in the order on the blister pack pill dispenser at the same time each day for **28** days.
- After taking the last pill on **Day 28** from the blister pack pill dispenser, start taking the first pill from a new blister pack, on the same day of the week as the first blister pack (Sunday). Take the first pill in the new blister pack whether or not you are having your period.
- Use non-hormonal back-up contraception such as condoms and spermicide for the first 7 days of the first cycle that you take Prevfem<sup>®</sup> or Tri-Previfem<sup>®</sup>.


**Day 1 Start:**

You will use a **Day 1 Start** if your doctor told you to take your first pill (Day 1) on the **first day of your period**.

- Take **1** pill every day in the order of the blister pack pill dispenser, at the same time each day, for **28** days.
- After taking the last pill on **Day 28** from the blister pack pill dispenser, start taking the first pill from a new blister pack, on the same day of the week as the first blister pack. Take the first pill in the new blister pack whether or not you are having your period.

**Previfem® and Tri-Previfem® come in a blister pack pill dispenser. Read the instructions below for using your blister pack pill dispenser.**

Your blister package consists of three parts, the calendar label, the sleeve and the blister pack containing 28 individually sealed pills. Note that the pills are arranged in four numbered rows of 7 pills, with the pre-printed days of the week printed above them. Refer to the sample of the blister pack below:



Previfem® consists of 21 blue “active” birth control pills and 7 light green “reminder” pills.

Tri-Previfem® consists of 7 white “active” pills, 7 light-blue “active” pills, 7 blue “active” pills and 7 light green “reminder” pills.

There are two ways to start taking birth-control pills, Sunday Start or Day 1 Start. Your healthcare professional will tell you which to use.

**How to use Blister Cards for the 28 tablets**

1. If Sunday Start, discard the sticker and take the first active pill on the first Sunday after your menstrual period begins. Due to the potential risk of becoming pregnant, use additional non-hormonal contraception (such as condom and spermicide) for the first seven days of your first cycle pack of Previfem® or Tri-Previfem®.
2. If Day 1 Start, pick the Days of the Week Sticker that starts the first day of your period. (This is the day you begin bleeding or spotting, even if it is midnight when bleeding begins.) When you have picked the right sticker, throw away the others and place the sticker on the blister card over the preprinted days of the week and make sure it lines up with the pills.
3. Remove the first pill by pushing down on the pill. Swallow the pill. You will take 1 pill every day, at the same time each day. Wait 24 hours to take your next pill. Continue to take 1 pill each day until all the pills have been taken.
4. It is important to take the correct pill each day and not miss any pills. To help you remember, take your pill at the same time as another daily activity, like turning off your alarm clock or brushing your teeth.
5. After taking the last pill, start a new blister pack the very next day, no matter when your next period starts.
6. Take the pills in each new package as before. Start with the pill on row #1 and take one pill each day, left to right, until the last pill has been taken.

Three Ways to Remember in What Order to Take the Pills

1. Follow the sticker with the days of the week (placed above the pills).
2. Always go from left to right.
3. Always finish all your pills.

**What should I do if I miss any Prevfem<sup>®</sup> or Tri-Prefvem<sup>®</sup> pills?**

**If you miss 1 pill in Weeks 1, 2, or 3, follow these steps:**

- Take it as soon as you remember. Take the next pill at your regular time. This means you may take 2 pills in 1 day.
- Then continue taking 1 pill every day until you finish the pack.
- You do not need to use a back-up birth control method if you have sex.

**If you miss 2 pills in Week 1 or Week 2 of your pack, follow these steps:**

- Take the 2 missed pills as soon as possible and the next 2 pills the next day.
- Then continue to take 1 pill every day until you finish the pack.
- Use a non-hormonal birth control method (such as a condom and spermicide) as a back-up if you have sex during the first 7 days after missing your pills.

**If you miss 2 pills in a row in Week 3, or you miss 3 or more pills in a row during Weeks 1, 2, or 3 of the pack, follow these steps:**

- **If you are a Day 1 Starter:**
  - Throw out the rest of the pill pack and start a new pack that same day.
  - You may not have your period this month but this is expected. However, if you miss your period 2 months in a row, call your healthcare provider because you might be pregnant.
  - You could become pregnant if you have sex during the first 7 days after you restart your pills. You **MUST** use a non-hormonal birth control method (such as a condom and spermicide) as a back-up if you have sex during the first 7 days after you restart your pills.
- **If you are a Sunday Starter:**
  - Keep taking 1 pill every day until Sunday. On Sunday, throw out the rest of the pack and start a new pack of pills that same day.
  - Use a non-hormonal birth control method (such as a condom and spermicide) as a back-up if you have sex during the first 7 days after you restart your pills.

**If you have any questions or are unsure about the information in this leaflet, call your healthcare provider.**

This Patient Information and Instructions for Use has been approved by the U.S. Food and Drug Administration.

All brand names listed are registered trademarks of their respective owners and are not trademarks of Par Pharmaceutical.

Manufactured In Canada By:

**Patheon Inc.**

Ontario, Canada L5N 7K9

Distributed by:

**Par Pharmaceutical**

Chestnut Ridge, NY 10977

Revised 08/2017

**Prevfem Tablets 28-Day Regimen**

NDC# 53002-1191-1  
 LIST# 0603-7642-01  
**PREVIFEM TABLETS 28-DAY REGIMEN**  
 28 TABLETS  
 LOT# 17355-400  
 EXP DATE: 01-31-2019  
 Shape & Color Markings  
 BLUE & LT GREEN 95 T48 (BLUE)  
 ROUND TABLETS 90 T42 (GREEN)  
 EACH BLUE TABLET CONTAINS:  
 NORGESTIMATE 0.25MG  
 ETHINYL ESTRADIOL 0.020MG  
 EACH LT GREEN TABLET IS INERT

28 TABLETS PATIENT NAME: \_\_\_\_\_  
**PREVIFEM TABLETS 28-DAY**  
 QUALITY Rx only  
 TAKE 1 TABLET DAILY  
 AS DIRECTED.

CLINIC NAME GOES HERE  
 Prescriber Name: \_\_\_\_\_  
 Date Dispensed: \_\_\_\_\_  
 IMPORTANT: READ PATIENT INFORMATION SHEET CAREFULLY BEFORE USING. TAKE ONLY AS DIRECTED BY YOUR PRESCRIBER.



173610401000 ITEM# 1191

1 TAB DAILY U0  
 LOT# 17355-001 EXP 01-31-2019  
 R# 173610401 - 000 FCA- 1191  
 28 ea PREVIFEM TABLETS 28-DAY  
 BILLING NDC# 00630-7642-01  
 R# 173610401 - 000  
 28 ea PREVIFEM TABLETS 28-DAY  
 BILLING NDC# 00630-7642-01  
 R# 173610401 - 000  
 28 ea PREVIFEM TABLETS 28-DAY

**PREVIFEM TABLETS 28-DAY REGIMEN**  
 28 TABLETS  
 DISCARD BY 01-31-2019  
 NDC# 53002-1191-1  
 R# 173610401 - 000

**PREVIFEM**  
 norgestimate and ethinyl estradiol kit

Product Information			
Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:53002-1191(NDC:0603-7642)

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:53002-1191-6	6 in 1 PACKAGE	09/01/2017	
1		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		
2	NDC:53002-1191-3	3 in 1 PACKAGE	09/01/2017	
2		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		
3	NDC:53002-1191-1	1 in 1 PACKAGE	09/01/2017	
3		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Quantity of Parts		
Part #	Package Quantity	Total Product Quantity
Part 1		21
Part 2		7

**Part 1 of 2**  
**PREVIFEM**  
 norgestimate and ethinyl estradiol tablet, film coated

Product Information	
Route of Administration	ORAL

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
NORGESTIMATE (UNII: C291HFX4DY) (NORGESTIMATE - UNII:C291HFX4DY)	NORGESTIMATE	0.25 mg
ETHINYL ESTRADIOL (UNII: 423D2T571U) (ETHINYL ESTRADIOL - UNII:423D2T571U)	ETHINYL ESTRADIOL	0.035 mg

### Inactive Ingredients

Ingredient Name	Strength
FD&C BLUE NO. 1 (UNII: HBR47K3TBD)	
HYPROMELLOSE 2910 (6 MPAS) (UNII: 0WZ8WG20P6)	
LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ)	
POLYETHYLENE GLYCOL 8000 (UNII: Q662QK8M3B)	
STARCH, CORN (UNII: O8232NY3SJ)	

### Product Characteristics

Color	BLUE	Score	no score
Shape	ROUND	Size	6mm
Flavor		Imprint Code	93;748
Contains			

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA076334	01/09/2004	

### Part 2 of 2

#### INERT

inert tablet, film coated

### Product Information

Route of Administration	ORAL
-------------------------	------

### Inactive Ingredients

Ingredient Name	Strength
FD&C BLUE NO. 2 (UNII: L06K8R7DQK)	
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)	

### Product Characteristics

Color	GREEN (light-green)	Score	no score
Shape	ROUND	Size	6mm

<b>Flavor</b>		<b>Imprint Code</b>	93;743
<b>Contains</b>			
<b>Marketing Information</b>			
<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
ANDA	ANDA076334	01/09/2004	
<b>Marketing Information</b>			
<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
ANDA	ANDA076334	01/09/2004	

**Labeler** - RPK Pharmaceuticals, Inc. (147096275)

**Establishment**

<b>Name</b>	<b>Address</b>	<b>ID/FEI</b>	<b>Business Operations</b>
RPK Pharmaceuticals, Inc.		147096275	RELABEL(53002-1191) , REPACK(53002-1191)

Revised: 9/2018

RPK Pharmaceuticals, Inc.