

**RIGHT REMEDIES EXTRA STRENGTH PAIN RELIEF PM CAPLET- acetaminophen,
diphenhydramine hydrochloride tablet
Strive Pharmaceuticals Inc.**

RIGHT REMEDIES EXTRA STRENGTH Pain Relief PM Caplet

Drug Facts

Active ingredients (in each caplet)

Acetaminophen 500 mg
Diphenhydramine HCl 25 mg

Purpose

Pain reliever
Nighttime sleep aid

Uses

• Temporary relief of occasional headaches and minor aches and pains with accompanying sleeplessness

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash. If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- with any other product containing diphenhydramine, even one used on skin
- in children under 12 years of age
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have • liver disease • a breathing problem such as emphysema or chronic bronchitis • trouble urinating due to an enlarged prostate gland • glaucoma

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

When using this product

- drowsiness will occur
- avoid alcoholic drinks
- do not drive a motor vehicle or operate machinery

Stop use and ask a doctor if

- sleeplessness persists continuously for more than 2 weeks. Insomnia may be a symptom of serious underlying medical illness
 - pain gets worse or lasts more than 10 days
 - fever gets worse or lasts more than 3 days
 - redness or swelling is present
 - new symptoms occur.
- These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: in case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not take more than directed (see overdose warning)**

adults and children 12 years and over	<ul style="list-style-type: none"> • take 2 caplets at bedtime • do not take more than 2 caplets of this product in 24 hours
children under 12 years	do not use

Other information

- store at room temperature between 20°-25°C (68°-77°F)
- protect from light, heat and moisture

Inactive ingredients

FD&C Blue No. 1, FD&C Blue No. 2, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol 400, polyvinylpyrrolidone, pregelatinized starch, stearic acid powder, titanium dioxide

Questions or comments?

1-888-577-8033 Monday - Friday 8am - 4pm EST

Compare to the active ingredients in Extra Strength **Tylenol® PM***

TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS

BROKEN OR MISSING

*This product is not manufactured or distributed by Kenvue, Inc., owner of the registered trademark Tylenol®.

Distributed by:

Strive Pharmaceuticals Inc.
East Brunswick, NJ 08816

PRODUCT OF INDIA

Packaging



Compare to the active ingredients in Extra Strength **TYLENOL[®] PM[®]**

NDC 70692-471-31

EXTRA STRENGTH Pain Relief PM

Acetaminophen 500 mg
Diphenhydramine HCl 25 mg

+ Pain reliever
Nighttime sleep aid



Actual Size

300
caplets

WARNING: EVIDENCE OF TAMPERING: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

Drug Facts

Active ingredients (in each caplet)
Acetaminophen 500 mg.....Pain reliever
Diphenhydramine HCl 25 mg.....Nighttime sleep aid

Uses ■ Temporary relief of occasional headaches and minor aches and pains with accompanying sleeplessness

Warnings
■ **Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take
■ more than 4,000 mg of acetaminophen in 24 hours ■ with other drugs containing acetaminophen
■ 3 or more alcoholic drinks every day while using this product
■ **Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:
■ skin reddening ■ blisters ■ rash. If a skin reaction occurs, stop use and seek medical help right away.
Do not use
■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
■ with any other product containing diphenhydramine, even one used on skin ■ in children under 12 years of age
■ if you have ever had an allergic reaction to this product or any of its ingredients
Ask a doctor before use if you have ■ liver disease ■ a breathing problem such as emphysema or chronic bronchitis ■ trouble urinating due to an enlarged prostate gland ■ glaucoma
Ask a doctor or pharmacist before use if you are
■ taking the blood thinning drug warfarin ■ taking sedatives or tranquilizers
When using this product
■ drowsiness will occur ■ avoid alcoholic drinks
■ do not drive a motor vehicle or operate machinery
Stop use and ask a doctor if
■ sleeplessness persists continuously for more than 2 weeks. Insomnia may be a symptom of serious underlying medical illness ■ pain gets worse or lasts more than 10 days
■ fever gets worse or lasts more than 3 days ■ redness or swelling is present ■ new symptoms occur. These could be signs of a serious condition.
If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children.
Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions
■ do not take more than directed (see overdose warning)
adults and children ■ take 2 caplets at bedtime
12 years and over ■ do not take more than 2 caplets of this product in 24 hours
children under 12 years do not use

Other information
■ store at room temperature between 20–25°C (68–77°F) ■ protect from light, heat and moisture

Inactive ingredients
FD&C Blue No. 1, FD&C Blue No. 2, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol 400, polyvinylpyrrolidone, pregelatinized starch, stearic acid powder, titanium dioxide

Questions or comments? 1-888-577-8033 Monday-Friday 8am-4pm EST

This product is not manufactured or distributed by Kenvue, Inc., owner of the registered trademark Tylenol[®].
L87069247131 REV:06-022025
Distributed by:
Kenvue Pharmaceuticals
East Brunswick, NJ 08816
PRODUCT OF INDIA
LOT:
EXP:
3 70692 0045 0



NO VARNISH

DRUG FACTS LABEL

Drug Facts	
Active ingredients (in each caplet)	Purpose
Acetaminophen 500 mg.....	Pain reliever
Diphenhydramine HCl 25 mg.....	Nighttime sleep aid
Uses ■ Temporary relief of occasional headaches and minor aches and pains with accompanying sleeplessness	
Warnings Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take ■ more than 4,000 mg of acetaminophen in 24 hours ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash. If a skin reaction occurs, stop use and seek medical help right away. Do not use ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ with any other product containing diphenhydramine, even one used on skin ■ in children under 12 years of age ■ if you have ever had an allergic reaction to this product or any of its ingredients Ask a doctor before use if you have ■ liver disease ■ a breathing problem such as emphysema or chronic bronchitis ■ trouble urinating due to an enlarged prostate gland ■ glaucoma Ask a doctor or pharmacist before use if you are ■ taking the blood thinning drug warfarin ■ taking sedatives or tranquilizers When using this product ■ drowsiness will occur ■ avoid alcoholic drinks ■ do not drive a motor vehicle or operate machinery Stop use and ask a doctor if ■ sleeplessness persists continuously for more than 2 weeks. Insomnia may be a symptom of serious underlying medical illness ■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present ■ new symptoms occur. These could be signs of a serious condition. If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.	
Directions ■ do not take more than directed (see overdose warning)	
adults and children	■ take 2 caplets at bedtime
12 years and over	■ do not take more than 2 caplets of this product in 24 hours
children under 12 years	do not use
Other information ■ store at room temperature between 20–25°C (68–77°F) ■ protect from light, heat and moisture	
Inactive ingredients FD&C Blue No. 1, FD&C Blue No. 2, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol 400, polyvinylpyrrolidone, pregelatinized starch, stearic acid powder, titanium dioxide	
Questions or comments? 1-888-577-8033 Monday-Friday 8am-4pm EST	

RIGHT REMEDIES EXTRA STRENGTH PAIN RELIEF PM CAPLET

acetaminophen, diphenhydramine hydrochloride tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:70692-471
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	500 mg
DIPHENHYDRAMINE HYDROCHLORIDE (UNII: TC2D6JAD40) (DIPHENHYDRAMINE - UNII:8GTS82S83M)	DIPHENHYDRAMINE HYDROCHLORIDE	25 mg

Inactive Ingredients

Ingredient Name	Strength
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
FD&C BLUE NO. 2 (UNII: L06K8R7DQK)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
STARCH, CORN (UNII: O8232NY3S)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	blue	Score	no score
Shape	OVAL	Size	18mm
Flavor		Imprint Code	S525
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:70692-471-31	300 in 1 BOTTLE; Type 0: Not a Combination Product	03/17/2025	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	03/17/2025	

Labeler - Strive Pharmaceuticals Inc. (080028013)

Revised: 3/2025

Strive Pharmaceuticals Inc.