

CREST DEEP CLEANSE GERM TREATMENT- cetylpyridinium chloride rinse
The Procter & Gamble Manufacturing Company

Crest Deep Cleanse Germ Treatment Rinse

Drug Facts

Active ingredient

Cetylpyridinium chloride 0.07%

Purpose

Antigingivitis/Antiplaque

Uses

- helps prevent and reduce plaque and gingivitis
- helps control plaque bacteria that contribute to the development of gingivitis and bleeding gums

Warning

Ask a dentist if symptoms persist or condition worsens after regular use.

Keep out of reach of children 6 years of age. If more than used for rinsing is accidentally swallowed, get medical help or contact a Poison Control Center right away.

Directions

- use after your normal brushing and flossing routine, rinse toothpaste from mouth prior to use
- adults and children 6 yrs. & older: Rinse for 30 seconds with 20 mL (4 teaspoonfuls) twice a day
- do not swallow
- children 6 years to under 12 years of age: supervise use
- children under 6 years of age: do not use

Inactive ingredients

water, glycerin, flavor, zinc lactate, methylparaben, sodium saccharin, sucralose, propylparaben, poloxamer 407

Questions?

1-800-285-9139

DISTR. BY PROCTER & GAMBLE, CINCINNATI, OH 45202

PRINCIPAL DISPLAY PANEL - 475 mL Bottle Label

DENTIST
DEVELOPED

Crest

CPC ANTIGINGIVITIS/ANTIPLAQUE MOUTHWASH

DEEP
CLEANSE
MOUTHWASH

GERM TREATMENT

MILD MINT
No Burn of Alcohol
ALCOHOL

0%

FREE

SLS

FREE

PLAQUE

24

HOUR

DEFENSE

475 mL (16 FL OZ)



Crest
CPC ANTIGINGIVITIS/ANTIPLAQUE MOUTHWASH

DEEP CLEANSE MOUTHWASH

GERM TREATMENT

MILD MINT

No Burn of Alcohol



475 mL (16 FL OZ)

GERM TREATMENT** IN 3 STEPS

POUR

Pour 20 mL of rinse in the dosing cup

SWISH

Swish around your mouth for 30 seconds then spit

REPEAT

Use twice per day for Germ Treatment

Drug Facts

Active Ingredient

Cetylpyridinium chloride 0.07%..... Antigingivitis/Antiplaque

Uses

- helps prevent and reduce plaque and gingivitis
- helps control plaque bacteria that contribute to the development of gingivitis and bleeding gums

Warnings

Ask a dentist if symptoms persist or condition worsens after regular use.

Keep out of reach of children under 6 years of age. If more than used for rinsing is accidentally swallowed, get medical help or contact a Poison Control Center right away.

Directions

| | |
|---|--|
| • use after your normal brushing and flossing routine, rinse toothpaste from mouth prior to use | Rinse for 30 seconds with 20 mL (4 teaspoonsful) |
| • adults and children 6 yrs. & older: | |
| • children 6 years to under 12 years of age: | • do not swallow |
| • children under 6 years of age: | supervise use |
| | do not use |

Inactive Ingredients water, glycerin, flavor, zinc lactate, methylparaben, sodium saccharin, sucrose, propylparaben, polyoxamer 407

Questions? 1-800-205-0139

*With twice daily use

**Kills the germs that cause plaque, gingivitis and bad breath

In some cases, antimicrobial rinses may cause surface staining to teeth. This is not permanent or harmful, and may be prevented by adequate brushing or removed at your next dental visit.

If you are not satisfied with your results, Crest will refund your purchase via pre-paid card. Simply return your receipt and copy of bottle UPC within 60 days of purchase. Call 1-877-789-8784 for more information.

CAP SEALED WITH PRINTED SLEEVE

DISTR. BY PROCTER & GAMBLE, CINCINNATI, OH 45202

P&G www.pg.com www.crest.com

Patents: www.pg.com/patents © 2024 P&G



CREST DEEP CLEANSE GERM TREATMENT

cetylpyridinium chloride rinse

Product Information

| | | | |
|-------------------------|----------------|--------------------|---------------|
| Product Type | HUMAN OTC DRUG | Item Code (Source) | NDC:84126-606 |
| Route of Administration | ORAL | | |

Active Ingredient/Active Moiety

| Ingredient Name | Basis of Strength | Strength |
|-----------------|-------------------|----------|
| | | |

| | | |
|--|--------------------------|----------------|
| CETYLPYRIDINIUM CHLORIDE (UNII: D9OM4SK49P) (CETYLPYRIDINIUM - UNII:CUB7JI0JV3) | CETYLPYRIDINIUM CHLORIDE | 0.07 g in 1 mL |
|--|--------------------------|----------------|

Inactive Ingredients

| Ingredient Name | Strength |
|--|----------|
| WATER (UNII: 059QF0KO0R) | |
| GLYCERIN (UNII: PDC6A3C00X) | |
| METHYLPARABEN (UNII: A2I8C7HI9T) | |
| PROPYLPARABEN (UNII: Z8IX2SC1OH) | |
| POLOXAMER 407 (UNII: TUF2IVW3M2) | |
| SUCRALOSE (UNII: 96K6UQ3ZD4) | |
| ZINC LACTATE (UNII: 2GXR25858Y) | |
| SACCHARIN SODIUM (UNII: SB8ZUX40TY) | |

Product Characteristics

| | | |
|----------|-------|--------------|
| Color | Score | |
| Shape | Size | |
| Flavor | MINT | Imprint Code |
| Contains | | |

Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|---|------------------|--|----------------------|--------------------|
| 1 | NDC:84126-606-16 | 475 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product | 01/01/2024 | |

Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|--------------------|--|----------------------|--------------------|
| OTC Monograph Drug | M022 | 01/01/2024 | |

Labeler - The Procter & Gamble Manufacturing Company (004238200)

Revised: 1/2026

The Procter & Gamble Manufacturing Company