ADAPALENE- adapalene gel YYBA Corp

Welmate Adapalene Gel USP, 0.1% Paraben Free

Drug Facts

Active ingredient

Adapalene USP, 0.1% (retinoid) ¹

1 read consumer information leaflet

Purpose

Acne treatment

Use

For the treatment of acne

Warnings

For external use only

Do not use

- on damaged skin (cuts, abrasions, eczema, sunburn)
- if you are allergic to adapalene or any of the ingredients in this product.

If pregnant or breast-feeding, ask a doctor before use.

When using this product

- limit sun exposure, including light from tanning beds, and use sunscreen when going outdoors
- do not wax to remove hair in areas where the product has been applied
- during the early weeks of use, your acne may appear to worsen before it improves (this is normal); continue using as directed, unless you get irritation that becomes severe
- irritation (redness, itching, dryness, burning) is more likely to occur:
 - in the first few weeks of use
 - if using more than one topical acne medication at a time
 - but irritation usually lessens with continued use of this product
- it may take up to 3 months of once daily use to see results
- avoid product contact with eyes, lips, and mouth. If contact occurs, immediately flush the area with water.
- wash hands after use

Stop use and ask doctor if

- you become pregnant, or are planning to become pregnant, while using the product
- you have symptoms of an allergic reaction (such as itching, rash, hives, swelling of the lips, eyelids, and shortness of breath)
- irritation becomes severe
- you see no improvement after 3 months of once daily use

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

Adults and children 12 years of age and older:

- use **once** daily
- clean the skin gently and pat dry before applying the product
- cover the entire affected area with a thin layer. For example, if your acne is on the face, apply the product to the entire face.
- do not use more than one time a day. Applying more than directed will not provide faster or better results, but may worsen skin irritation.

Children under 12 years of age: ask a doctor

Other information

- TAMPER EVIDENT: DO NOT USE IF THE SEAL ON THE TUBE IS PUNCTURED OR NOT VISIBLE.
- store at 20° to 25°C (68° to 77°F) [see USP Controlled Room Temperature].
- protect from freezing

Inactive ingredients

carbomer homopolymer type C, edetate disodium, phenoxyethanol, poloxamer 182, propylene glycol, purified water, sodium hydroxide.

Questions?

1-866-923-4914

Distributed by: Wellspring Airmont, NY 10952, U.S.A.

Made in Canada


PRINCIPAL DISPLAY PANEL - 45 g Tube Carton

Compare to the active ingredient in Differin [®] Gel [†]

NDC 73581-019-06

Adapalene Gel USP, 0.1% Acne Treatment

NET WT 45 g (1.6 oz)



ADAPALENE

adapalene gel

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:73581-019
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety			
	Ingredient Name	Basis of Strength	Strength
	ADAPALENE (UNII: 1L4806J2QF) (ADAPALENE - UNII:1L4806J2QF)	ADAPALENE	1 mg in 1 g

Inactive Ingredients			
Ingredient Name	Strength		
CARBOMER HOMOPOLYMER TYPE C (ALLYL PENTAERYTHRITOL CROSSLINKED) (UNII: 4Q93RCW27E)			
EDETATE DISODIUM (UNII: 7FLD91C86K)			
PHENOXYETHANOL (UNII: HIE492ZZ3T)			
POLOXAMER 182 (UNII: JX0HIX6OAG)			
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)			
WATER (UNII: 059QF0KO0R)			
SODIUM HYDROXIDE (UNII: 55X04QC32I)			

Packaging				
#	tem Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:73581-019- 06	1 in 1 CARTON	03/26/2025	
1		45 g in 1 TUBE; Type 0: Not a Combination Product		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA215940	03/26/2025	

Labeler - YYBA Corp (006339772)

Establishment				
Name	Address	ID/FEI	Business Operations	
Taro Pharmaceuticals Inc.		206263295	manufacture(73581-019)	

Revised: 3/2025 YYBA Corp