

**METHYL SALICYLATE- methyl salicylate 10% patch**  
**Florrx Pharmaceutical Corp.**

-----

**Directions**

Adults and children 18 years of age and older:

Apply 1 patch to the affected area of intact skin up to 3 times a day. Do not leave patch on for more than 8 hours at a time.

- Clean and dry affected area
- Open pouch and remove one patch
- Remove any protective film and apply directly to affected area of pain. Apply immediately after removal from the protective envelope.
- Wash hands with soap and water after handling the patches.
- Reseal pouch containing unused patches after each use. Do not store patch outside the sealed envelope.

Fold used patches so that the adhesive side sticks to itself and safely discard used patches or pieces of cut patches where children and pets cannot get to them.

**Children under 18 years: Ask a physician**

**Purpose**

External Analgesic

**Uses**

For the temporary relief of minor aches and pains of muscles and joints, such as simple backache, lumbago, arthritis, neuralgia, strains, bruises, and sprains.

**Warnings**

For external use only. Use only as directed or as directed by a healthcare professional. Read all warnings and directions before use. Discontinue use at least one hour prior to bath, shower, or swimming; do not use immediately after bath, shower, or swimming.

**Do not use**

- On wounds, cuts, damaged or infected skin
- On eyes, mouth, genitals, or any other mucous membranes

Allergy Alert: if you are allergic to capsicum or chili peppers or any inactive ingredient of this product, contact a doctor before use.

**When using this product**

- You may experience a burning sensation. The intensity of this reaction varies among individuals and may be severe. With regular use, this sensation generally disappears after several days.
- Avoid contact with the eyes, lips, nose and mucous membranes
- You may experience a burning sensation. The intensity of this reaction varies among individuals and may be severe. With regular use, this sensation generally disappears after several days.
- Avoid contact with the eyes, lips, nose and mucous membranes
- Do not tightly wrap or bandage the treated area
- Do not apply heat to the treated area immediately before or after use

**Stop use and ask a physician if:**

- allergic reaction occurs
- condition worsens or does not improve within 7 days
- symptoms clear up and return within a few days
- redness, irritation, swelling, pain or other symptoms begin or increase

**KEEP OUT OF REACH OF CHILDREN**

**CONSULT PHYSICIAN FOR CHILDREN UNDER 18.**

**Active Ingredient**

Methyl Salicylate 10%

**Other Information**

May be applied under occlusive dressing.

Store at 20-25°C (68-77°F); excursions permitted to 15-30°C (59-86°F). See USP Controlled Room Temperature

**Inactive Ingredients**

Polysorbate 80, Glycerin, Sodium Cellulose Glycolate, Sodium Polyacrylate, Vitamin E, EDTA Disodium Salt, Kaolin, Water, Methylparaben



## METHYL SALICYLATE

methyl salicylate 10% patch

### Product Information

|                         |                |                    |               |
|-------------------------|----------------|--------------------|---------------|
| Product Type            | HUMAN OTC DRUG | Item Code (Source) | NDC:84460-103 |
| Route of Administration | TOPICAL        |                    |               |

### Active Ingredient/Active Moiety

| Ingredient Name   | Basis of Strength | Strength      |
|---|-------------------|---------------|
| METHYL SALICYLATE (UNII: LAV5U5022Y) (SALICYLIC ACID - UNII:O414PZ4LPZ) | METHYL SALICYLATE | 10 g in 100 g |

### Inactive Ingredients

| Ingredient Name                     | Strength |
|-------------------------------------|----------|
| EDETATE DISODIUM (UNII: 7FLD91C86K) |          |
| KAOLIN (UNII: 24H4NWX5CO)           |          |

| <b>METHYLPARABEN</b> (UNII: A2I8C7HI9T)                 |                  |   |                      |                    |
|---|------------------|---|----------------------|--------------------|
| <b>SODIUM POLYACRYLATE (8000 MW)</b> (UNII: 285CYO341L) |                  |   |                      |                    |
| <b>CARBOXYMETHYLCELLULOSE SODIUM</b> (UNII: K679OBS311) |                  |   |                      |                    |
| <b>GLYCERIN</b> (UNII: PDC6A3C0OX)                      |                  |   |                      |                    |
| <b>WATER</b> (UNII: 059QF0KO0R)                         |                  |   |                      |                    |
| <b>POLYSORBATE 80</b> (UNII: 6OZP39ZG8H)                |                  |   |                      |                    |
| <b>ALPHA-TOCOPHEROL</b> (UNII: H4N855PNZ1)              |                  |   |                      |                    |
|   |                  |   |                      |                    |
| <b>Packaging</b>  |                  |   |                      |                    |
| #   | Item Code        | Package Description                                 | Marketing Start Date | Marketing End Date |
| 1   | NDC:84460-103-01 | 2 in 1 BOX  | 02/10/2025           |                    |
| 1   |                  | 5 in 1 POUCH  |                      |                    |
| 1   |                  | 100 g in 1 PATCH; Type 0: Not a Combination Product |                      |                    |
|   |                  |   |                      |                    |
| <b>Marketing Information</b>                            |                  |   |                      |                    |
| Marketing Category                                      |                  | Application Number or Monograph Citation            | Marketing Start Date | Marketing End Date |
| OTC Monograph Drug                                      |                  | 505G(a)(3)  | 02/10/2025           |                    |

**Labeler** - Florrax Pharmaceutical Corp. (119257612)

Revised: 2/2025

Florrax Pharmaceutical Corp.