GOOD NEIGHBOR PHARMACY 24 HOUR NASAL ALLERGY- triamcinolone acetonide spray, metered Praxis, LLC

Good Neighbor Pharmacy 24 Hour Nasal Allergy Spray Drug Facts

Active ingredient (in each spray)

Triamcinolone acetonide 55 mcg (glucocorticoid)

Purpose

Allergy symptom reliever

Uses

temporarily relieves these symptoms of hay fever or other upper respiratory allergies:

- nasal congestion
- runny nose
- sneezing
- itchy nose

Warnings

Do not use

- in children under 2 years of age
- if you have ever had an allergic reaction to any of the ingredients

Ask a doctor before use if you

- have had recent nose ulcers or nose surgery
- have had a nose injury that has not healed
- are using a steroid medicine for asthma, allergies or skin rash
- have an eye infection
- have or had glaucoma or cataracts

When using this product

- the growth rate of some children may be slower
- some symptoms may get better on the first day of treatment. It may take up to one week of daily use to feel the most symptom relief.
- do not share this bottle with anyone else as this may spread germs
- remember to tell your doctor about all the medicines you take, including this one

Stop use and ask a doctor if

- you have, or come into contact with someone who has, chickenpox, measles or tuberculosis
- you have or develop symptoms of an infection such as a persistent fever

- you have any change in vision
- you have severe or frequent nosebleeds

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

Read insert (inside package) on how to:

- get a new bottle ready (primed) before first use
- prime bottle again if not used for more than 2 weeks
- use the spray
- clean the spray nozzle

ADULTS	AND CHILDREN 12 YEARS OF AGE AND OLDER		
adults and children 12 years of age and older	 once daily, spray 2 times into each nostril while sniffing gently once your allergy symptoms improve, reduce to 1 spray in each nostril per day 		
C	HILDREN 2 TO UNDER 12 YEARS OF AGE		
-	some children may be slower while using this product. Talk to if your child needs to use the spray for longer than two months a		
children 6 to under 12 years of age	 an adult should supervise use once daily, spray 1 time into each nostril while sniffing gently if allergy symptoms do not improve, increase to 2 sprays in each nostril per day. Once allergy symptoms improve, reduce to 1 spray in each nostril per day. 		
children 2 to under 6 years of age	 an adult should supervise use once daily, spray 1 time into each nostril while sniffing gently 		
children under 2 years of age	• do not use		
• do not spray into ey	, do not double the next dose yes or mouth do not improve after one week, stop using and talk to a doctor common cold		

Other information

- do not use if printed bottle wrap is torn or opened
- keep package and insert. They contain important information.
- store between 20 °-25 °C (68 °-77 °F)

Inactive ingredients

benzalkonium chloride, carboxymethylcellulose sodium, dextrose anhydrous, edetate disodium dihydrate, hydrochloric acid or sodium hydroxide (for pH adjustment), microcrystalline cellulose, polysorbate 80, purified water

Questions or comments?

1-800-719-9260

Consumer information

Triamcinolone Acetonide Nasal Spray 55 mcg per spray


IMPORTANT INFORMATION

- **Read both sides of this insert** for complete instructions on how to get a bottle ready (primed), how to use the spray bottle and how to clean the spray nozzle.
- Keep this insert as it contains important information.

STEPS TO GET A NEW BOTTLE READY FOR USE (PRIMED)

1. Before first use, a new bottle must be primed.


- 1. Remove cap and safety clip (*Figure A*).
- 2. Shake bottle.
- 3. Press and release spray nozzle until a fine mist is produced (as shown in Figure B). This may take several times. Take care not to spray in face.




• Repeat steps for priming a bottle (see above).

USE INSTRUCTIONS

- 1. Blow nose gently to clear nostrils.
- 2. Remove cap and safety clip, then shake bottle.
- 3. Hold bottle with thumb under bottle and spray nozzle between fingers (as shown in Figure B).
- 4. Press against the outside of your nose with your finger to close off one nostril (as shown in Figure C).
- 5. Place the tip of the spray nozzle into the other nostril. The spray nozzle should not reach far into the nose. Aim nozzle toward back of nose (as shown in Figure C).



DO NOTspray toward nasal septum (the wall between the 2 nostrils) (as shown in Figure D).



1. 6. While sniffing gently, spray into nostril.

IMPORTANT: For complete dosing instructions, See "Directions for use" on next side.

- 1. 7. Repeat steps "4" through "6" for the other nostril.
- 2. 8. After using the nasal spray, wipe nozzle with a tissue, replace safety clip, and

replace cap by pressing it down over the spray nozzle.

NOTE: Avoid blowing nose for 15 minutes after use.

If nozzle does not spray properly, see cleaning instructions on next side.

(continued on side 2)

SIDE 1

DIRECTIONS FOR USE

	t use, you must get a new bottle ready
(primed).If bottle is n	not used for more than 2 weeks, prime
bottle again	· · ·
Dosing	
ADULTS AND	CHILDREN 12 YEARS OF AGE AND OLDER
adults and children 12	 once daily, spray 2 times into each nostril while sniffing gently
years of age and older	 once your allergy symptoms improve, reduce to 1 spray in each nostril per day
CHILDE	REN 2 TO UNDER 12 YEARS OF AGE
this product.	te of some children may be slower while using Talk to your child's doctor if your child needs to for longer than two months a year.
children 6 to under 12 years of age	 an adult should supervise use once daily, spray 1 time into each nostril while sniffing gently if allergy symptoms do not improve, increase to 2 sprays in each nostril per day. Once allergy symptoms improve, reduce to 1 spray in each nostril per day.
children 2 to under 6 years of age	 an adult should supervise use once daily, spray 1 time into each nostril while sniffing gently
children under 2 years of age	• do not use


Other important information for use

- Do not use more than directed.
- If you forget a dose, do **not**double the next dose.
- Do not spray into eyes or mouth.
- If you get the spray in your eyes, rinse well with water.
- If allergy symptoms do not improve after one week, stop using and talk to a doctor.
- Do not share this bottle with anyone else as this may spread

g	el	'n	ns.
3		•••	


IF PUMP DOES NOT SPRAY PROPERLY, THE NOZZLE MAY BE BLOCKED


- 1. Never try to unblock nozzle with a pin or any other object (as shown in Figure E).
- 2. Clean the nozzle as shown below.



CLEANING INSTRUCTIONS

- 1. Gently pull spray nozzle away from bottle (as shown in Figure F).
- 2. Rinse **SPRAY NOZZLE ONLY** under warm water (as shown in Figure G).
- 3. Shake or tap to remove excess water.
- 4. Re-attach spray nozzle to bottle.
- 5. Press and release spray nozzle until a fine spray is produced, taking care not to spray in face.





Triamcinolone Acetonide Nasal Spraypump is now ready to use. Where can I get more information? 1-800-719-9260 Store between 20°-25°C (68°-77°F) Made in Israel Distributed By Perrigo ® Allegan, MI 49010 SIDE 2

: 44300 00 J5

Package/Label Principal Display Panel

Compare to Nasacort [®]Allergy 24HR active ingredient GOOD NEIGHBOR PHARMACY [®] Original Prescription Strength MULTI-SYMPTOM 24 Hour Nasal Allergy Spray triamcinolone acetonide nasal spray, 55 mcg per spray Allergy Symptom Reliever (Glucocorticoid)* 24 Hour Relief of: • Nasal Congestion

- Runny Nose
- Sneezing
- Itchy Nose

24 Hour

*Triamcinolone acetonide is a steroid medicine known as a glucocorticoid.

Non-Drowsy

Scent-Free

Alcohol-Free

120 Sprays

0.57 fl oz (16.9 mL)



GOOD NEIGHBOR PHARMACY 24 HOUR NASAL ALLERGY

triamcinolone acetonide spray, metered

Product Information					
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:59368-207		
Route of Administration	NASAL				

		Ingredient Name			Basis of St	trength	Strength
TRIAMCINOLO - UNII:F446C597		IDE (UNII: F446C597KA) (TRIAMCIN	OLONE ACE	TONIDE	TRIAMCINOLO ACETONIDE	-	55 ug
Inactive In	gredients						
		Ingredient Name				S	Strength
BENZALKONIU	M CHLORID	E (UNII: F5UM2KM3W7)					
CARBOXYMET	HYLCELLUL	DSE SODIUM, UNSPECIFIED (UN	III: K679OBS	5311)			
ANHYDROUS D	EXTROSE (l	JNII: 5SL0G7R0OK)					
EDETATE DISC							
		I: QTT17582CB)					
SODIUM HYDR							
		ULOSE (UNII: OP1R32D61U)					
POLYSORBATE	-)ZP39ZG8H)					
WATER (UNII: C	S9QFUKUUK)						
Product Ch	aracteris	tics					
Color		white (off white)	Sco	Score			
Shape		Size					
Flavor			Imp	orint Co	ode		
Contains							
Packaging							
# Item Code	Package Description				-		rketing d Date
1 NDC:59368-207-01	1 in 1 CARTON			-	10/24/2016		
1	120 in 1 BOTTLE, SPRAY; Type 2: Prefilled Drug Delivery Device/System (syringe, patch, etc.)						
Marketin	g Infor	mation					
	Marketing Application Number or Monograp Category Citation		raph	Marketing Start Ma Date			ting End ate
	ANDA078104			10/24/2016			

Labeler - Praxis, LLC (016329513)

Establishment					
Name	Address	ID/FEI	Business Operations		
Praxis, LLC		016329513	manufacture(59368-207) , label(59368-207) , pack(59368-207)		