PINK BISMUTH - bismuth subsalicylate tablet, chewable Discount Drug Mart, Inc

Pink Bismuth Chewable Tablets

ACTIVE INGREDIENT(in each tablet)

Bismuth subsalicylate 262 mg (total salicylate 102 mg per tablet)

PURPOSE

Upset stomach reliever and anti-diarrheal

USE(S)

relieves:

- heartburn
- indigestion
- nausea
- upset stomach
- diarrhea

WARNINGS

Reye's Syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behaviour with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's Syndrome, a rare but serious illness.

Allergy alert: Contains salicylate. Do not take if you are

- allergic to salicylates (including aspirin)
- taking other salicylate products

DO NOT USE IF YOU HAVE

- bloody or black stool
- an ulcer
- a bleeding problem

ASK A DOCTOR BEFORE USE IF YOU HAVE

- fever
- mucus in the stool

ASK A DOCTOR OR PHARMACIST BEFORE USE IF YOU ARE

taking any drug for

- anticoagulation (thinning of the blood)
- diabetes
- gout
- arthritis

WHEN USING THIS PRODUCT

a temporary, but harmless darkening of the stool and/or tongue may occur.

STOP USE AND ASK DOCTOR IF

- symptoms get worse
- ringing in the ears or loss of hearing occurs
- diarrhea lasts more than 2 days

IF PREGNANT OR BREAST-FEEDING,

ask a health professional before use.

KEEP OUT OF REACH OF CHILDREN

In case of overdose, get medical help or contact a Poison Control Center immediately.

DIRECTIONS

- chew or dissolve in mouth
- adults and children 12 years and over: 2 tablets every 1/2 to 1 hour as needed
- do not take more than 8 doses (16 tablets) in 24 hours
- use until diarrhea stops but not more than 2 days
- children under 12 years: ask a doctor
- drink plenty of clear fluids to help prevent dehydration caused diarrhea

OTHER INFORMATION

each tablet contains:

- sodium less than 1 mg
- salicylate 102 mg
- very low sodium
- avoid excessive heat (over 104°F or 40°C)
- TAMPER EVIDENT: Do not use if individual compartments are torn on open.

INACTIVE INGREDIENTS

calcium carbonate, D&C red 27 aluminum lake, flavor, magnesium stearate, mannitol, pregelatinized starch, saccharin sodium

PRINCIPAL DISPLAY PANEL

*Compare to the Active Ingredient in Pepto-Bismol®

DISCOUNT drug mart FOOD FAIR

Pink Bismuth Tablets

Soothing Relief For:

- Upset stomach
- Indigestion
- Nausea
- Heartburn
- Diarrhea

30 Chewable Tablets





Compare to the Active Ingredient in Pepto-Bismol

Soothing Relief For:

- Upset Stomach Indigestion
- Nausea Heartburn Diarrhea

30 Chewable Tablets



723396

Drug Facts

Active ingredient (in each tablet) Bismuth subsalicylate 262 mg

Purpose ..Upset stomach reliever

(total salicylate 102 mg per tablet)

and anti-diarrheal

Uses relieves: ■ diarrhea ■ heartburn ■ indigestion ■ nausea ■ upset stomach associated with these symptoms

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Contains salicylate. Do not take if you are ■ allergic to salicylates (including aspirin)

taking other salicylate products

Do not use if you have | bloody or black stool | an ulcer a bleeding problem

Ask a doctor before use if you have ■ fever ■ mucus in the stool

Ask a doctor or pharmacist before use if you are taking any drug for ■ anticoagulation (thinning of the blood)
■ diabetes
■ gout
■ arthritis

When using this product a temporary, but harmless darkening of the stool and/or tongue may occur.

Stop use and ask a doctor if symptoms get worse ringing in the ears or loss of hearing occurs

diarrhea lasts more than 2 days

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center immediately.

Directions ■ chew or dissolve in mouth ■ adults and children 12 years and over: 2 tablets every 1/2 to 1 hour as needed do not take more than 8 drink plenty of fluids to help prevent dehydration which may accompany



Other information = each tablet contains:

very low sodium = avoid excessive heat (over

Drug Facts (continued

individual compartments are torn or open.

Inactive ingredients calcium carbonate, D&C red 27 aluminum lake, flavor, magnesium stearate mannitol, pregelatinized starch, saccharin sodium.

PINK BISMUTH

bismuth subsalicylate tablet, chewable

Product Information

HUMAN OTC DRUG NDC:53943-122 **Product Type Item Code (Source)**

ORAL Route of Administration

Active Ingredient/Active Moiety

Ingredient Name Basis of Strength Strength

BISMUTH SUBSALICYLATE (UNII: 62TEY51RR1) (SALICYLIC ACID -UNII:O414PZ4LPZ)

BISMUTH SUBSALICYLATE

262 mg

Inactive Ingredients

Ingredient Name	Strength

CALCIUM CARBONATE (UNII: H0G9379FGK)

D&C RED NO. 27 ALUMINUM LAKE (UNII: ZK64F7XSTX)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MANNITOL (UNII: 3OWL53L36A)	
STARCH, CORN (UNII: O8232NY3SJ)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	

Product Characteristics			
Color	PINK	Score	no score
Shape	ROUND	Size	16mm
Flavor		Imprint Code	G122
Contains			

P	Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date	
1	NDC:53943-122- 35	5 in 1 BOX	02/26/2025		
1		6 in 1 BLISTER PACK; Type 0: Not a Combination Product			

Marketing Information				
Marketing Application Number or Monograph Category Citation		Marketing Start Date	Marketing End Date	
OTC Monograph Drug	M008	02/26/2025		

Labeler - Discount Drug Mart, Inc (047741335)

Establishment			
Name	Address	ID/FEI	Business Operations
Guardian Drug Company		119210276	MANUFACTURE(53943-122)

Revised: 2/2025 Discount Drug Mart, Inc