

**COCCI-CURE C FOR CALVES- amprolium solution**  
**Aspen Veterinary Resources**

-----  
**Cocci-Cure® C FOR CALVES**

(amprolium)

**9.6% Oral Solution Coccidiostat**

**Active Ingredient: amprolium.....9.6%**

**Restricted Drug (California) - Use Only as Directed**

Approved by FDA under ANADA # 200-482

**Net Contents: 1 Gallon (128 fl oz) (3.785 L)**

**INDICATIONS:** An aid in the treatment and prevention of coccidiosis caused by *Eimeria bovis* and *E. zuernii* in calves. For a satisfactory diagnosis a microscopic examination of the feces should be done before treatment. When treating outbreaks, drug should be administered promptly after diagnosis is determined.

**DOSAGE AND ADMINISTRATION:**

**USE DIRECTIONS:**

**5 Day Treatment Protocol**

Daily Dosage: 10 mg amprolium/kg  
(10 mg per 2.2 lb body weight)

**21 Day Prevention Protocol**

Daily Dosage: 5 mg amprolium/kg  
(5 mg per 2.2 lb body weight)

**USE DIRECTIONS:** There are **96 mg** of amprolium in every **1 mL** of Cocci-Cure® C FOR CALVES.

1 fl oz = 29.57 mL

1 fl oz = 2 measuring tablespoons

8 fl oz = ½ pint (236.56 mL)

16 fl oz = 1 pint (473.12 mL)

**IN DRINKING WATER:** Mix Cocci-Cure® C FOR CALVES 9.6% Oral Solution daily with fresh drinking water.

**5-DAY TREATMENT:** Add Cocci-Cure® C FOR CALVES 9.6% Oral Solution to drinking water at the rate of 16 fl oz/100 gal. At the usual rate of water consumption this will provide an intake of approximately 10 mg amprolium/kg (2.2 lb) body weight. Offer this solution as the only source of water for 5 days. Use on a herd basis only; when one or more calves show signs of coccidiosis, it is likely that the rest of the group has been exposed, and all calves in the group should be treated.

**21-DAY PREVENTION:** During periods of exposure or when experience indicates that coccidiosis is likely to be a hazard, add Cocci-Cure® C FOR CALVES 9.6% Oral Solution to drinking water at the rate of 8 fl oz/100 gal. At usual rates of water consumption, this

will provide an intake of approximately 5 mg amprolium/kg (2.2 lb) body weight. Offer this solution as the only source of water for 21 days.

**AS A DRENCH:** Cocci-Cure<sup>®</sup> C FOR CALVES 9.6% Oral Solution drench solutions may be stored in a clean, closed, labeled container for up to 3 days.

**5-DAY TREATMENT:** Add 3 fl oz Cocci-Cure<sup>®</sup> C FOR CALVES 9.6% Oral Solution to 1 pt of water and, with a dosage syringe, give 1 fl oz of this drench solution for each 100 lb (45 kg) body weight. This will provide a dose of approximately 10 mg amprolium/kg (2.2) body weight. Give daily for 5 days. Use on a herd basis only; when one or more calves show signs of coccidiosis, it is likely that the rest of the group has been exposed, and all calves should be treated.

**21-DAY PREVENTION:** During periods of exposure or when experience indicates that coccidiosis is likely to be a hazard, add 1 1/2 fl oz of Cocci-Cure<sup>®</sup> C FOR CALVES 9.6% Oral Solution to 1 pt of water and, with a dose syringe, give 1 fl oz of this solution for each 100 lb (45 kg) body weight. This will provide a dose of approximately 5 mg amprolium/kg (2.2 lb) body weight. Give daily for 21 days.

**RESIDUE WARNING:** Withdraw 24 hours before slaughter. A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal.

**WARNING: NOT FOR HUMAN USE. Keep this and all drugs out of reach of children.**

The Safety Data Sheet (SDS) contains more detailed occupational safety information. To report adverse drug events, for technical assistance or to obtain a copy of the SDS, contact the manufacturer at 1-800-792-1238. For additional information about adverse drug experience reporting for animal drugs, contact FDA at 1-888-FDA-VETS or online at [www.fda.gov/reportanimalae](http://www.fda.gov/reportanimalae).

**PRECAUTIONS: FOR ORAL USE IN ANIMALS ONLY. MAY CAUSE EYE IRRITATION.** For irritation, flush with plenty of water; get medical attention.

**STORAGE:** Store container in an upright position. Store between 5°C - 25°C (41°F - 77°F), with brief excursions to 40°C.

**Sodium Benzoate 0.1% added as preservative.**

**INDICATIONS:** An aid in the treatment and prevention of coccidiosis caused by *Eimeria bovis* and *E. zuernii* in calves. For a satisfactory diagnosis a microscopic examination of the feces should be done before treatment. When treating outbreaks, drug should be administered promptly after diagnosis is determined.

**DOSAGE AND ADMINISTRATION:**

**5-Day Treatment Protocol**  
Daily Dosage: 10 mg amprolium/kg (10 mg per 2.2 lb body weight)

**21-Day Prevention Protocol**  
Daily Dosage: 5 mg amprolium/kg (5 mg per 2.2 lb body weight)

**USE DIRECTIONS:** There are 96 mg of amprolium in every 1 mL of Coccid-Cure® C FOR CALVES.  
1 fl oz = 29.57 mL  
1 fl oz = 2 measuring tablespoons  
8 fl oz = 1/2 pint (236.56 mL)  
16 fl oz = 1 pint (473.12 mL)

**IN DRINKING WATER:** Mix Coccid-Cure® C FOR CALVES 9.6% Oral Solution daily with fresh drinking water.

**5-DAY TREATMENT:** Add Coccid-Cure® C FOR CALVES 9.6% Oral Solution to drinking water at the rate of 16 fl oz/100 gal. At the usual rate of water consumption this will provide an intake of approximately 10 mg amprolium/kg (2.2 lb) body weight. Offer this solution as the only source of water for 5 days. Use on a herd basis only; when one or more calves show signs of coccidiosis, it is likely that the rest of the group has been exposed, and all calves in the group should be treated.

**21-DAY PREVENTION:** During periods of exposure or when experience indicates that coccidiosis is likely to be a hazard, add Coccid-Cure® C FOR CALVES 9.6% Oral Solution to drinking water at the rate of 8 fl oz/100 gal. At usual rates of water consumption, this will provide an intake of approximately 5 mg amprolium/kg (2.2 lb) body weight. Offer this solution as the only source of water for 21 days.

Manufactured for:  
Aspen Veterinary Resources®, LTD.  
Liberty, MO 64068  
www.aspenveterinaryresources.com  
A1478  
1AMP026 8AMP015 Rev. 06/22

NDC 46066-900-01

aspEN  
VETERINARY RESOURCES, LTD.

**Cocci-Cure® C FOR CALVES**

**(amprolium)**  
9.6% Oral Solution  
Coccidiostat

Active Ingredient: amprolium.....9.6%

**Restricted Drug (California) - Use only as Directed**  
Approved by FDA under ANADA # 200-482

**AS A DRENCH:** Coccid-Cure® C FOR CALVES 9.6% Oral Solution drench solutions may be stored in a clean, closed, labeled container for up to 3 days.

**5-DAY TREATMENT:** Add 3 fl oz Coccid-Cure® C FOR CALVES 9.6% Oral Solution to 1 pt of water and, with a dose syringe, give 1 fl oz of this drench solution for each 100 lb (45 kg) body weight. This will provide a dose of approximately 10 mg amprolium/kg (2.2 lb) body weight. Give daily for 5 days. Use on a herd basis only; when one or more calves show signs of coccidiosis, it is likely that the rest of the group has been exposed, and all calves in the group should be treated.

**21-DAY PREVENTION:** During periods of exposure or when experience indicates that coccidiosis is likely to be a hazard, add 1 1/2 fl oz of Coccid-Cure® C FOR CALVES 9.6% Oral Solution to 1 pt of water and, with a dose syringe, give 1 fl oz of this drench solution for each 100 lb (45 kg) body weight. This will provide a dose of approximately 5 mg amprolium/kg (2.2 lb) body weight. Give daily for 21 days.

**RESIDUE WARNING:** Withdraw 24 hours before slaughter. A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal.

**WARNING: NOT FOR HUMAN USE. Keep this and all drugs out of reach of children.**  
The Safety Data Sheet (SDS) contains more detailed occupational safety information. To report adverse drug events, for technical assistance or to obtain a copy of the SDS, contact the manufacturer at 1-800-792-1238. For additional information about adverse drug experience reporting for animal drugs, contact FDA at 1-888-FDA-VEETS or online at www.fda.gov/reportanimalac.

**PRECAUTIONS: FOR ORAL USE IN ANIMALS ONLY. MAY CAUSE EYE IRRITATION.** For irritation, flush with plenty of water; get medical attention.

**STORAGE:** Store container in an upright position. Store between 5°C-25°C (41°F-77°F), with brief excursions to 40°C.

Sodium Benzoate 0.1% added as preservative

TAKE TIME  OBSERVE LABEL DIRECTIONS

Lot: \_\_\_\_\_ Exp: \_\_\_\_\_

Net Contents:  
1 Gallon (128 fl oz) (3.785 L)

## COCCI-CURE C FOR CALVES

amprolium solution

### Product Information

|                                |                 |                           |               |
|--------------------------------|-----------------|---------------------------|---------------|
| <b>Product Type</b>            | OTC ANIMAL DRUG | <b>Item Code (Source)</b> | NDC:46066-900 |
| <b>Route of Administration</b> | ORAL            |                           |               |

### Active Ingredient/Active Moiety

| Ingredient Name  | Basis of Strength | Strength      |
|--|-------------------|---------------|
| AMPROLIUM (UNII: 95CO6N199Q) (AMPROLIUM ION - UNII:H2T307KMZR) | AMPROLIUM ION     | 96 mg in 1 mL |

### Packaging

| # | Item Code        | Package Description | Marketing Start Date | Marketing End Date |
|---|------------------|---------------------|----------------------|--------------------|
| 1 | NDC:46066-900-01 | 3785 mL in 1 JUG    |                      |                    |

### Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|--------------------|--|----------------------|--------------------|
| ANADA              | ANADA200482                              | 12/04/2012           |                    |

**Labeler** - Aspen Veterinary Resources (627265361)

**Registrant** - Bimeda Inc. (060492923)

---

## Establishment

| Name        | Address | ID/FEI    | Business Operations |
|-------------|---------|-----------|---------------------|
| Bimeda Inc. |         | 060492923 | manufacture         |

Revised: 1/2023

Aspen Veterinary Resources