

BLOOD CLOTTING FIRST AID- benzethonium chloride and lidocaine aerosol, spray

Unifirst First Aid Corporation

Medi-First Blood Clotting Spray

Drug Facts

☐ *Active Ingredients*

Benzethonium Chloride 0.2% w/w

Lidocaine 4% w/w

Purpose

Topical antiseptic

Topical anesthetic

Uses

- for help in control of superficial bleeding
- for temporary relief of pain and to help protect against infection in minor cuts, scrapes, and burns

Warnings

For external use only

Flammable

- keep away from fire or flame
- contents under pressure
- do not puncture or incinerate container

Do not use

- in or near eyes or other mucus membranes
-
- in case of serious burns
-
- in case of deep or puncture wounds

-
- for a prolonged period of time
-
- on a large portion of the body
-
- on raw surfaces
-
- on blistered areas

Stop use and ask a doctor if

- condition worsens or symptoms persist for more than 7 days
-
- condition clears up and recurs within a few days
-
- redness, swelling, or irritation occurs

Keep out of reach of children. .

If swallowed, get medical help or contact a Poison Control Center right away

Directions

- clean the affected area
-
- shake can well before using
-
- hold 6-8 inches from surface
-
- spray until area is covered and superficial bleeding is stopped
-
- scab-like cover will gradually disappear or may be removed with soap and water
-
- for adult institutional use only
-
- not intended for use on children under 12 years of age

Other information

- avoid inhaling
- use only as directed
- intentional misuse by deliberately concentrating and inhaling the contents may be harmful or fatal
- Store at room temperature 59-86°F (15-30°C)
- do not expose to temperatures above 120°F

Inactive ingredients

isobutane, isopropyl myristate, karaya, n-butane, propane, silica, sorbitan monooleate, talc, tragacanth.

Questions or comments?

1-800-634-7680

Medi-First Blood Clotting Spray Label

Medi-First® Blood Clotting Spray

- Aids in control of bleeding
- Temporarily relieves pain of minor cuts, burns and scrapes
- Helps prevent infection
- Contains no fluorocarbons
- Aerosol
- Warning: FLAMMABLE
- Contents under pressure, read Back Panel Carefully
- Net Weight 3 oz (85g)

Drug Facts (continued)	
Directions	
<ul style="list-style-type: none"> ■ clean the affected area ■ shake can well before using ■ hold 6-8 inches from surface ■ spray until area is covered and superficial bleeding is stopped ■ scab-like cover will gradually disappear or may be removed with soap and water ■ for adult institutional use only ■ not intended for use on children under 12 years of age 	
Other information	
<ul style="list-style-type: none"> ■ avoid inhaling ■ use only as directed ■ intentional misuse by deliberately concentrating and inhaling the contents may be harmful or fatal ■ store at room temperature 59-86°F (15-30°C) ■ do not expose to temperatures above 120° F 	
Inactive ingredients	
isobutane, isopropyl myristate, karaya, n-butane, propane, silica, sorbitan monooleate, talc, tragacanth.	
Questions or comments? 1-800-634-7680	
Reorder #22617	Rev. 7/25/18
Manufactured for Medique Products Fort Myers, FL 33967 USA	

MEDI-FIRST®
Blood Clotting
Spray

- Aids in control of superficial bleeding
- Temporarily relieves pain of minor cuts, burns and scrapes
- Helps prevent infection
- Contains no fluorocarbons

Aerosol

WARNING: FLAMMABLE
CONTENTS UNDER PRESSURE, READ BACK PANEL CAREFULLY

NET WEIGHT 3 OZ (85g)

Drug Facts	
Active ingredients	Purposes
Benzethonium Chloride 0.2% w/w.....	Topical antiseptic
Lidocaine 4% w/w.....	Topical anesthetic
Uses	
<ul style="list-style-type: none"> ■ for help in control of superficial bleeding ■ for temporary relief of pain and to help protect against infection in minor cuts, scrapes and minor burns 	
Warnings	
For external use only.	
Flammable	
<ul style="list-style-type: none"> ■ keep away from fire or flame ■ contents under pressure ■ do not puncture or incinerate container 	
Do not use	
<ul style="list-style-type: none"> ■ in or near eyes or other mucous membranes ■ in case of serious burns ■ in case of deep or puncture wounds ■ for a prolonged period of time ■ on a large portion of the body ■ on raw surfaces ■ on blistered areas 	
Stop use and ask a doctor if	
<ul style="list-style-type: none"> ■ condition worsens or symptoms persist for more than 7 days ■ condition clears up and recurs within a few days ■ redness, swelling, or irritation occurs 	
Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away. ▶	

BLOOD CLOTTING FIRST AID

benzethonium chloride and lidocaine aerosol, spray

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:47682-226
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
LIDOCAINE (UNII: 98PI200987) (LIDOCAINE - UNII:98PI200987)	LIDOCAINE	3400 mg in 85 g
BENZETHONIUM CHLORIDE (UNII: PH41D05744) (BENZETHONIUM - UNII:1VU15B70BP)	BENZETHONIUM CHLORIDE	170 mg in 85 g

Inactive Ingredients

Ingredient Name	Strength
ISOBUTANE (UNII: BXR49TP611)	
ISOPROPYL MYRISTATE (UNII: ORE8K4LNJS)	
KARAYA GUM (UNII: 73W9IQY50Q)	
BUTANE (UNII: 6LV4FOR43R)	
PROPANE (UNII: T75W9911L6)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
SORBITAN MONOOLEATE (UNII: 06XEA2VD56)	
TALC (UNII: 7SEV7J4R1U)	
TRAGACANTH (UNII: 2944357O2O)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:47682-226-17	85 g in 1 BOTTLE, SPRAY; Type 0: Not a Combination Product	04/05/2013	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M003	04/05/2013	

Labeler - Unifirst First Aid Corporation (832947092)**Establishment**

Name	Address	ID/FEI	Business Operations
Dixon Investments, Inc.		115315822	manufacture(47682-226)

Revised: 9/2025

Unifirst First Aid Corporation