

# **IVABRADINE- ivabradine tablet, film coated**

## **Ingenus Pharmaceuticals, LLC**

-----

### **HIGHLIGHTS OF PRESCRIBING INFORMATION**

**These highlights do not include all the information needed to use IVABRADINE TABLETS safely and effectively. See full prescribing information for IVABRADINE TABLETS IVABRADINE tablets, for oral use**  
**Initial U.S. Approval: 2015**

### **INDICATIONS AND USAGE**

Ivabradine is a hyperpolarization-activated cyclic nucleotide-gated channel blocker indicated:

- To reduce the risk of hospitalization for worsening heart failure in adult patients with stable, symptomatic chronic heart failure with reduced left ventricular ejection fraction. (1.1)

### **DOSAGE AND ADMINISTRATION**

#### **Adult patients**

● Starting dose is 2.5 (vulnerable adults) or 5 mg twice daily with food. After 2 weeks of treatment, adjust dose based on heart rate. The maximum dose is 7.5 mg twice daily. (2.1)

### **DOSAGE FORMS AND STRENGTHS**

- Tablets: 5 mg, 7.5 mg (3)

### **CONTRAINDICATIONS**

- Acute decompensated heart failure (4)
- Clinically significant hypotension (4)
- Sick sinus syndrome, sinoatrial block or 3<sup>rd</sup>degree AV block, unless a functioning demand pacemaker is present (4)
- Clinically significant bradycardia (4)
- Severe hepatic impairment (4)
- Heart rate maintained exclusively by the pacemaker (4)
- In combination with strong cytochrome CYP3A4 inhibitors (4)

### **WARNINGS AND PRECAUTIONS**

- Fetal toxicity: Females should use effective contraception. (5.1)
- Monitor patients for atrial fibrillation. (5.2)
- Monitor heart rate decreases and bradycardia symptoms during treatment. (5.3)
- Not recommended in patients with 2<sup>nd</sup>degree AV block. (5.3)

### **ADVERSE REACTIONS**

Most common adverse reactions occurring in  $\geq 1\%$  of patients are bradycardia, hypertension, atrial fibrillation and luminous phenomena (phosphenes). (6)

**To report SUSPECTED ADVERSE REACTIONS, contact Ingenus Pharmaceuticals, LLC at 1-877-748-1970 or FDA at 1-800-FDA-1088 or [www.fda.gov/medwatch](http://www.fda.gov/medwatch).**

### **DRUG INTERACTIONS**

- Avoid CYP3A4 inhibitors or inducers. (7.1)
- Negative chronotropes increase risk of bradycardia; monitor heart rate. (7.2)

### **USE IN SPECIFIC POPULATIONS**

- Lactation: Breastfeeding not recommended. (8.2)

**See 17 for PATIENT COUNSELING INFORMATION and Medication Guide.**

**Revised: 9/2023**

## **1 INDICATIONS AND USAGE**

1.1 Heart Failure in Adult Patients

## **2 DOSAGE AND ADMINISTRATION**

2.1 Adults

## **3 DOSAGE FORMS AND STRENGTHS**

## **4 CONTRAINDICATIONS**

## **5 WARNINGS AND PRECAUTIONS**

5.1 Fetal Toxicity

5.2 Atrial Fibrillation

5.3 Bradycardia and Conduction Disturbances

## **6 ADVERSE REACTIONS**

6.1 Clinical Trials Experience

6.2 **Post marketing Experience**

## **7 DRUG INTERACTIONS**

7.1 Cytochrome P450-Based Interactions

7.2 Negative Chronotropes

7.3 Pacemakers in Adults

## **8 USE IN SPECIFIC POPULATIONS**

8.1 Pregnancy

8.2 Lactation

8.3 Females and Males of Reproductive Potential

8.4 Pediatric Use

8.5 Geriatric Use

8.6 Hepatic Impairment

8.7 Renal Impairment

## **10 OVERDOSAGE**

## **11 DESCRIPTION**

## **12 CLINICAL PHARMACOLOGY**

12.1 Mechanism of Action

12.2 Pharmacodynamics

12.3 Pharmacokinetics

## **13 NONCLINICAL TOXICOLOGY**

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

13.2 Animal Toxicology and/or Pharmacology

## **14 CLINICAL STUDIES**

14.1 Heart Failure in Adult Patients

## **16 HOW SUPPLIED/STORAGE AND HANDLING**

## **17 PATIENT COUNSELING INFORMATION**

\* Sections or subsections omitted from the full prescribing information are not listed.

---

## **FULL PRESCRIBING INFORMATION**

### **1 INDICATIONS AND USAGE**

#### **1.1 Heart Failure in Adult Patients**

Ivabradine tablets are indicated to reduce the risk of hospitalization for worsening heart failure in adult patients with stable, symptomatic chronic heart failure with left ventricular ejection fraction  $\leq 35\%$ , who are in sinus rhythm with resting heart rate  $\geq 70$  beats per minute and either are on maximally tolerated doses of beta-blockers or have a contraindication to beta-blocker use.

## 2 DOSAGE AND ADMINISTRATION

### 2.1 Adults

The recommended starting dose of ivabradine tablets is 5 mg twice daily with food. Assess patient after two weeks and adjust dose to achieve a resting heart rate between 50 and 60 beats per minute (bpm) as shown in Table 1. Thereafter, adjust dose as needed based on resting heart rate and tolerability. The maximum dose is 7.5 mg twice daily. In adult patients unable to swallow tablets, Corlanor oral solution can be used [see *Clinical Pharmacology (12.3)*].

In patients with a history of conduction defects or other patients in whom bradycardia could lead to hemodynamic compromise, initiate therapy at 2.5 mg twice daily before increasing the dose based on heart rate [see *Warnings and Precautions (5.3)*].

**Table 1. Dose Adjustment for Adults**

Heart Rate	Dose Adjustment
> 60 bpm	Increase dose by 2.5 mg (given twice daily) up to a maximum dose of 7.5 mg twice daily
50-60 bpm	Maintain dose
< 50 bpm or signs and symptoms of bradycardia	Decrease dose by 2.5 mg (given twice daily); if current dose is 2.5 mg twice daily, discontinue therapy*

\*[see *Warnings and Precautions (5.3)*]

## 3 DOSAGE FORMS AND STRENGTHS

### Tablets

Ivabradine tablets 5 mg: Light salmon-colored, capsule-shaped, film-coated scored tablet debossed with "15" divided by score on one side and plain on other side. The tablet is scored and can be divided into equal halves to provide a 2.5 mg dose.

Ivabradine tablets 7.5 mg: Light salmon-colored, round-shaped, film-coated tablet debossed with "1" on one side & 7.5 on other side.

## 4 CONTRAINDICATIONS

Ivabradine tablets are contraindicated in patients with:

- Acute decompensated heart failure
- Clinically significant hypotension
- Sick sinus syndrome, sinoatrial block or 3<sup>rd</sup> degree AV block, unless a functioning

demand pacemaker is present

- Clinically significant bradycardia [see *Warnings and Precautions (5.3)*]
- Severe hepatic impairment [see *Use in Specific Populations (8.6)*]
- Pacemaker dependence (heart rate maintained exclusively by the pacemaker) [see *Drug Interactions (7.3)*]
- Concomitant use of strong cytochrome P450 3A4 (CYP3A4) inhibitors [see *Drug Interactions (7.1)*]

## **5 WARNINGS AND PRECAUTIONS**

### **5.1 Fetal Toxicity**

Ivabradine tablets may cause fetal toxicity when administered to a pregnant woman based on findings in animal studies. Embryo-fetal toxicity and cardiac teratogenic effects were observed in fetuses of pregnant rats treated during organogenesis at exposures 1 to 3 times the human exposures ( $AUC_{0-24hr}$ ) at the maximum recommended human dose (MRHD) [see *Use in Specific Populations (8.1)*]. Advise females of reproductive potential to use effective contraception when taking ivabradine tablets [see *Use in Specific Populations (8.3)*].

### **5.2 Atrial Fibrillation**

Ivabradine tablets increases the risk of atrial fibrillation. In the Systolic Heart Failure Treatment with the I<sub>f</sub> Inhibitor Ivabradine Trial (SHIFT), the rate of atrial fibrillation was 5.0% per patient-year in patients treated with ivabradine tablets and 3.9% per patient-year in patients treated with placebo [see *Clinical Studies (14)*]. Regularly monitor cardiac rhythm. Discontinue ivabradine tablets if atrial fibrillation develops.

### **5.3 Bradycardia and Conduction Disturbances**

#### Adult Patients

Bradycardia, sinus arrest, and heart block have occurred with ivabradine tablets. The rate of bradycardia was 6.0% per patient-year in patients treated with ivabradine tablets (2.7% symptomatic; 3.4% asymptomatic) and 1.3% per patient-year in patients treated with placebo. Risk factors for bradycardia include sinus node dysfunction, conduction defects (e.g., 1<sup>st</sup> or 2<sup>nd</sup> degree atrioventricular block, bundle branch block), ventricular dyssynchrony, and use of other negative chronotropes (e.g., digoxin, diltiazem, verapamil, amiodarone). Bradycardia may increase the risk of QT prolongation which may lead to severe ventricular arrhythmias, including torsade de pointes, especially in patients with risk factors such as use of QTc prolonging drugs [see *Adverse Reactions (6.2)*].

Concurrent use of verapamil or diltiazem will increase ivabradine tablets exposure, may themselves contribute to heart rate lowering, and should be avoided [see *Clinical Pharmacology (12.3)*]. Avoid use of ivabradine tablets in patients with 2<sup>nd</sup> degree atrioventricular block unless a functioning demand pacemaker is present [see *Contraindications (4)*].

## **6 ADVERSE REACTIONS**

Clinically significant adverse reactions that appear in other sections of the labeling include:

- Atrial Fibrillation [see *Warnings and Precautions (5.2)*]
- Bradycardia and Conduction Disturbances [see *Warnings and Precautions (5.3)*]

## 6.1 Clinical Trials Experience

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in practice.

### Adult Patients with Heart Failure

In SHIFT, safety was evaluated in 3,260 patients treated with ivabradine tablets and 3,278 patients given placebo. The median duration of ivabradine tablets exposure was 21.5 months.

The most common adverse drug reactions in the SHIFT trial are shown in Table 2 [see *Warnings and Precautions (5.2), (5.3)*].

**Table 2. Adverse Drug Reactions with Rates  $\geq$  1.0% Higher on Ivabradine than Placebo occurring in  $>$  1% on Ivabradine in SHIFT**

	<b>Ivabradine N = 3,260</b>	<b>Placebo N = 3,278</b>
Bradycardia	10 %	2.2 %
Hypertension, Blood Pressure Increased	8.9 %	7.8 %
Atrial fibrillation	8.3 %	6.6 %
Phosphenes, visual brightness	2.8 %	0.5 %

### Luminous Phenomena (Phosphenes)

Phosphenes are phenomena described as a transiently enhanced brightness in a limited area of the visual field, halos, image decomposition (stroboscopic or kaleidoscopic effects), colored bright lights, or multiple images (retinal persistency). Phosphenes are usually triggered by sudden variations in light intensity. Ivabradine tablets can cause phosphenes, thought to be mediated through ivabradine tablets effects on retinal photoreceptors [see *Clinical Pharmacology (12.1)*]. Onset is generally within the first 2 months of treatment, after which they may occur repeatedly. Phosphenes were generally reported to be of mild to moderate intensity and led to treatment discontinuation in  $<$  1% of patients; most resolved during or after treatment.

## 6.2 Post marketing Experience

Because these reactions are reported voluntarily from a population of uncertain size, it is not always possible to estimate their frequency reliably or establish a causal relationship to drug exposure.

The following adverse reactions have been identified in adults during post-approval use of ivabradine tablets: syncope, hypotension, torsade de pointes, ventricular fibrillation,

ventricular tachycardia, angioedema, erythema, rash, pruritus, urticaria, vertigo, and diplopia, and visual impairment.

## **7 DRUG INTERACTIONS**

### **7.1 Cytochrome P450-Based Interactions**

Ivabradine is primarily metabolized by CYP3A4. Concomitant use of CYP3A4 inhibitors increases ivabradine plasma concentrations and use of CYP3A4 inducers decreases them. Increased plasma concentrations may exacerbate bradycardia and conduction disturbances.

The concomitant use of strong CYP3A4 inhibitors is contraindicated [see *Contraindications (4) and Clinical Pharmacology (12.3)*]. Examples of strong CYP3A4 inhibitors include azole antifungals (e.g., itraconazole), macrolide antibiotics (e.g., clarithromycin, telithromycin), HIV protease inhibitors (e.g. nelfinavir), and nefazodone.

Avoid concomitant use of moderate CYP3A4 inhibitors when using ivabradine tablets. Examples of moderate CYP3A4 inhibitors include diltiazem, verapamil, and grapefruit juice [see *Warnings and Precautions (5.3) and Clinical Pharmacology (12.3)*].

Avoid concomitant use of CYP3A4 inducers when using ivabradine tablets. Examples of CYP3A4 inducers include St. John's wort, rifampicin, barbiturates, and phenytoin [see *Clinical Pharmacology (12.3)*].

### **7.2 Negative Chronotropes**

Most patients receiving ivabradine tablets will also be treated with a beta-blocker. The risk of bradycardia increases with concomitant administration of drugs that slow heart rate (e.g., digoxin, amiodarone, beta-blockers). Monitor heart rate in patients taking ivabradine tablets with other negative chronotropes.

### **7.3 Pacemakers in Adults**

Ivabradine tablets dosing is based on heart rate reduction, targeting a heart rate of 50 to 60 beats per minute in adults [see *Dosage and Administration (2.1)*]. Patients with demand pacemakers set to a rate  $\geq 60$  beats per minute cannot achieve a target heart rate  $< 60$  beats per minute, and these patients were excluded from clinical trials [see *Clinical Studies (14.1)*]. The use of ivabradine tablets is not recommended in patients with demand pacemakers set to rates  $\geq 60$  beats per minute.

## **8 USE IN SPECIFIC POPULATIONS**

### **8.1 Pregnancy**

#### Risk Summary

Based on findings in animals, ivabradine tablets may cause fetal harm when administered to a pregnant woman. There are no adequate and well-controlled studies of ivabradine tablets in pregnant women to inform any drug-associated risks. In animal reproduction studies, oral administration of ivabradine to pregnant rats during organogenesis at a dosage providing 1 to 3 times the human exposure ( $AUC_{0-24hr}$ ) at the MRHD resulted in

embryo-fetal toxicity and teratogenicity manifested as abnormal shape of the heart, interventricular septal defect, and complex anomalies of primary arteries. Increased post-natal mortality was associated with these teratogenic effects in rats. In pregnant rabbits, increased post-implantation loss was noted at an exposure ( $AUC_{0-24hr}$ ) 5 times the human exposure at the MRHD. Lower doses were not tested in rabbits. The background risk of major birth defects for the indicated population is unknown. The estimated background risk of major birth defects in the U.S. general population is 2 to 4%, however, and the estimated risk of miscarriage is 15 to 20% in clinically recognized pregnancies. Advise a pregnant woman of the potential risk to the fetus.

### Clinical Considerations

#### *Disease-associated Maternal and/or Embryo-fetal Risk*

Stroke volume and heart rate increase during pregnancy, increasing cardiac output, especially during the first trimester. Pregnant patients with left ventricular ejection fraction less than 35% on maximally tolerated doses of beta-blockers may be particularly heart rate dependent for augmenting cardiac output. Therefore, pregnant patients who are started on ivabradine tablets, especially during the first trimester, should be followed closely for destabilization of their congestive heart failure that could result from heart rate slowing.

Monitor pregnant women with chronic heart failure in 3<sup>rd</sup> trimester of pregnancy for preterm birth.

### Data

#### *Animal Data*

In pregnant rats, oral administration of ivabradine during the period of organogenesis (gestation day 6-15) at doses of 2.3, 4.6, 9.3, or 19 mg/kg/day resulted in fetal toxicity and teratogenic effects. Increased intrauterine and post-natal mortality and cardiac malformations were observed at doses  $\geq 2.3$  mg/kg/day (equivalent to the human exposure at the MRHD based on  $AUC_{0-24hr}$ ). Teratogenic effects including interventricular septal defect and complex anomalies of major arteries were observed at doses  $\geq 4.6$  mg/kg/day (approximately 3 times the human exposure at the MRHD based on  $AUC_{0-24hr}$ ).

In pregnant rabbits, oral administration of ivabradine during the period of organogenesis (gestation day 6-18) at doses of 7, 14, or 28 mg/kg/day resulted in fetal toxicity and teratogenicity. Treatment with all doses  $\geq 7$  mg/kg/day (equivalent to the human exposure at the MRHD based on  $AUC_{0-24hr}$ ) caused an increase in post-implantation loss. At the high dose of 28 mg/kg/day (approximately 15 times the human exposure at the MRHD based on  $AUC_{0-24hr}$ ), reduced fetal and placental weights were observed, and evidence of teratogenicity (ectrodactylia observed in 2 of 148 fetuses from 2 of 18 litters) was demonstrated.

In the pre- and post-natal study, pregnant rats received oral administration of ivabradine at doses of 2.5, 7, or 20 mg/kg/day from gestation day 6 to lactation day 20. Increased post-natal mortality associated with cardiac teratogenic findings was observed in the F1 pups delivered by dams treated at the high dose (approximately 15 times the human exposure at the MRHD based on  $AUC_{0-24hr}$ ).

## **8.2 Lactation**

## Risk Summary

There is no information regarding the presence of ivabradine in human milk, the effects of ivabradine on the breastfed infant, or the effects of the drug on milk production. Animal studies have shown, however, that ivabradine is present in rat milk [see *Data*]. Because of the potential risk to breastfed infants from exposure to ivabradine tablets, breastfeeding is not recommended.

## Data

Lactating rats received daily oral doses of [14C]-ivabradine (7 mg/kg) on post-parturition days 10 to 14; milk and maternal plasma were collected at 0.5 and 2.5 hours post-dose on day 14. The ratios of total radio activity associated with [14C]-ivabradine or its metabolites in milk vs. plasma were 1.5 and 1.8, respectively, indicating that ivabradine is transferred to milk after oral administration.

## **8.3 Females and Males of Reproductive Potential**

### Contraception

#### *Females*

Ivabradine tablets may cause fetal harm, based on animal data. Advise females of reproductive potential to use effective contraception during ivabradine tablets treatment [see *Use in Specific Populations (8.1)*].

## **8.4 Pediatric Use**

The safety and efficacy of ivabradine tablets have not been established in patients less than 6 months of age.

## **8.5 Geriatric Use**

No pharmacokinetic differences have been observed in elderly ( $\geq 65$  years) or very elderly ( $\geq 75$  years) patients compared to the overall population. However, ivabradine tablets has only been studied in a limited number of patients  $\geq 75$  years of age.

## **8.6 Hepatic Impairment**

No dose adjustment is required in patients with mild or moderate hepatic impairment. Ivabradine tablets is contraindicated in patients with severe hepatic impairment (Child-Pugh C) as it has not been studied in this population and an increase in systemic exposure is anticipated [see *Contraindications (4) and Clinical Pharmacology (12.3)*].

## **8.7 Renal Impairment**

No dosage adjustment is required for patients with creatinine clearance 15 to 60 mL/min. No data are available for patients with creatinine clearance below 15 mL/min [see *Clinical Pharmacology (12.3)*].

## **10 OVERDOSAGE**

Overdose may lead to severe and prolonged bradycardia. In the event of bradycardia with poor hemodynamic tolerance, temporary cardiac pacing may be required.

Supportive treatment, including intravenous (IV) fluids, atropine, and intravenous beta-stimulating agents such as isoproterenol, may be considered.


## 11 DESCRIPTION

Ivabradine tablets contains ivabradine as the active pharmaceutical ingredient. Ivabradine is a hyperpolarization-activated cyclic nucleotide-gated channel blocker that reduces the spontaneous pacemaker activity of the cardiac sinus node by selectively inhibiting the  $I_f$  current, resulting in heart rate reduction with no effect on ventricular repolarization and no effects on myocardial contractility.

The chemical name for ivabradine hydrochloride is:

3-(3-(((7S)-3,4-Dimethoxybicyclo[4.2.0]octa-1,3,5-trien-7-yl)methyl) methyl amino} propyl)-1,3,4,5-tetrahydro-7,8-dimethoxy-2H-3-benzazepin-2-one,hydrochloride. The molecular formula is  $C_{27}H_{36}N_2O_5 \cdot HCl$ , and the molecular weight (free base +HCl) is 505.1 (468.6 + 36.5). The chemical structure of ivabradine is shown in Figure 1.

**Figure 1. Chemical Structure of Ivabradine**



### Tablets

Ivabradine tablets are supplied in 5 mg and 7.5 mg tablets for oral administration. The tablets contain 5 mg and 7.5 mg of ivabradine, as active ingredient, equivalent to 5.39 mg and 8.09 mg of ivabradine hydrochloride, respectively. The tablets contain the following inactive ingredients: lactose monohydrate, maltodextrin, maize starch b, colloidal silicon dioxide, magnesium stearate, hypromellose, titanium dioxide, polyethylene glycol 4000, yellow iron oxide and red iron oxide.

## 12 CLINICAL PHARMACOLOGY

### 12.1 Mechanism of Action

Ivabradine tablets blocks the hyperpolarization-activated cyclic nucleotide-gated (HCN) channel responsible for the cardiac pacemaker  $I_f$  current, which regulates heart rate. In clinical electrophysiology studies, the cardiac effects were most pronounced in the sinoatrial (SA) node, but prolongation of the AH interval has occurred as has PR interval prolongation. There was no effect on ventricular repolarization and no effects on myocardial contractility [see *Clinical Pharmacology (12.2)*].

Ivabradine tablets can also inhibit the retinal current  $I_h$ .  $I_h$  is involved in curtailing retinal responses to bright light stimuli. Under triggering circumstances (e.g., rapid changes in luminosity), partial inhibition of  $I_h$  by ivabradine tablets may underlie the luminous

phenomena experienced by patients. Luminous phenomena (phosphenes) are described as a transient enhanced brightness in a limited area of the visual field [see *Adverse Reactions (6.1)*].

## 12.2 Pharmacodynamics

Ivabradine tablets causes a dose-dependent reduction in heart rate. The size of the effect is dependent on the baseline heart rate (i.e., greater heart rate reduction occurs in subjects with higher baseline heart rate). At recommended doses, heart rate reduction is approximately 10 bpm at rest and during exercise. Analysis of heart rate reduction vs. dose indicates a plateau effect at doses > 20 mg twice daily. In a study of subjects with preexisting conduction system disease (first- or second-degree AV block or left or right bundle branch block) requiring electrophysiologic study, IV ivabradine (0.20 mg/kg) administration slowed the overall heart rate by approximately 15 bpm, increased the PR interval (29 msec), and increased the AH interval (27 msec).

Ivabradine tablets does not have negative inotropic effects. Ivabradine increases the uncorrected QT interval with heart rate slowing but does not cause rate-corrected prolongation of QT.

## 12.3 Pharmacokinetics

The peak concentration ( $C_{max}$ ) and area under the plasma concentration time curve (AUC) are similar for ivabradine and S 18982 between oral solution and tablets for the same dose.

### Absorption and Bioavailability

Following oral administration, peak plasma ivabradine concentrations are reached in approximately 1 hour under fasting conditions. The absolute oral bioavailability of ivabradine is approximately 40% because of first-pass elimination in the gut and liver.

Food delays absorption by approximately 1 hour and increases plasma exposure by 20% to 40%. Ivabradine tablets should be taken with food [see *Dosage and Administration (2)*].

Ivabradine is approximately 70% plasma protein bound, and the volume of distribution at steady state is approximately 100 L.

### Metabolism and Excretion

The pharmacokinetics of ivabradine are linear over an oral dose range of 0.5 mg to 24 mg. Ivabradine is extensively metabolized in the liver and intestines by CYP3A4-mediated oxidation. The major metabolite is the N-desmethylated derivative (S 18982), which is equipotent to ivabradine and circulates at concentrations approximately 40% that of ivabradine. The N-desmethylated derivative is also metabolized by CYP3A4. Ivabradine plasma levels decline with a distribution half-life of 2 hours and an effective half-life of approximately 6 hours.


The total clearance of ivabradine is 24 L/h, and renal clearance is approximately 4.2 L/h, with ~ 4% of an oral dose excreted unchanged in urine. The excretion of metabolites occurs to a similar extent via feces and urine.

### Drug Interactions

The effects of coadministered drugs (CYP3A4 inhibitors, substrates, inducers, and other

concomitantly administered drugs) on the pharmacokinetics of ivabradine tablets were studied in several single- and multiple dose studies. Pharmacokinetic measures indicating the magnitude of these interactions are presented in Figure 2.

**Figure 2. Impact of Coadministered Drugs on the Pharmacokinetics of Ivabradine tablets**



Digoxin exposure did not change when concomitantly administered with ivabradine. No dose adjustment is required when ivabradine is concomitantly administered with digoxin.

### Effect of Ivabradine on Metformin Pharmacokinetics

Ivabradine, dosed at 10 mg twice daily to steady state, did not affect the pharmacokinetics of metformin (an organic cation transporter [OCT2] sensitive substrate). The geometric mean (90% confidence interval [CI]) ratios of  $C_{max}$  and  $AUC_{inf}$  of metformin, with and without ivabradine were 0.98 [0.83–1.15] and 1.02 [0.86–1.22], respectively. No dose adjustment is required for metformin when administered with ivabradine tablets.

### Specific Populations

#### *Age*

No pharmacokinetic differences ( $AUC$  or  $C_{max}$ ) have been observed between elderly ( $\geq 65$  years) or very elderly ( $\geq 75$  years) patients and the overall patient population [see *Use in Specific Populations (8.5)*].

#### *Hepatic impairment*

In patients with mild (Child-Pugh A) and moderate (Child-Pugh B) hepatic impairment, the pharmacokinetics of ivabradine tablets were similar to that in patients with normal hepatic function. No data are available in patients with severe hepatic impairment (Child-Pugh C) [see *Contraindications (4)*].

#### *Renal impairment*

Renal impairment (creatinine clearance from 15 to 60 mL/min) has minimal effect on the

pharmacokinetics of ivabradine tablets. No data are available for patients with creatinine clearance below 15 mL/min.

## 13 NONCLINICAL TOXICOLOGY

### 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

There was no evidence of carcinogenicity when mice and rats received ivabradine up to 104 weeks by dietary administration. High doses in these studies were associated with mean ivabradine exposures of at least 37 times higher than the human exposure ( $AUC_{0-24hr}$ ) at the MRHD.

Ivabradine tested negative in the following assays: bacterial reverse mutation (Ames) assay, *in vivo* bone marrow micronucleus assay in both mouse and rat, *in vivo* chromosomal aberration assay in rats, and *in vivo* unscheduled DNA synthesis assay in rats. Results of the *in vitro* chromosomal aberration assay were equivocal at concentrations approximately 1,500 times the human  $C_{max}$  at the MRHD. Ivabradine tested positive in the mouse lymphoma assays and *in vitro* unscheduled DNA synthesis assay in rat hepatocytes at concentrations greater than 1,500 times the human  $C_{max}$  at the MRHD.

Reproduction toxicity studies in animals demonstrated that ivabradine did not affect fertility in male or female rats at exposures 46 to 133 times the human exposure ( $AUC_{0-24hr}$ ) at the MRHD.

### 13.2 Animal Toxicology and/or Pharmacology

Reversible changes in retinal function were observed in dogs administered oral ivabradine at total doses of 2, 7, or 24 mg/kg/day (approximately 0.6 to 50 times the human exposure at the MRHD based on  $AUC_{0-24hr}$ ) for 52 weeks. Retinal function assessed by electroretinography demonstrated reductions in cone system responses, which reversed within a week post-dosing, and were not associated with damage to ocular structures as evaluated by light microscopy. These data are consistent with the pharmacological effect of ivabradine related to its interaction with hyperpolarization-activated  $I_h$  currents in the retina, which share homology with the cardiac pacemaker  $I_f$  current.

## 14 CLINICAL STUDIES

### 14.1 Heart Failure in Adult Patients

#### SHIFT

The Systolic Heart failure treatment with the  $I_f$  inhibitor ivabradine Trial (SHIFT) was a randomized, double-blind trial comparing ivabradine tablets and placebo in 6558 adult patients with stable New York Heart Association (NYHA) class II to IV heart failure, left ventricular ejection fraction  $\leq 35\%$ , and resting heart rate  $\geq 70$  bpm. Patients had to have been clinically stable for at least 4 weeks on an optimized and stable clinical regimen, which included maximally tolerated doses of beta-blockers and, in most cases, ACE inhibitors or ARBs, spironolactone, and diuretics, with fluid retention and symptoms of congestion minimized. Patients had to have been hospitalized for heart failure within

12 months prior to study entry.

The underlying cause of CHF was coronary artery disease in 68% of patients. At baseline, approximately 49% of randomized subjects were NYHA class II, 50% were NYHA class III, and 2% were NYHA class IV. The mean left ventricular ejection fraction was 29%. All subjects were initiated on ivabradine tablets 5 mg (or matching placebo) twice daily and the dose was increased to 7.5 mg twice daily or decreased to 2.5 mg twice daily to maintain the resting heart rate between 50 and 60 bpm, as tolerated. The primary endpoint was a composite of the first occurrence of either hospitalization for worsening heart failure or cardiovascular death.

Most patients (89%) were taking beta-blockers, with 26% on guideline-defined target daily doses. The main reasons for not receiving the target beta-blocker doses at baseline were hypotension (45% of patients not at target), fatigue (32%), dyspnea (14%), dizziness (12%), history of cardiac decompensation (9%), and bradycardia (6%). For the 11% of patients not receiving any beta-blocker at baseline, the main reasons were chronic obstructive pulmonary disease, hypotension, and asthma. Most patients were also taking ACE inhibitors and/or angiotensin II antagonists (91%), diuretics (83%), and anti-aldosterone agents (60%). Few patients had an implantable cardioverter-defibrillator (ICD) (3.2%) or a cardiac resynchronization therapy (CRT) device (1.1%). Median follow-up was 22.9 months. At 1 month, 63%, 26%, and 8% of ivabradine tablets -treated patients were taking 7.5, 5, and 2.5 mg BID, whereas 3% had withdrawn from the drug, primarily for bradycardia.

SHIFT demonstrated that ivabradine tablets reduced the risk of the combined endpoint of hospitalization for worsening heart failure or cardiovascular death based on a time-to-event analysis (hazard ratio: 0.82, 95% confidence interval [CI]: 0.75, 0.90,  $p < 0.0001$ ) (Table 3). The treatment effect reflected only a reduction in the risk of hospitalization for worsening heart failure; there was no favorable effect on the mortality component of the primary endpoint. In the overall treatment population, ivabradine tablets had no statistically significant benefit on cardiovascular death.

**Table 3. SHIFT - Incidence of the Primary Composite Endpoint and Components**

Endpoint	Ivabradine tablets (N = 3,241) %			Placebo (N = 3,264) %			Hazard	
	n	%	PY	n	%	PY	Ratio	[95%CI] p-value
Primary composite endpoint: Time to first hospitalization for worsening heart failure or cardiovascular death <sup>a</sup>	793	24.5	14.5	937	28.7	17.7	0.82	[0.75 , < 0.90] 0.0001
Hospitalization for worsening	505	15.6	9.2	660	20.2	12.5		

heart failure Cardiovascular death as first event	288	8.9	4.8	277	8.5	4.7		
Subjects with events at any Time								
Hospitalization for worsening heart failure <sup>b</sup>	514	15.9	9.4	672	20.6	12.7	0.74	[0.66 , 0.83]
Cardiovascular death <sup>b</sup>	449	13.9	7.5	491	15.0	8.3	0.91	[0.80 , 1.03]

<sup>a</sup> Patients who died on the same calendar day as their first hospitalization for worsening heart

failure are counted under cardiovascular death.

<sup>b</sup> Analyses of the components of the primary composite endpoint were not prospectively planned to be adjusted for multiplicity.


N: number of patients at risk; n: number of patients having experienced the endpoint; %: incidence rate =  $(n/N) \times 100$ ; % PY: annual incidence rate =  $(n/\text{number of patient-years}) \times 100$ ; CI: Confidence interval

The hazard ratio between treatment groups (ivabradine/placebo) was estimated based on an adjusted Cox proportional hazards model with beta-blocker intake at randomization (yes/no) as

a covariate; p-value: Wald test

The Kaplan-Meier curve (Figure 3) shows time to first occurrence of the primary composite endpoint of hospitalization for worsening heart failure or cardiovascular death in the overall study.

**Figure 3. SHIFT: Time to First Event of Primary Composite Endpoint**




Number of Subjects at Risk:

Placebo	3264	2868	2489	2061	1089	439	17
Corlanor	3241	2928	2600	2173	1191	447	16

A wide range of demographic characteristics, baseline disease characteristics, and baseline concomitant medications were examined for their influence on outcomes. Many of these results are shown in Figure 4. Such analyses must be interpreted cautiously, as differences can reflect the play of chance among a large number of analyses.

Most of the results show effects consistent with the overall study result. Ivabradine tablets benefit on the primary endpoint in SHIFT appeared to decrease as the dose of beta-blockers increased, with little if any benefit demonstrated in patients taking guideline-defined target doses of beta-blockers.

**Figure 4. Effect of Treatment on Primary Composite Endpoint in Subgroups**



Note: The figure above presents effects in various subgroups, all of which are baseline characteristics. The 95% confidence limits that are shown do not take into account the number of comparisons made and may not reflect the effect of a particular factor after adjustment for all other factors. Apparent homogeneity or heterogeneity among groups

should not be over-interpreted.

### BEAUTIFUL and SIGNIFY: No benefit in stable coronary artery disease with or without stable heart failure

The Morbidity-mortality Evaluation of the I<sub>f</sub> Inhibitor Ivabradine in Patients with Coronary Disease and Left Ventricular Dysfunction Trial (BEAUTIFUL) was a randomized, double-blind, placebo-controlled trial in 10,917 adult patients with coronary artery disease, impaired left ventricular systolic function (ejection fraction <40%) and resting heart rate ≥ 60 bpm. Patients had stable symptoms of heart failure and/or angina for at least 3 months and were receiving conventional cardiovascular medications at stable doses for at least 1 month. Beta-blocker therapy was not required, nor was there a protocol mandate to achieve any specific dosing targets for patients who were taking beta-blockers. Patients were randomized 1:1 to ivabradine tablets or placebo at an initial dose of 5 mg twice daily with the dose increased to 7.5 mg twice daily depending on resting heart rate and tolerability. The primary endpoint was the composite of time to first cardiovascular death, hospitalization for acute myocardial infarction, or hospitalization for new-onset or worsening heart failure. Most patients were NYHA class II (61.4%) or class III (23.2%) - none were class IV. Through a median follow-up of 19 months, ivabradine tablets did not significantly affect the primary composite endpoint (HR 1.00, 95% CI = 0.91, 1.10).

The Study Assessing the Morbi-mortality Benefits of the I<sub>f</sub> Inhibitor Ivabradine in Patients with Coronary Artery Disease Trial (SIGNIFY) was a randomized, double-blind trial administering ivabradine tablets or placebo to 19,102 adult patients with stable coronary artery disease but without clinically evident heart failure (NYHA class I). Beta-blocker therapy was not required. Ivabradine tablets was initiated at a dose of 7.5 mg twice daily and the dose could be increased to as high as 10 mg twice daily or down-titrated to 5.0 mg twice daily to achieve a target heart rate of 55 to 60 bpm. The primary endpoint was a composite of the first occurrence of either cardiovascular death or myocardial infarction. Through a median follow-up of 24.1 months, ivabradine tablets did not significantly affect the primary composite endpoint (HR 1.08, 95% CI = 0.96, 1.20).

## **16 HOW SUPPLIED/STORAGE AND HANDLING**

Ivabradine tablets 5 mg tablets are formulated as Light salmon-colored, Capsule-shaped, film-coated scored tablets debossed with "I5" divided by the score on one side and plain on the other.

They are supplied as follows:

- Bottles of 60 tablets with child-resistant closure, (NDC 50742-362-60)
- Bottles of 180 tablets with child-resistant closure, (NDC 50742-362-18)

Ivabradine tablets 7.5 mg tablets are formulated as Light salmon-colored, Round-shaped, film-coated tablets debossed with "I" on one side and "7.5" on the other side.

They are supplied as follows:

- Bottles of 60 tablets with child-resistant closure, (NDC 50742-363-60)
- Bottles of 180 tablets with child-resistant closure, (NDC 50742-363-18)

### Storage

Store ivabradine tablets at 25°C (77°F); excursions permitted to 15°C - 30°C (59°F -

86°F) [see USP Controlled Room Temperature].

## 17 PATIENT COUNSELING INFORMATION

Advise the patient to read the FDA-approved patient labeling [see *Medication Guide and Instructions for Use*].

### ● Fetal Toxicity

Advise pregnant women of the potential risks to a fetus.

Advise females of reproductive potential to use effective contraception and to notify their healthcare provider with a known or suspected pregnancy [see *Warnings and Precautions (5.1) and Use in Specific Populations (8.1), (8.3)*].

### ● Low Heart Rate

Advise patients to report significant decreases in heart rate or symptoms such as dizziness, fatigue, or hypotension [see *Warnings and Precautions (5.3)*].

### ● Atrial Fibrillation

Advise patients to report symptoms of atrial fibrillation, such as heart palpitations or racing, chest pressure, or worsened shortness of breath [see *Warnings and Precautions (5.2)*].

### ● Phosphenes

Advise patients about the possible occurrence of luminous phenomena (phosphenes). Advise patients to use caution if they are driving or using machines in situations where sudden changes in light intensity may occur, especially when driving at night. Advise patients that phosphenes may subside spontaneously during continued treatment with ivabradine tablets [see *Adverse Reactions (6.1)*].

### ● Drug Interactions

Advise patients to avoid ingestion of grapefruit juice and St. John's wort [see *Drug Interactions (7.1)*].

### ● Intake with Food

Advise patients to take ivabradine tablets twice daily with food [see *Dosage and Administration (2)*].

### **Manufactured for:**

Ingenus Pharmaceuticals, LLC

Orlando, FL 32811-7193

Made in India

Dispense with Medication Guide available at: [www.ingenus.com/medguide/ivabradine-tablets.pdf](http://www.ingenus.com/medguide/ivabradine-tablets.pdf)



## SPL MEDGUIDE

Dispense with Medication Guide available at: [www.ingenus.com/medguide/ivabradine-tablets.pdf](http://www.ingenus.com/medguide/ivabradine-tablets.pdf)

### MEDICATION GUIDE

#### Ivabradine (eye VAB ra deen) Tablets

##### **What is the most important information I should know about ivabradine tablets?**

**Ivabradine tablets may cause serious side effects in adults and children, including:**

- **Harm to an unborn baby.** Females who are able to get pregnant:
  - Must use effective birth control during treatment with ivabradine tablets.
  - Tell your doctor right away if you become pregnant during treatment with ivabradine tablets.
- **Increased risk of irregular or rapid heartbeat (atrial fibrillation or heart rhythm problems).** Tell your doctor if you feel any of the following symptoms of an irregular or rapid heartbeat:
  - heart is pounding or racing (palpitations).
  - chest pressure.
  - worsened shortness of breath.
  - near fainting or fainting.
- **Slower than normal heart rate (bradycardia).** Tell your doctor if you have:
  - a slowing of heart rate, **or**
  - symptoms of a slow heart rate such as dizziness, fatigue, lack of energy. In young children signs and symptoms of slow heart rate may include: poor feeding, difficulty breathing or turning blue

##### **What is ivabradine tablets ?**

Ivabradine tablets is a prescription medicine used:

- to treat adults who have chronic (lasting a long time) heart failure, with symptoms, to reduce their risk of hospitalization for worsening heart failure.

##### **Who should not take ivabradine tablets ?**

**Do not** take ivabradine tablets if you have:

- symptoms of heart failure that recently worsened
- very low blood pressure (hypotension)
- certain heart conditions: sick sinus syndrome, sinoatrial block, or 3<sup>rd</sup> degree atrioventricular block
- a slow resting heart rate before treatment with ivabradine tablets. Ask your doctor what a slow resting heart rate is for you.
- certain liver problems
- been prescribed any medicines that can increase the effects of ivabradine tablets.

Ask your doctor if you are not sure if you have any of the medical conditions listed above.

##### **What should I tell my doctor before taking ivabradine tablets ?**

**Before you take ivabradine tablets, tell your doctor about all of your medical conditions,**

## Including if you:

- have any other heart problems, including heart rhythm problems, a slow heart rate, or a heart conduction problem.
- are breastfeeding or planning to breastfeed. It is not known if ivabradine passes into breast milk. You and your doctor should decide if you will take ivabradine tablets or breastfeed; do not do both.
- are pregnant or planning to become pregnant. See "**What is the most important information I should know about ivabradine tablets ? - Harm to an unborn baby**" section.

Tell your doctor about all the medicines you take, including prescription and over the counter medicines, vitamins, and herbal supplements. Ivabradine tablets may affect the way other medicines work, and other medicines may affect how ivabradine tablets works. This could cause serious side effects.

## How should you take ivabradine tablets ?

- Take ivabradine tablets exactly as your doctor tells you.
- **Do not** stop taking ivabradine tablets without talking with your doctor.
- Ivabradine tablets comes as a tablet.
- Tell your doctor if you have trouble swallowing tablets.
- Your doctor may change your dose of ivabradine tablets during treatment.
- Take ivabradine tablets 2 times each day with food.
- If you miss a dose of ivabradine tablets, do not give another dose. Give the next dose at the usual time.
- If you or your child take too much ivabradine tablets, call your doctor or go to the nearest emergency room right away.

## What should you avoid while taking ivabradine tablets ?

- Avoid drinking grapefruit juice and taking St. John's wort during treatment with ivabradine tablets. These can affect the way ivabradine tablets works and may cause serious side effects.

## What are the possible side effects of ivabradine tablets ?

**Ivabradine tablets may cause serious side effects. See "What is the most important information I should know about ivabradine tablets ?"**

The most common side effects of ivabradine tablets are:

- increased blood pressure
- temporary brightness in part of your field of vision. This is usually caused by sudden changes in light (luminous phenomena). This brightness usually happens within the first 2 months of treatment with ivabradine tablets and may go away during or after treatment with ivabradine tablets. Be careful when driving or operating machinery where sudden changes in light can happen, especially when driving at night.

These are not all the side effects of ivabradine tablets. Ask your doctor or pharmacist for more information. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

## How should I store ivabradine tablets ?

- Store ivabradine tablets at room temperature between 68°F to 77°F (20°C to 25°C).

**Keep ivabradine tablets and all medicines out of the reach of children.**

## General information about the safe and effective use of ivabradine tablets .

Medicines are sometimes prescribed for purposes other than those listed in a

Medication Guide. Do not use ivabradine tablets for a condition for which it was not prescribed. Do not give ivabradine tablets to other people, even if they have the same symptoms that you have. It may harm them. You can ask your doctor or pharmacist for information about ivabradine tablets that is written for health professionals.

### What are the ingredients in ivabradine tablets ?

**Active ingredient:** ivabradine

**Inactive ingredients:**


**Tablet:** lactose monohydrate, maltodextrin, maize starch b, colloidal silicon dioxide, magnesium stearate, hypromellose, titanium dioxide, polyethylene glycol 4000, yellow iron oxide and red iron oxide.

### Manufactured for:

Ingenus Pharmaceuticals, LLC

Orlando, FL 32811-7193

Made in India



For more information contact Ingenus Pharmaceuticals, LLC at 1-877-748-1970

This Medication Guide has been approved by the U.S. Food and Drug Administration.

**Revised:** 09/2023

## PACKAGE LABEL.PRINCIPAL DISPLAY PANEL

212730 362-60 v1/09-2023 MH/DRUGS/PD/182

**Manufactured for:**  
Ingenus Pharmaceuticals, LLC  
Orlando, FL 32811-7193  
Made in India

**ingenus**  
NDC 50742-362-60  
**Ivabradine  
Tablets**  
**5 mg**

**PHARMACIST:**  
Dispense the Medication  
Guide to each patient.

**Rx Only**

60 Film-coated tablets

Each film-coated tablet contains 5 mg ivabradine equivalent to 5.39 mg ivabradine as hydrochloride.  
Oral Use.  
Store at 25°C (77°F); excursions permitted to 15°C to 30°C (between 59°F to 86°F) [see USP for controlled room temperature].  
Keep out of the sight and reach of children.  
This package is child-resistant.

**Recommended Dosage:**  
See prescribing information.  
Print Medication Guides at [www.ingenus.com/medguide/ivabradine-tablets.pdf](http://www.ingenus.com/medguide/ivabradine-tablets.pdf)

N 3 50742136260118

### Ivabradine Tablets, 5 mg - Bottle Label 60 count

212731 363-60 v1/09-2023 MH/DRUGS/PD/182

**Manufactured for:**  
Ingenus Pharmaceuticals, LLC  
Orlando, FL 32811-7193  
Made in India

**ingenus**  
NDC 50742-363-60  
**Ivabradine  
Tablets**  
**7.5 mg**

**PHARMACIST:**  
Dispense the Medication  
Guide to each patient.

**Rx Only**

60 Film-coated tablets

Each film-coated tablet contains 7.5 mg ivabradine equivalent to 8.085 mg ivabradine as hydrochloride.  
Oral Use.  
Store at 25°C (77°F); excursions permitted to 15°C to 30°C (between 59°F to 86°F) [see USP for controlled room temperature].  
Keep out of the sight and reach of children.  
This package is child-resistant.

**Recommended Dosage:**  
See prescribing information.  
Print Medication Guides at [www.ingenus.com/medguide/ivabradine-tablets.pdf](http://www.ingenus.com/medguide/ivabradine-tablets.pdf)

N 3 50742136360115

### Ivabradine Tablets, 7.5 mg - Bottle Label-60 count

# IVABRADINE

ivabradine tablet, film coated

## Product Information

<b>Product Type</b>	HUMAN PRESCRIPTION DRUG	<b>Item Code (Source)</b>	NDC:50742-362
<b>Route of Administration</b>	ORAL		

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>IVABRADINE HYDROCHLORIDE</b> (UNII: TP19837BZK) (IVABRADINE - UNII:3H48L0LPZQ)	IVABRADINE	5 mg

## Inactive Ingredients

Ingredient Name	Strength
<b>LACTOSE MONOHYDRATE</b> (UNII: EWQ57Q8I5X)	
<b>MALTODEXTRIN</b> (UNII: 7CVR7L4A2D)	
<b>STARCH, CORN</b> (UNII: O8232NY3S)	
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	
<b>HYPROMELLOSE 2910 (15 MPA.S)</b> (UNII: 36SFW2JZ0W)	
<b>POLYETHYLENE GLYCOL 4000</b> (UNII: 4R4HFI6D95)	
<b>FERRIC OXIDE YELLOW</b> (UNII: EX438O2MRT)	
<b>FERRIC OXIDE RED</b> (UNII: 1K09F3G675)	

## Product Characteristics

<b>Color</b>	PINK (Light salmon-colored)	<b>Score</b>	2 pieces
<b>Shape</b>	CAPSULE	<b>Size</b>	8mm
<b>Flavor</b>		<b>Imprint Code</b>	I;5
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:50742-362-60	60 in 1 BOTTLE; Type 0: Not a Combination Product	07/15/2024	

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA214051	07/15/2024	

# IVABRADINE

ivabradine tablet, film coated

## Product Information

<b>Product Type</b>	HUMAN PRESCRIPTION DRUG	<b>Item Code (Source)</b>	NDC:50742-363
<b>Route of Administration</b>	ORAL		

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>IVABRADINE HYDROCHLORIDE</b> (UNII: TP19837BZK) (IVABRADINE - UNII:3H48L0LPZQ)	IVABRADINE	7.5 mg

## Inactive Ingredients

Ingredient Name	Strength
<b>LACTOSE MONOHYDRATE</b> (UNII: EWQ57Q8I5X)	
<b>MALTODEXTRIN</b> (UNII: 7CVR7L4A2D)	
<b>STARCH, CORN</b> (UNII: O8232NY3SJ)	
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	
<b>HYPROMELLOSE 2910 (15 MPA.S)</b> (UNII: 36SFW2JZ0W)	
<b>POLYETHYLENE GLYCOL 4000</b> (UNII: 4R4HFI6D95)	
<b>FERRIC OXIDE YELLOW</b> (UNII: EX438O2MRT)	
<b>FERRIC OXIDE RED</b> (UNII: 1K09F3G675)	

## Product Characteristics

<b>Color</b>	PINK (Light salmon-colored)	<b>Score</b>	no score
<b>Shape</b>	ROUND	<b>Size</b>	7mm
<b>Flavor</b>		<b>Imprint Code</b>	l;7;5
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:50742-363-60	60 in 1 BOTTLE; Type 0: Not a Combination Product	07/15/2024	

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA214051	07/15/2024	

**Labeler** - Ingenus Pharmaceuticals, LLC (833250017)

**Registrant** - Ingenus Pharmaceuticals, LLC (833250017)

Revised: 2/2025

Ingenus Pharmaceuticals, LLC