

**EXTRA STRENGTH TYLENOL SEVERE COUGH PLUS SORE THROAT DAY-
acetaminophen and dextromethorphan hydrobromide liquid
Kenvue Brands LLC**

Extra Strength TYLENOL® Severe Cough + Sore Throat Day

Drug Facts

Active ingredients(in each 30 mL)

Acetaminophen 1,000 mg

Dextromethorphan HBr 30 mg

Purpose

Pain reliever/fever reducer

Cough suppressant

Uses

- temporarily relieves these common cold/flu symptoms:
- minor aches and pains
- headache
- sore throat
- cough
- temporarily reduces fever

Warnings

Liver Warning

This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert

acetaminophen may cause severe skin reactions.

Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning

If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains

acetaminophen, ask a doctor or pharmacist.

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- liver disease
- persistent or chronic cough such as occurs with smoking, asthma or emphysema
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are

taking the blood thinning drug warfarin

When using this product

do not exceed recommended dose

Stop use and ask a doctor if

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts

These could be signs of a serious condition.

If pregnant or breast-feeding,

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not take more than directed(see overdose warning)**
- **shake well before using**
- mL = milliliter
- use only enclosed dosing cup designed for use with this product. Do not use any other dosing device.

adults and children 12 years and over	<ul style="list-style-type: none">• take 30 mL in the dosing cup provided every 6 hours while symptoms last• do not take more than 90 mL in 24 hours, unless directed by a doctor• do not use for more than 10 days unless directed by a doctor
children under 12 years	ask a doctor

Other information

- each 30 mL contains: **sodium 19 mg**
- store between **20-25°C (68-77°F)**
- **do not use if neck band imprinted with “INSTANT COOLING SENSATION” or foil inner seal imprinted with**

“TYLENOL” is broken or missing

Inactive ingredients

anhydrous citric acid, carboxymethylcellulose sodium, FD&C red no 40, flavors, glycerin,microcrystalline cellulose and carboxymethylcellulose sodium,poloxamer, polyethylene glycol,propylene glycol, purified water,sodium benzoate, sorbitol, sucralose

Questions or comments?

call **1-877-895-3665** (toll-free) or
215-273-8755 (collect)

PRINICIPAL DISPLAY PANEL

NEW

NDC 50580-140-08

EXTRA STRENGHT TYLENOL ®

FOR ADULTS

SEVERE COUGH + SORE THROAT

Acetaminophen,

Dextromethorphan HBr, Pain Reliever-Fever Reducer, Cough Suppressant

- **HEAD+BODY ACHES**
- **FEVER+ SORE THROAT**
- **COUGH**

DAY

COOLING CHERRY

8 fl oz (240 mL)



30057937

Do not use if neck band imprinted with
"INSTANT COOLING SENSATION" or
foil inner seal imprinted with
"TYLENOL" is broken or missing

**DO NOT USE WITH OTHER MEDICINES
CONTAINING ACETAMINOPHEN**



3 00450 56508 2

Drug Facts

Active ingredients Purpose (in each 30 mL)

Acetaminophen 1,000 mg Pain reliever/
fever reducer

Dextromethorphan HBr 30 mg Cough
suppressant

Uses

- temporarily relieves these common cold/flu symptoms:
 - minor aches and pains
 - headache
 - sore throat
 - cough
- temporarily reduces fever

Drug Facts (continued)

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: acetaminophen may cause severe skin reactions.

Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Drug Facts (continued)

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for

Drug Facts (continued)

2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- liver disease
- persistent or chronic cough such as occurs with smoking, asthma or emphysema
- cough that occurs with too much phlegm (mucus)

Drug Facts (continued)

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

When using this product do not exceed recommended dose

Stop use and ask a doctor if

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts

These could be signs of a serious condition.

Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than directed (see overdose warning)
- shake well before using
- mL = milliliter

Drug Facts (continued)

- use only enclosed dosing cup designed for use with this product. Do not use any other dosing device.

adults
and
children
12 years
and over

- take 30 mL in the dosing cup provided every 6 hours while symptoms last
- do not take more than 90 mL in 24 hours, unless directed by a doctor
- do not use for more than 10 days unless directed by a doctor

children
under
12 years

ask a doctor

Drug Facts (continued)

Other information

- each 30 mL contains: **sodium 19 mg**
- store between 20-25°C (68-77°F)
- do not use if neck band imprinted with "INSTANT COOLING SENSATION" or foil inner seal imprinted with "TYLENOL" is broken or missing

Inactive ingredients

anhydrous citric acid, carboxymethylcellulose sodium, FD&C red no 40, flavors, glycerin, microcrystalline cellulose and carboxymethylcellulose sodium, poloxamer, polyethylene glycol, propylene glycol, purified water, sodium benzoate, sorbitol, sucralose

Drug Facts (continued)

Questions or comments?

call 1-877-895-3665 (toll-free) or
215-273-8755 (collect)

PARENTS:

Learn about teen medicine abuse

www.StopMedicineAbuse.org

Extra Strength
TYLENOL
FOR ADULTS
SEVERE COUGH

+ SORE THROAT
Day

How can we help?
1-877-895-3665

Acetaminophen
(1,000 mg per 30 mL)

Dextromethorphan HBr
(30 mg per 30 mL)

8 fl oz (240 mL)

Distributed by:
Kenvue Brands LLC
Summit, NJ 07901

© Kenvue Brands LLC 2025

30057937

EXTRA STRENGTH TYLENOL SEVERE COUGH PLUS SORE THROAT DAY				
acetaminophen and dextromethorphan hydrobromide liquid				
Product Information				
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:50580-140	
Route of Administration	ORAL			
Active Ingredient/Active Moiety				
Ingredient Name		Basis of Strength	Strength	
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)		ACETAMINOPHEN	1000 mg in 30 mL	
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)		DEXTROMETHORPHAN HYDROBROMIDE	30 mg in 30 mL	
Inactive Ingredients				
Ingredient Name			Strength	
WATER (UNII: 059QF0KO0R)				
CARBOXYMETHYLCELLULOSE SODIUM, UNSPECIFIED (UNII: K679OBS311)				
FD&C RED NO. 40 (UNII: WZB9127XOA)				
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)				
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)				
SODIUM BENZOATE (UNII: OJ245FE5EU)				
SUCRALOSE (UNII: 96K6UQ3ZD4)				
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)				
GLYCERIN (UNII: PDC6A3C0OX)				
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)				
SORBITOL (UNII: 506T60A25R)				
Product Characteristics				
Color	red	Score		
Shape		Size		
Flavor	BERRY	Imprint Code		
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:50580-140-08	240 mL in 1 BOTTLE; Type 0: Not a Combination Product	06/06/2025	
Marketing Information				
Marketing		Application Number or Monograph	Marketing Start	Marketing End

Category	Citation	Date	Date
OTC Monograph Drug	M012	06/06/2025	

Labeler - Kenvue Brands LLC (118772437)

Revised: 3/2025 Kenvue Brands LLC