

RITE AID ACETAMINOPHEN- acetaminophen tablet
Praxis, LLC

Rite Aid Corporation Acetaminophen Drug Facts

Active ingredient (in each caplet)

Acetaminophen 500 mg

Purpose

Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
- the common cold
- headache
- backache
- minor pain of arthritis
- toothache
- muscular aches
- premenstrual and menstrual cramps
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

liver disease

Ask a doctor or pharmacist before use if you are

taking the blood thinning drug warfarin

Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

Overdose warning:In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not take more than directed (see overdose warning)**

| | |
|---------------------------------------|--|
| adults and children 12 years and over | <ul style="list-style-type: none">• take 2 caplets every 6 hours while symptoms last• do not take more than 6 caplets in 24 hours, unless directed by a doctor• do not use for more than 10 days unless directed by a doctor |
| children under 12 years | ask a doctor |

Other information

- store at 20-25°C (68-77°F)

Inactive ingredients

carnauba wax, corn starch*, croscarmellose sodium*, hypromellose, polyethylene glycol, povidone, pregelatinized starch, sodium starch glycolate*, stearic acid

*may contain one or more of these ingredients

Questions or comments?

1-800-719-9260

Principal Display Panel

Compare to the active ingredient in Extra Strength Tylenol® Caplets

FREE FROM

GLUTEN FREE

CAFFEINE FREE

EXTRA STRENGTH

ACETAMINOPHEN

ACETAMINOPHEN CAPLETS, 500 mg



ACTUAL SIZE

PAIN RELIEVER/FEVER REDUCER

For adults

100 CAPLETS





Drug Facts

Active ingredient (in each caplet) **Purpose**

Acetaminophen 500 mg.....Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
 - the common cold ■ headache
 - minor pain of arthritis ■ backache
 - muscular aches ■ toothache
 - premenstrual and menstrual cramps
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening ■ blisters ■ rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have liver disease

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days
 - fever gets worse or lasts more than 3 days
 - new symptoms occur ■ redness or swelling is present
- These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Drug Facts (continued)

Keep out of reach of children. Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

■ do not take more than directed (see overdose warning)

adults and children 12 years and over

- take 2 caplets every 6 hours while symptoms last
- do not take more than 6 caplets in 24 hours, unless directed by a doctor
- do not use for more than 10 days unless directed by a doctor

children under 12 years

ask a doctor


Other information ■ store at 20-25°C (68-77°F)

Inactive ingredients carnauba wax, corn starch*, croscarmellose sodium*, hypromellose, polyethylene glycol, povidone, pregelatinized starch, sodium starch glycolate*, stearic acid *may contain one or more of these ingredients

Questions or comments? 1-800-719-9260

****This product is not manufactured or distributed by Johnson & Johnson Consumer Inc., distributor of Extra Strength Tylenol® Caplets.**





RITE AID ACETAMINOPHEN

acetaminophen tablet

Product Information

| | | | |
|--------------------------------|----------------|---------------------------|---------------|
| Product Type | HUMAN OTC DRUG | Item Code (Source) | NDC:59368-227 |
| Route of Administration | ORAL | | |

Active Ingredient/Active Moiety

| Ingredient Name | Basis of Strength | Strength |
|---|-------------------|----------|
| ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D) | ACETAMINOPHEN | 500 mg |

Inactive Ingredients

| Ingredient Name | Strength |
|--|----------|
| CARNAUBA WAX (UNII: R12CBM0EIZ) | |
| STARCH, CORN (UNII: O8232NY3SJ) | |
| CROSCARMELLOSE SODIUM (UNII: M28OL1HH48) | |
| HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO) | |
| POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A) | |
| POVIDONE, UNSPECIFIED (UNII: FZ989GH94E) | |
| STEARIC ACID (UNII: 4ELV7Z65AP) | |

Product Characteristics

| | | | |
|-----------------|-------|---------------------|----------|
| Color | white | Score | no score |
| Shape | OVAL | Size | 16mm |
| Flavor | | Imprint Code | L484 |
| Contains | | | |

Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|---|------------------|---------------------|----------------------|--------------------|
| 1 | NDC:59368-227-02 | 1 in 1 CARTON | 09/08/2020 | |

| | | | | |
|---|------------------|--|------------|--|
| 1 | | 225 in 1 BOTTLE; Type 0: Not a Combination Product | | |
| 2 | NDC:59368-227-04 | 1 in 1 CARTON | 07/11/2022 | |
| 2 | | 50 in 1 BOTTLE; Type 0: Not a Combination Product | | |
| 3 | NDC:59368-227-01 | 1 in 1 CARTON | 07/11/2022 | |
| 3 | | 100 in 1 BOTTLE; Type 0: Not a Combination Product | | |
| 4 | NDC:59368-227-03 | 500 in 1 BOTTLE; Type 0: Not a Combination Product | 07/11/2022 | |

Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|--------------------|--|----------------------|--------------------|
| OTC Monograph Drug | M013 | 09/04/2020 | |

Labeler - Praxis, LLC (016329513)

Establishment

| Name | Address | ID/FEI | Business Operations |
|-------------|---------|-----------|---|
| Praxis, LLC | | 016329513 | manufacture(59368-227) , pack(59368-227) , label(59368-227) |

Revised: 1/2023

Praxis, LLC