SYLVIRENE LIQUID WART CORN CALLUS REMOVER- salicylic acid 17% liquid wart corn callus remover liquid Jiangxi Hemei Pharmaceutical Co., Ltd

84010-079

Active Ingredient

Salicylic acid 17%

Purpose

Wart remover

Use

For the removal of common warts and plantar warts. Common warts have a rough "cauliflower-like" surface and are easily recognizable. Plantar warts are located only on the soles of the feet, tender to touch, with an interrupted footprint pattern.

Warnings

For external use only. Flammable.

Keep away from fire and flame.

Do not use

on irritated skin or on any area that is infected or reddened.On moles, birthmarks, warts with hair growing from them, genital warts, or warts on the face or mucous membranes. If you have diabetes or poor blood circulation.

When Using

Avoid contact with eyes. If product gets into the eye, flush with Water for 15 minutes. Non-edible. Cap tightly and store at room temperature, away from heat.

Stop Use

Stop use and ask a doctor if discomfort persists.

Ask Doctor

Stop use and ask a doctor if discomfort persists.

Keep Out Of Reach Of Children

If swallowed, get medical help or contact the Poison Control Center(1-800-222-1222) right away.

Directions

Wash the affected area.

May soak wart in warm water for 5 minutes. Dry area thoroughly.

Use a cotton swab to apply a sufficient amount to cover each wart. Let it dry.

Repeat this procedure once or twice daily as needed (until wart is removed) for upto 12 weeks.

Other information

Store at room temperature. Avoid excessive heat $(37^{\circ}\text{C}/99^{\circ}\text{F})$. Skin discoloration may occur during or after use.

Inactive ingredients

Water, Polyethylene Glycol, Glycerin, Borneol, Podofilox

PRINCIPAL DISPLAY PANEL



Stop use and ask a doctor if discomfort persists.

Keep out of reach of children.If swallowed, get medical help or contact the Poison Control Center (1-800-222-1222) right away. **Directions**

Wash the affected area.

May soak wart in warm water for 5 minutes.
Dry area thoroughly.

Use a cotton swab to apply a sufficient amount to cover each wart. Let it dry.

Repeat this procedure once or twice daily as needed (until wart is removed) for up to 12 weeks.

Other information

Store at room temperature.

Avoid excessive heat (37°C / 99°F). Skin discoloration may occur during or after use.

Inactive ingredients:

Water
Polyethylene Glycol·····16%
Glycerin · · · · · · · · · · · · 8%
Borneol·····6%
Podofilox······0.5%
*



SYLVIRENE TM

LIQUID WART CORN&CALLUS REMOVER



- ☑ Prevents Spread
- ☑ Targeting Removal
- ☑ Safe For Adults

MEDICAL STRENGTH NET WT: 30ml / 1fl oz

Drug Facts Active ingredient:

Purpose

Salicylic acid 17%····· Wart remover

UsesFor the removal of common warts and plantar warts.

Common warts have a rough "cauliflower -like" surface and are easily recognizable. Plantar warts are located only on the soles of the feet, tender to touch, with an interrupted footprint pattern.

Warnings

For external use only. Flammable. Keep away from fire and flame.

Do not use

on irritated skin or on any area that is infected or reddened.On moles, birthmarks, warts with hair growing from them, genital warts, or warts on the face or mucous membranes.If you have diabetes or poor blood circulation.

When using this product

Avoid contact with eyes. If product gets into the eye, flush with Water for 15 minutes.Non-edible.
Cap tightly and store at room temperature,

SYLVIRENE TM

LIQUID WART CORN&CALLUS REMOVER



- ☑ Prevents Spread
- ☑ Targeting Removal
- ☑ Safe For Chinldren & Adults

MEDICAL STRENGTH NET WT: 30ml / 1fl oz

SYLVIRENE LIQUID WART CORN CALLUS REMOVER

salicylic acid 17% liquid wart corn callus remover liquid

Product Information

Product Type HUMAN OTC DRUG Item Code (Source) NDC:84010-079

Route of Administration TOPICAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
CALLOWING ACID (UNIII) O 41 4 D.Z. 41 D.Z.) (CALLOVING ACID. LINIII) O 41 4 D.Z. 41 D.Z.)	CALICYLIC ACID	17 - 1- 100

SALICYLIC ACID (UNII: O414PZ4LPZ) (SALICYLIC ACID - UNII:O414PZ4LPZ) SALICYLIC ACID 17 g in 100 mL

Inactive Ingredients			
Ingredient Name	Strength		
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)			
WATER (UNII: 059QF0KO0R)			
GLYCERIN (UNII: PDC6A3C0OX)			
BORNEOL (UNII: M89NIB437X)			
PODOFILOX (UNII: L36H50F353)			

l	Packaging				
	#	Item Code	Package Description	Marketing Start Date	Marketing End Date
		NDC:84010-079- 01	30 mL in 1 BOTTLE; Type 0: Not a Combination Product	02/21/2025	

Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
OTC Monograph Drug	M028	02/21/2025		

Labeler - Jiangxi Hemei Pharmaceutical Co., Ltd (724892056)

Establishment					
Name	Address	ID/FEI	Business Operations		
Jiangxi Hemei Pharmaceutical Co., Ltd		724892056	manufacture(84010-079)		

Revised: 2/2025 Jiangxi Hemei Pharmaceutical Co., Ltd