

SYLVIRENE TATTOO NUMBING- lidocaine4% tattoo numbing cream
Jiangxi Hemei Pharmaceutical Co., Ltd

84010-080 Update

Active Ingredient

Lidocaine 4%

Purpose

Local Anesthetic

Use

For temporary relief of local discomfort, itching, pain, soreness, or burning related to the tattoo process.

Warnings

For external use only.

Do not use

In children less than 2 months of age on open skin wounds. Do not exceed the recommended daily dosage unless directed by a doctor.

When Using

Avoid contact with eyes. If this happens, rinse thoroughly with water.
If symptoms persist for more than 7 days or subside but occur again within 3 days, stop use and consult a physician.

Stop Use

Stop use and seek medical attention if signs of an allergy occur, such as redness, irritation, swelling, pain, or other severe symptoms. If symptoms do not subside completely within 7 days, seek medical help.

Ask Doctor

Stop use and seek medical attention if signs of an allergy occur, such as redness, irritation, swelling, pain, or other severe symptoms. If symptoms do not subside completely within 7 days, seek medical help.

Keep Out Of Reach Of Children

Keep out of reach of children.

Directions

Before applying the numbing cream, thoroughly wash your hands and wear disposable tattoo gloves.

Clean the area of the body receiving the tattoo. Dry completely.

Apply a generous amount of numbing cream to the cleaned area and rub it in thoroughly in a circular motion for 30 to 60 seconds. Then, cover with plastic wrap.

Leave the cream and wrap in place for 30 to 45 minutes. Wipe away the cream and thoroughly clean the area to prepare for a pain-free tattoo.

Other information

Store between 15-30°C(59-86 F). Avoid freezing and excessive heat above 40 °C(104°F).

Inactive ingredients

Water, Carbomer 934, Propylene Glycol, Soybean Lecithin, Alpha-Tocopherol Acetate, Benzyl Alcohol.

PRINCIPAL DISPLAY PANEL

Inactive Ingredients				
Ingredient Name			Strength	
WATER (UNII: 059QF0KO0R)				
.ALPHA.-TOCOPHEROL ACETATE (UNII: 9E8X80D2L0)				
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)				
CARBOMER 934 (UNII: Z135WT9208)				
SOYBEAN LECITHIN (UNII: 1DI56QDM62)				
BENZYL ALCOHOL (UNII: LKG8494WBH)				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:84010-080-01	20 g in 1 TUBE; Type 0: Not a Combination Product	02/21/2025	
Marketing Information				
Marketing Category		Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug		M017	02/21/2025	

Labeler - Jiangxi Hemei Pharmaceutical Co., Ltd (724892056)

Establishment			
Name	Address	ID/FEI	Business Operations
Jiangxi Hemei Pharmaceutical Co., Ltd		724892056	manufacture(84010-080)

Revised: 3/2025

Jiangxi Hemei Pharmaceutical Co., Ltd