

ACETAMINOPHEN GELCAPS- acetaminophen tablet
TARGET CORPORATION

673T Target Acetaminophen Gelcaps

Drug Facts

Active ingredient (in each gelcap)

Acetaminophen USP 500 mg

Purpose

Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
 - headache
 - muscular aches
 - backache
 - minor pain of arthritis
 - the common cold
 - toothache
 - premenstrual and menstrual cramps
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if you have liver disease

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not take more than directed (see overdose warning)**

adults and children 12 years and over	<ul style="list-style-type: none">• take 2 gelcaps every 6 hours while symptoms last• do not take more than 6 gelcaps in 24 hours, unless directed by a doctor• do not use for more than 10 days unless directed by a doctor
children under 12 years	ask a doctor

Other information

- each gelcap contains: **sodium 1 mg**
- store between 20-25°C (68-77°F). Avoid excessive heat above 40°C (104°F)

Inactive ingredients croscarmellose sodium, colloidal anhydrous silica, D&C red #33, FD&C blue #1, FD&C red #40, FD&C yellow #6, ferric oxide red, ferric oxide yellow, ferrosoferric oxide, gelatin, hypromellose, polyethylene glycol, povidone, pregelatinized starch, stearic acid, talc, titanium dioxide

Questions or comments? Call **1-800-910-6874**

Compare to active ingredient in
Extra Strength Tylenol®
Rapid Release Gels*

Acetaminophen

500 mg Gelcaps,
Pain Reliever / Fever Reducer
For Adults

DO NOT USE WITH OTHER MEDICINES
CONTAINING ACETAMINOPHEN



225 GELCAPS



Actual Size
225 Gelcaps

TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING

Drug Facts

Active ingredient (in each gelcap) **Purpose**
Acetaminophen USP 500 mg.....Pain reliever/fever reducer

Uses

temporarily relieves minor aches and pains due to:
■ headache ■ muscular aches ■ backache
■ minor pain of arthritis ■ the common cold
■ toothache ■ premenstrual and menstrual cramps
temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take
■ more than 4,000 mg of acetaminophen in 24 hours
■ with other drugs containing acetaminophen
■ 3 or more alcoholic drinks every day while using this product
Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include:
■ skin reddening ■ blisters ■ rash
If a skin reaction occurs, stop use and seek medical help right away.

Do not use
■ with any other drug containing acetaminophen

Drug Facts (continued)

(prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
■ if you are allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if you have liver disease

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

Stop use and ask a doctor if

■ pain gets worse or lasts more than 10 days
■ fever gets worse or lasts more than 3 days
■ new symptoms occur
■ redness or swelling is present
These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. **Keep out of reach of children.**

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

Do not take more than directed (see overdose warning)

Drug Facts (continued)

adults and children 12 years and over
■ take 2 gelcaps every 6 hours while symptoms last
■ do not take more than 6 gelcaps in 24 hours, unless directed by a doctor
■ do not use for more than 10 days unless directed by a doctor

children under 12 years ask a doctor

Other information ■ each gelcap contains: sodium 1 mg ■ store at 20-25°C (68-77°F). Avoid excessive heat above 40°C (104°F) ■ avoid high humidity

Inactive ingredients

colloidal anhydrous silica, croscarmellose sodium, D&C red #33, FD&C blue #1, FD&C red #40, FD&C yellow #6, ferric oxide red, ferric oxide yellow, ferrous ferric oxide, gelatin, hypromellose, polyethylene glycol, povidone, pregelatinized starch, stearic acid, talc, titanium dioxide

Questions or comments? Call 1-800-910-8874

*This product is not manufactured or distributed by Kenvue Inc., owner of the registered trademark Extra Strength Tylenol® Rapid Release Gels.

Contains No Aspirin
Distributed by Target Corporation
Minneapolis, MN 55403
NDC 11673-674-34
094 01 0048 R00 C-002262-01-034-0000



Made in India
TM & ©2025 Target Brands, Inc.
673T 0425

Varnish Omit

3

11673-674-34 5

LOT:
EXP.:

Contains No Aspirin

Adhesive Area

Drug Facts (continued)

Do not use

■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
■ if you are allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if you have liver disease

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

Stop use and ask a doctor if

■ pain gets worse or lasts more than 10 days
■ fever gets worse or lasts more than 3 days
■ new symptoms occur
■ redness or swelling is present
These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. **Keep out of reach of children.**

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

Do not take more than directed (see overdose warning)

adults and children 12 years and over
■ take 2 gelcaps every 6 hours while symptoms last
■ do not take more than 6 gelcaps in 24 hours, unless directed by a doctor
■ do not use for more than 10 days unless directed by a doctor

children under 12 years ask a doctor

Other information

■ each gelcap contains: sodium 1 mg
■ store at 20-25°C (68-77°F). Avoid excessive heat above 40°C (104°F)
■ avoid high humidity

Inactive ingredients

colloidal anhydrous silica, croscarmellose sodium, D&C red #33, FD&C blue #1, FD&C red #40, FD&C yellow #6, ferric oxide red, ferric oxide yellow, ferrous ferric oxide, gelatin, hypromellose, polyethylene glycol, povidone, pregelatinized starch, stearic acid, talc, titanium dioxide

Questions or comments? Call 1-800-910-8874

1/16" Margin

Compare to active ingredient in Extra
Strength Tylenol® Rapid Release Gels*

Acetaminophen

500 mg Gelcaps,
Pain Reliever / Fever Reducer

For Adults



100 GELCAPS

DO NOT USE WITH OTHER MEDICINES
CONTAINING ACETAMINOPHEN



Actual Size
100 Gelcaps

TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING

Drug Facts

Active ingredient (in each gelcap) **Purpose**
Acetaminophen USP 500 mg.....Pain reliever/fever reducer

Uses

temporarily relieves minor aches and pains due to:
■ headache ■ muscular aches ■ backache
■ minor pain of arthritis ■ the common cold
■ toothache ■ premenstrual and menstrual cramps
temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take
■ more than 4,000 mg of acetaminophen in 24 hours
■ with other drugs containing acetaminophen
■ 3 or more alcoholic drinks every day while using this product

Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include:
■ skin reddening ■ blisters ■ rash
If a skin reaction occurs, stop use and seek medical help right away.

Drug Facts (continued under label)

NDC 11673-674-73

094 01 0547 R00 C-002262-01-034-0000



This product is not manufactured or distributed by Kenvue Inc., owner of the registered trademark Extra Strength Tylenol® Rapid Release Gels.

Dist. by Target Corp.

Mpls, MN 55403

Made in India

TM & ©2025

Target Brands, Inc.

673T 0425

LOT:

EXP.:

Varnish Omit Area

Acetaminophen

500 mg Gelcaps,
Pain Reliever /
Fever Reducer
For Adults

DO NOT USE WITH OTHER MEDICINES
CONTAINING ACETAMINOPHEN



50 GELCAPS

RETAIN CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION

TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL
SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING

Drug Facts

Active ingredient (in each gelcap)

Acetaminophen USP 500 mg Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
 - headache ■ muscular aches ■ backache
 - minor pain of arthritis ■ the common cold ■ toothache
 - premenstrual and menstrual cramps
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take ■ more than 4,000 mg of acetaminophen in 24 hours ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product

Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash. If a skin reaction occurs, stop use and seek medical help right away. If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. **Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

■ do not take more than directed (see overdose warning) adults and children 12 years and over ■ take 2 gelcaps every 6 hours while symptoms last ■ do not take more than 6 gelcaps in 24 hours, unless directed by a doctor ■ do not use for more than 10 days unless directed by a doctor children under 12 years: ask a doctor

Other information ■ each gelcap contains: sodium 1 mg ■ store at 20-25°C (68-77°F). Avoid excessive heat above 40°C (104°F) ■ avoid high humidity

Contains No Aspirin

Questions or comments? Call 1-800-910-6874 6/3T 0425

Dist. by Target Corp., Mpls., MN 55403 NDC167674/08

Made in India 094 01 0548 R0Q

TM & ©2025 Target Brands, Inc. C-002262-01-034-0000

LOT:

EXP.:

Varnish Omit Area

Active Ingredient/Active Moiety			
Ingredient Name		Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)		ACETAMINOPHEN	500 mg

Inactive Ingredients	
Ingredient Name	Strength
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
POVIDONE (UNII: FZ989GH94E)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TALC (UNII: 7SEV7J4R1U)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
D&C RED NO. 33 (UNII: 9DBA0SBB0L)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
FERROSOFERRIC OXIDE (UNII: XM0M87F357)	
GELATIN (UNII: 2G86QN327L)	
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
STARCH, CORN (UNII: O8232NY3SJ)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	

Product Characteristics			
Color	red (Red opaque and Blue gray opaque hard gelatin shells)	Score	2 pieces
Shape	OVAL	Size	19mm
Flavor		Imprint Code	
Contains			

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:11673-674-05	1 in 1 CARTON	07/10/2025	
1		50 in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:11673-674-34	225 in 1 BOTTLE; Type 0: Not a Combination Product	07/10/2025	
3	NDC:11673-674-73	100 in 1 BOTTLE; Type 0: Not a Combination Product	07/10/2025	

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date

OTC Monograph Drug	M013	07/10/2025	
--------------------	------	------------	--

Labeler - TARGET CORPORATION (006961700)

Registrant - TIME CAP LABORATORIES, INC. (037052099)

Establishment

Name	Address	ID/FEI	Business Operations
MARKSANS PHARMA LIMITED		677604129	manufacture(11673-674)

Revised: 7/2025

TARGET CORPORATION