

HYDROCODONE BITARTRATE AND ACETAMINOPHEN- hydrocodone bitartrate and acetaminophen tablet

Apotheca Inc.

Hydrocodone Bitartrate and Acetaminophen

Tablets USP 5 mg/325 mg

C-II

Rx only

BOXED WARNING


Hepatotoxicity

Acetaminophen has been associated with cases of acute liver failure, at times resulting in liver transplant and death. Most of the cases of liver injury are associated with the use of acetaminophen at doses that exceed 4000 milligrams per day, and often involve more than one acetaminophen-containing product.


DESCRIPTION

Hydrocodone bitartrate and acetaminophen is supplied in tablet form for oral administration.

Hydrocodone bitartrate is an opioid analgesic and antitussive and occurs as fine, white crystals or as a crystalline powder. It is affected by light. The chemical name is 4,5 α -Epoxy-3-methoxy-17-methylmorphinan-6-one tartrate (1:1) hydrate (2:5). It has the following structural formula:



Acetaminophen, 4'-Hydroxyacetanilide, a slightly bitter, white, odorless, crystalline powder, is a non-opiate, non-salicylate analgesic and antipyretic. It has the following structural formula:



Each tablet contains:

Hydrocodone Bitartrate..... 5 mg

Acetaminophen..... 325 mg

In addition, each tablet contains the following inactive ingredients: colloidal silicon dioxide, croscarmellose sodium, crospovidone, microcrystalline cellulose, povidone, pregelatinized starch, stearic acid and sugar spheres which are composed of starch derived from corn, sucrose, and FD&C Yellow #6. Meets USP Dissolution Test 1.

CLINICAL PHARMACOLOGY

Hydrocodone is a semisynthetic narcotic analgesic and antitussive with multiple actions qualitatively similar to those of codeine. Most of these involve the central nervous system and smooth muscle. The precise mechanism of action of hydrocodone and other opiates is not known, although it is believed to relate to the existence of opiate receptors in the central nervous system. In addition to analgesia, narcotics may produce drowsiness, changes in mood and mental clouding.

The analgesic action of acetaminophen involves peripheral influences, but the specific mechanism is as yet undetermined. Antipyretic activity is mediated through hypothalamic heat regulating centers. Acetaminophen inhibits prostaglandin synthetase. Therapeutic doses of acetaminophen have negligible effects on the cardiovascular or respiratory systems; however, toxic doses may cause circulatory failure and rapid, shallow breathing.

Pharmacokinetics: The behavior of the individual components is described below.

Hydrocodone: Following a 10 mg oral dose of hydrocodone administered to five adult male subjects, the mean peak concentration was 23.6 ± 5.2 ng/mL. Maximum serum levels were achieved at 1.3 ± 0.3 hours and the half-life was determined to be 3.8 ± 0.3 hours. Hydrocodone exhibits a complex pattern of metabolism including O-demethylation, N-demethylation and 6-keto reduction to the corresponding 6- α - and 6- β - hydroxymetabolites. See **OVERDOSAGE** for toxicity information.

Acetaminophen: Acetaminophen is rapidly absorbed from the gastrointestinal tract and is distributed throughout most body tissues. The plasma half-life is 1.25 to 3 hours, but may be increased by liver damage and following overdosage. Elimination of acetaminophen is principally by liver metabolism (conjugation) and subsequent renal excretion of metabolites. Approximately 85% of an oral dose appears in the urine within 24 hours of administration, most as the glucuronide conjugate, with small amounts of other conjugates and unchanged drug. See **OVERDOSAGE** for toxicity information.

INDICATIONS AND USAGE

Hydrocodone bitartrate and acetaminophen tablets are indicated for the relief of moderate to moderately severe pain.

CONTRAINDICATIONS

This product should not be administered to patients who have previously exhibited hypersensitivity to hydrocodone or acetaminophen.

Patients known to be hypersensitive to other opioids may exhibit cross-sensitivity to hydrocodone.

WARNINGS

Hepatotoxicity: Acetaminophen has been associated with cases of acute liver failure, at times resulting in liver transplant and death. Most of the cases of liver injury are associated with the use of acetaminophen at doses that exceed 4000 milligrams per day, and often involve more than one acetaminophen-containing product. The excessive intake of acetaminophen may be intentional to cause

self-harm or unintentional as patients attempt to obtain more pain relief or unknowingly take other acetaminophen-containing products.

The risk of acute liver failure is higher in individuals with underlying liver disease and in individuals who ingest alcohol while taking acetaminophen.

Instruct patients to look for acetaminophen or APAP on package labels and not to use more than one product that contains acetaminophen. Instruct patients to seek medical attention immediately upon ingestion of more than 4000 milligrams of acetaminophen per day, even if they feel well.

Serious skin reactions: Rarely, acetaminophen may cause serious skin reactions such as acute generalized exanthematous pustulosis (AGEP), Stevens-Johnson Syndrome (SJS), and toxic epidermal necrolysis (TEN), which can be fatal. Patients should be informed about the signs of serious skin reactions, and use of the drug should be discontinued at the first appearance of skin rash or any other sign of hypersensitivity.

Hypersensitivity/anaphylaxis: There have been post-marketing reports of hypersensitivity and anaphylaxis associated with use of acetaminophen. Clinical signs included swelling of the face, mouth, and throat, respiratory distress, urticaria, rash, pruritus, and vomiting. There were infrequent reports of life-threatening anaphylaxis requiring emergency medical attention. Instruct patients to discontinue Hydrocodone Bitartrate and Acetaminophen Tablets, USP immediately and seek medical care if they experience these symptoms. Do not prescribe Hydrocodone Bitartrate and Acetaminophen Tablets, USP for patients with acetaminophen allergy.

Respiratory Depression: At high doses or in sensitive patients, hydrocodone may produce dose-related respiratory depression by acting directly on the brain stem respiratory center. Hydrocodone also affects the center that controls respiratory rhythm, and may produce irregular and periodic breathing.

Head Injury and Increased Intracranial Pressure: The respiratory depressant effects of narcotics and their capacity to elevate cerebrospinal fluid pressure may be markedly exaggerated in the presence of head injury, other intracranial lesions or a pre-existing increase in intracranial pressure. Furthermore, narcotics produce adverse reactions which may obscure the clinical course of patients with head injuries.

Acute Abdominal Conditions: The administration of narcotics may obscure the diagnosis or clinical course of patients with acute abdominal conditions.

PRECAUTIONS

General:

Special Risk Patients: As with any narcotic analgesic agent, hydrocodone bitartrate and acetaminophen tablets should be used with caution in elderly or debilitated patients and those with severe impairment of hepatic or renal function, hypothyroidism, Addison's disease, prostatic hypertrophy or urethral stricture. The usual precautions should be observed and the possibility of respiratory depression should be kept in mind.

Cough Reflex: Hydrocodone suppresses the cough reflex; as with all narcotics, caution should be exercised when hydrocodone bitartrate and acetaminophen tablets are used postoperatively and in patients with pulmonary disease.

Information for Patients/Caregivers:

- Do not take Hydrocodone Bitartrate and Acetaminophen Tablets, USP if you are allergic to any of its ingredients.
- If you develop signs of allergy such as a rash or difficulty breathing stop taking Hydrocodone Bitartrate and Acetaminophen Tablets, USP and contact your healthcare provider immediately.
- Do not take more than 4000 milligrams of acetaminophen per day. Call your doctor if you took

more than the recommended dose.

Hydrocodone, like all narcotics, may impair the mental and/or physical abilities required for the performance of potentially hazardous tasks such as driving a car or operating machinery; patients should be cautioned accordingly.

Alcohol and other CNS depressants may produce an additive CNS depression, when taken with this combination product, and should be avoided.

Hydrocodone may be habit-forming. Patients should take the drug only for as long as it is prescribed, in the amounts prescribed, and no more frequently than prescribed.

Laboratory Tests:

In patients with severe hepatic or renal disease, effects of therapy should be monitored with serial liver and/or renal function tests.

Drug Interactions:

Patients receiving other narcotics, antihistamines, antipsychotics, antianxiety agents, or other CNS depressants (including alcohol) concomitantly with hydrocodone bitartrate and acetaminophen tablets may exhibit an additive CNS depression. When combined therapy is contemplated, the dose of one or both agents should be reduced.

The use of MAO inhibitors or tricyclic antidepressants with hydrocodone preparations may increase the effect of either the antidepressant or hydrocodone.

Drug/Laboratory Test Interactions:

Acetaminophen may produce false-positive test results for urinary 5-hydroxyindoleacetic acid.

Carcinogenesis, Mutagenesis, Impairment of Fertility:

No adequate studies have been conducted in animals to determine whether hydrocodone or acetaminophen have a potential for carcinogenesis, mutagenesis, or impairment of fertility.

Pregnancy:

Teratogenic Effects: *Pregnancy Category C:* There are no adequate and well-controlled studies in pregnant women. Hydrocodone bitartrate and acetaminophen tablets should be used during pregnancy only if the potential benefit justifies the potential risk to the fetus.

Nonteratogenic Effects: Babies born to mothers who have been taking opioids regularly prior to delivery will be physically dependent. The withdrawal signs include irritability and excessive crying, tremors, hyperactive reflexes, increased respiratory rate, increased stools, sneezing, yawning, vomiting, and fever. The intensity of the syndrome does not always correlate with the duration of maternal opioid use or dose. There is no consensus on the best method of managing withdrawal.

Labor and Delivery:

As with all narcotics, administration of this product to the mother shortly before delivery may result in some degree of respiratory depression in the newborn, especially if higher doses are used.

Nursing Mothers:

Acetaminophen is excreted in breast milk in small amounts, but the significance of its effects on nursing infants is not known. It is not known whether hydrocodone is excreted in human milk. Because many drugs are excreted in human milk and because of the potential for serious adverse reactions in nursing infants from hydrocodone and acetaminophen, a decision should be made whether to discontinue nursing or to discontinue the drug, taking into account the importance of the drug to the mother.

Pediatric Use:

Safety and effectiveness in pediatric patients have not been established.

Geriatric Use:

Clinical studies of hydrocodone bitartrate 5 mg and acetaminophen 500 mg did not include sufficient numbers of subjects aged 65 and over to determine whether they respond differently from younger subjects. Other reported clinical experience has not identified differences in responses between the elderly and younger patients. In general, dose selection for an elderly patient should be cautious, usually starting at the low end of the dosing range, reflecting the greater frequency of decreased hepatic, renal, or cardiac function, and of concomitant disease or other drug therapy.

ADVERSE REACTIONS

The most frequently reported adverse reactions are lightheadedness, dizziness, sedation, nausea and vomiting. These effects seem to be more prominent in ambulatory than in nonambulatory patients, and some of these adverse reactions may be alleviated if the patient lies down.

Other adverse reactions include:

Central Nervous System: Drowsiness, mental clouding, lethargy, impairment of mental and physical performance, anxiety, fear, dysphoria, psychic dependence, mood changes.

Gastrointestinal System: Prolonged administration of hydrocodone bitartrate and acetaminophen tablets may produce constipation.

Genitourinary System: Ureteral spasm, spasm of vesical sphincters and urinary retention have been reported with opiates.

Respiratory Depression: Hydrocodone bitartrate may produce dose-related respiratory depression by acting directly on the brain stem respiratory center (see **OVERDOSAGE**).

Special Senses: Cases of hearing impairment or permanent loss have been reported predominantly in patients with chronic overdose.

Dermatological: Skin rash, pruritus.

The following adverse drug events may be borne in mind as potential effects of acetaminophen: allergic reactions, rash, thrombocytopenia, agranulocytosis. Potential effects of high dosage are listed in the **OVERDOSAGE** section.

DRUG ABUSE AND DEPENDENCE**Controlled Substance:**

Hydrocodone Bitartrate and Acetaminophen Tablets are classified as a Schedule II controlled substance.

Abuse and Dependence:

Psychic dependence, physical dependence, and tolerance may develop upon repeated administration of narcotics; therefore, this product should be prescribed and administered with caution. However, psychic dependence is unlikely to develop when hydrocodone bitartrate and acetaminophen tablets are used for a short time for the treatment of pain.

Physical dependence, the condition in which continued administration of the drug is required to prevent the appearance of a withdrawal syndrome, assumes clinically significant proportions only after several weeks of continued narcotic use, although some mild degree of physical dependence may develop after a few days of narcotic therapy. Tolerance, in which increasingly large doses are required in order to

produce the same degree of analgesia, is manifested initially by a shortened duration of analgesic effect, and subsequently by decreases in the intensity of analgesia. The rate of development of tolerance varies among patients.

OVERDOSAGE

Following an acute overdosage, toxicity may result from hydrocodone or acetaminophen.

Signs and Symptoms:

Hydrocodone: Serious overdose with hydrocodone is characterized by respiratory depression (a decrease in respiratory rate and/or tidal volume, Cheyne-Stokes respiration, cyanosis), extreme somnolence progressing to stupor or coma, skeletal muscle flaccidity, cold and clammy skin, and sometimes bradycardia and hypotension. In severe overdosage, apnea, circulatory collapse, cardiac arrest and death may occur.

Acetaminophen: In acetaminophen overdosage: dose-dependent, potentially fatal hepatic necrosis is the most serious adverse effect. Renal tubular necrosis, hypoglycemic coma, and coagulation defects may also occur. Early symptoms following a potentially hepatotoxic overdose may include: nausea, vomiting, diaphoresis and general malaise. Clinical and laboratory evidence of hepatic toxicity may not be apparent until 48 to 72 hours post-ingestion.

Treatment: A single or multiple drug overdose with hydrocodone and acetaminophen is a potentially lethal polydrug overdose, and consultation with a regional poison control center is recommended. Immediate treatment includes support of cardiorespiratory function and measures to reduce drug absorption. Oxygen, intravenous fluids, vasopressors, and other supportive measures should be employed as indicated. Assisted or controlled ventilation should also be considered.

For hydrocodone overdose, primary attention should be given to the reestablishment of adequate respiratory exchange through provision of a patent airway and the institution of assisted or controlled ventilation. The narcotic antagonist naloxone hydrochloride is a specific antidote against respiratory depression which may result from overdosage or unusual sensitivity to narcotics, including hydrocodone. Since the duration of action of hydrocodone may exceed that of the antagonist, the patient should be kept under continued surveillance, and repeated doses of the antagonist should be administered as needed to maintain adequate respiration. A narcotic antagonist should not be administered in the absence of clinically significant respiratory or cardiovascular depression.

Gastric decontamination with activated charcoal should be administered just prior to N-acetylcysteine (NAC) to decrease systemic absorption if acetaminophen ingestion is known or suspected to have occurred within a few hours of presentation. Serum acetaminophen levels should be obtained immediately if the patient presents 4 hours or more after ingestion to assess potential risk of hepatotoxicity; acetaminophen levels drawn less than 4 hours post-ingestion may be misleading. To obtain the best possible outcome, NAC should be administered as soon as possible where impending or evolving liver injury is suspected. Intravenous NAC may be administered when circumstances preclude oral administration.

Vigorous supportive therapy is required in severe intoxication. Procedures to limit the continuing absorption of the drug must be readily performed since the hepatic injury is dose dependent and occurs early in the course of intoxication.

DOSAGE AND ADMINISTRATION

Dosage should be adjusted according to the severity of the pain and the response of the patient. However, it should be kept in mind that tolerance to hydrocodone can develop with continued use and that the incidence of untoward effects is dose related.

The usual adult dosage is one or two tablets every four to six hours as needed for pain. The total daily dosage should not exceed 8 tablets.

HOW SUPPLIED

Hydrocodone Bitartrate and Acetaminophen Tablets USP, contain Hydrocodone Bitartrate 5mg and Acetaminophen 325mg.

NDC 12634-264-00 Bottles of 10

NDC 12634-264-40 Bottles of 40

NDC 12634-264-50 Bottles of 50

NDC 12634-264-57 Blister Pack of 20

NDC 12634-264-59 Blister Pack of 30

NDC 12634-264-61 Blister pack of 10

NDC 12634-264-64 Blister Pack of 4

NDC 12634-264-65 Blister Pack of 5

NDC 12634-264-66 Blister Pack of 6

NDC 12634-264-71 Bottles of 30

NDC 12634-264-80 Bottles of 20

NDC 12634-264-82 Bottles of 12

NDC 12634-264-85 Bottles of 15

NDC 12634-264-91 Blister Pack of 1

NDC 12634-264-92 Bottles of 2

NDC 12634-264-94 Bottles of 4

NDC 12634-264-95 Bottles of 5

NDC 12634-264-96 Bottles of 6

Storage:

Store at controlled room temperature 15°-30°C (59°-86°F).

Dispense in a tight, light-resistant container with a child-resistant closure.

A Schedule CII Narcotic.

Manufactured by:

Mikart, Inc.

Atlanta, GA 30318

Repackaged and Distributed by:

Apotheca Inc.

Phoenix, AZ 85006 USA

Rev. 01/2014

Principal Display Panel

NDC 12634-264-71

CII

Hydrocodone

Bitartrate and

Acetaminophen

Tablets USP

5 mg/325 mg

See insert for full prescribing information.

Hydrocodone Bitartrate And Acetaminophen 5/325mg
NDC: 12634-264-71 30 Tablets CII
LOT: EXP:

Hydrocodone Bitartrate And Acetaminophen 5/325mg
NDC: 12634-264-71 30 Tablets CII
LOT: EXP:

Hydrocodone Bitartrate And Acetaminophen 5/325mg
NDC: 12634-264-71 30 Tablets CII
LOT: EXP:

Repackaged & Distributed by:
Apotheca, Inc Phoenix, AZ 85006

93522026471*

**Hydrocodone Bitartrate
And Acetaminophen 5/325mg**
30 Tablets CII
NDC: 12634-264-71

LOT: EXP:
COMPARE TO: VICODIN*
Rx Only
Dispensed In A Tight, Light Resistant
Container, As Defined in The
USP, Using A Child Resistant Closure.

WINDSOR PHARMACEUTICALS
APOTHECA, INC.
SPECIALISTS IN GENERIC PHARMACEUTICALS
Manufacturer • Distributor • Private Label
REV. DATE: 01/2014

BULK SOURCE DATA
Manufactured By:
Milkart, Inc.
Distributed By:
Watson Laboratories, Inc.

PRODUCT ID:
Debossed "WATSON" 3202
Bulk NDC: 0591-3202-01

Call your doctor for Medical Advice about Side Effects.
You may Report Side Effects to FDA at: 1-800-FDA-1088
May cause drowsiness. Alcohol may intensify this effect.
Use care when operating a car or dangerous machinery.
Federal law PROHIBITS the transfer of this drug to any
person other than the patient for whom it was prescribed.
Keep this and all Medication Out of the Reach of Children.
Not to exceed 12 tablets in a 24 hour period.

Directions:
Take ___ tablet(s) by mouth ___
times a day, as directed by physician.
Store at 20-25 C (68-77 F) [See USP Controlled
Room Temperature]. Protect from light and moisture.

HYDROCODONE BITARTRATE AND ACETAMINOPHEN

hydrocodone bitartrate and acetaminophen tablet

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:12634-264(NDC:0591-3202)
Route of Administration	ORAL	DEA Schedule	CII

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
HYDROCODONE BITARTRATE (UNII: NO70W886KK) (HYDROCODONE - UNII:6YK4Y3WQ7)	HYDROCODONE BITARTRATE	5 mg
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg

Inactive Ingredients

Ingredient Name	Strength
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
STARCH, CORN (UNII: O8232NY3SJ)	
CROSCARMELOLOSE SODIUM (UNII: M28OL1HH48)	

CROSPVIDONE (UNII: 68401960MK)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
POVIDONES (UNII: FZ989GH94E)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
SUCROSE (UNII: C151H8M554)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	

Product Characteristics

Color	orange (Specks) , white	Score	2 pieces
Shape	CAPSULE	Size	14mm
Flavor		Imprint Code	WATSON;3202
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:12634-264-00	10 in 1 BOTTLE; Type 0: Not a Combination Product	06/25/1997	
2	NDC:12634-264-40	40 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	06/25/1997	
3	NDC:12634-264-50	50 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	06/25/1997	
4	NDC:12634-264-57	20 in 1 BLISTER PACK; Type 0: Not a Combination Product	06/25/1997	
5	NDC:12634-264-59	30 in 1 BLISTER PACK; Type 0: Not a Combination Product	06/25/1997	
6	NDC:12634-264-61	10 in 1 BLISTER PACK; Type 0: Not a Combination Product	06/25/1997	
7	NDC:12634-264-64	4 in 1 BLISTER PACK; Type 0: Not a Combination Product	06/25/1997	
8	NDC:12634-264-65	5 in 1 BLISTER PACK; Type 0: Not a Combination Product	06/25/1997	
9	NDC:12634-264-66	6 in 1 BLISTER PACK; Type 0: Not a Combination Product	06/25/1997	
10	NDC:12634-264-71	30 in 1 BOTTLE; Type 0: Not a Combination Product	06/25/1997	
11	NDC:12634-264-80	20 in 1 BOTTLE; Type 0: Not a Combination Product	06/25/1997	
12	NDC:12634-264-82	12 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	06/25/1997	
13	NDC:12634-264-85	15 in 1 BOTTLE; Type 0: Not a Combination Product	06/25/1997	
14	NDC:12634-264-91	1 in 1 BLISTER PACK; Type 0: Not a Combination Product	06/25/1997	
15	NDC:12634-264-92	2 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	06/25/1997	
16	NDC:12634-264-94	4 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	06/25/1997	
17	NDC:12634-264-95	5 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	06/25/1997	
18	NDC:12634-264-96	6 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	06/25/1997	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA040099	06/25/1997	

Labeler - Apotheca Inc. (051457844)

Establishment

Name	Address	ID/FEI	Business Operations
Apotheca Inc.		051457844	repack(12634-264) , relabel(12634-264)

Revised: 5/2016

Apotheca Inc.