

**REDICARE NON-ASPIRIN 500MG- redicare non-aspirin, acetaminophen  
500mg tablet  
Redicare LLC**

**REDICARE NON-ASPIRIN 500MG**

***Drug Facts***

***Active Ingredient (in each tablet)***

Acetaminophen 500mg

***Purpose***

Pain reliever/fever reducer

***Uses:***

temporarily relieves minor aches and pains due to: ■ the common cold ■ headache ■ backache ■ minor pain of arthritis ■ toothache ■ muscular aches ■ premenstrual and menstrual cramps ■ temporarily reduces fever

***Warnings:***

**Liver warning:**

This product contains acetaminophen. Severe liver damage may occur if you take ■ more than 4,000mg of acetaminophen in 24 hours ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product

**Allergy alert:**

Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash

If a skin reaction occurs, stop use and seek medical help right away.

**Do not use**

■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.  
■ if you are allergic to acetaminophen or any of the inactive ingredients in this product

**Ask a doctor before use if you have**

liver disease

**Ask a doctor or pharmacist before use if**

you are taking the blood thinning drug warfarin

**Stop use and ask a doctor if**

■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days ■ new symptoms occur ■ redness or swelling is present. These could be signs of a serious condition

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

**Overdose warning:**

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

***Directions:***

**■ do not take more than directed (see overdose warning)**

**Adults and children 12 years and over:** ■ take 2 tablets every 6 hours while symptoms last ■ do not take more than 6 tablets in 24 hours, unless directed by a doctor ■ do not take for more than 10 days unless directed by a doctor

**Children under 12 years:** ask a doctor

***Other Information:***

■ store between 20°-25°C (68°-77°F)


■ do not use if imprinted safety seal under cap is broken or missing

***Inactive Ingredients:***

corn starch, povidone, pregelatinized starch, sodium starch glycolate, stearic acid

***Questions or Comments?***

Call 1-866-561-5650 Monday to Friday 9am-5pm EST



**Drug Facts**

**Active Ingredient (in each tablet)** Acetaminophen 500mg **Purpose** Pain reliever/fever reducer

**Uses:** temporarily relieves minor aches and pains due to: • the common cold • headache • backache • minor pain of arthritis • toothache • menstrual aches • premenstrual and menstrual cramps • temporarily reduces fever

**Warnings:**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take • more than 4,000mg of acetaminophen in 24 hours • with other drugs containing acetaminophen • 5 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include • skin redness • blisters • rash. If a skin reaction occurs, stop use and seek medical help right away.

**Do not use •** with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. • If you are allergic to acetaminophen or any of the inactive ingredients in this product.

**Ask a doctor before use if you have liver disease.**

**Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.**

**Stop use and ask a doctor if • pain gets worse or lasts more than 10 days • fever gets worse or lasts more than 3 days • new symptoms occur • redness or swelling is present. These could be signs of a serious condition.**

**Non-Aspirin**

**Pain Reliever/Fever Reducer**  
Similar to the ingredients in Tylenol® \*

**Drug Facts (Continued)**  
**Inactive Ingredients:** corn starch, povidone, pregelatinized starch, sodium starch glycolate, etc.  
**Questions or Comments?**  
Call 1-800-551-5650, Monday to Friday 9am-5pm EST

\*This product is not manufactured or distributed by McNeil Consumer Healthcare, owner of the registered trademark Tylenol.

**Redicare LLC**  
Cargens, NY 10820  
800.551.5650  
info@redicare.com  
www.redicare.com

NDC 719  
0109  
Made in U



**NON-ASPIRIN**  
2 TABLETS/PACKAGE

**Active ingredient:** (in each tablet)  
Acetaminophen 500mg.

**Purpose:** Pain reliever/fever reducer.

**Uses:** Temporarily relieves minor aches and pains due to: the common cold, headache, backache, minor pain of arthritis, toothache, muscular aches, premenstrual and menstrual cramps, temporarily reduces fever.

**Directions: Adults and children 12 years and over:** Take 2 tablets every 6 hours while symptoms last. Do not take more than 6 tablets in 24 hours, unless directed by a doctor. **Children under 12 years:** ask a doctor.

MANUFACTURED FOR REDICARE LLC, CONGERS, NY 10920

**PACKET NOT FOR INDIVIDUAL SALE**

FOR COMPLETE WARNINGS/CAUTIONS: SEE BOX

**WARNINGS: Liver Warning:** This product contains acetaminophen. Severe liver damage may occur if you take more than 4,000mg of acetaminophen in 24 hours, with other drugs containing acetaminophen, 3 or more alcoholic drinks every day while using this product. **Allergy Alert:** Acetaminophen may cause severe skin reactions. Symptoms may include: skin reddening, blisters, rash. If a skin reaction occurs, stop use and seek medical help right away. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Store between 20°-25°C (68°-77°F).



1.75"x1.075"  
Back

## REDICARE NON-ASPIRIN 500MG

redicare non-aspirin, acetaminophen 500mg tablet

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:71105-011
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	500 mg

### Inactive Ingredients

Ingredient Name	Strength
<b>STARCH, CORN</b> (UNII: O8232NY3SJ)	
<b>POVIDONE</b> (UNII: FZ989GH94E)	
<b>SODIUM STARCH GLYCOLATE TYPE A CORN</b> (UNII: AG9B65PV6B)	

STEARIC ACID (UNII: 4ELV7Z65AP)

Product Characteristics

Color	white	Score	no score
Shape	ROUND	Size	12mm
Flavor		Imprint Code	AZ;011
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:71105-011-02	50 in 1 BOX	02/27/2025	
1	NDC:71105-011-01	2 in 1 PACKET; Type 0: Not a Combination Product		



## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	02/27/2025	

**Labeler** - Redicare LLC (800149346)

Revised: 3/2025

Redicare LLC