

IBUPROFEN- ibuprofen capsule, liquid filled
BJWC (Berkley & Jensen / BJ's)

Bjs Ibuprofen Soft Gelatin Capsules 200 mg

Drug Facts

Active ingredient (in each capsule)

Solubilized ibuprofen equal to

200 mg ibuprofen (NSAID)*

(present as the free acid and potassium salt)

*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever/Fever reducer

Uses

- temporarily relieves minor aches and pains due to:
 - headache
 - toothache
 - backache
 - menstrual cramps
 - the common cold
 - muscular aches
 - minor pain of arthritis
- temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems

- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
 - slurred speech
 - leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- **do not take more than directed**
- **the smallest effective dose should be used**
- adults and children 12 years and over: take 1 capsule every 4 to 6 hours while symptoms persist
- if pain or fever does not respond to 1 capsule, 2 capsules may be used
- do not exceed 6 capsules in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

Other information

- each capsule contains: **potassium 20 mg**
- read all warnings and directions before use
- store at 20-25°C (68-77°F)
- avoid excessive heat above 40°C (104°F).

Inactive ingredients

ammonium hydroxide, FD&C green no. 3, gelatin, iron oxide black, medium chain triglycerides, polyethylene glycol, potassium hydroxide, propylene glycol, purified water, shellac, sorbitan monooleate, sorbitol sorbitan

Questions or comments? Call **1-877-290-4008**

COMPARE TO THE ACTIVE INGREDIENT IN ADVIL® LIQUID-GELS[®]

NDC 68391-610-99

berkley jensen.

480
SOFTGELS[™]

IBUPROFEN
CAPSULES, 200mg

PAIN RELIEVER/
FEVER REDUCER (NSAID)

ACTUAL SIZE

480
SOFTGELS[™]

([™]LIQUID-FILLED CAPSULES)

Drug Facts

Active ingredient (in each capsule) Purpose
Sulfonated ibuprofen (NSAID) 200 mg (present as the free acid and potassium salt)
Pain reliever/fever reducer
*nonsteroidal anti-inflammatory drug

Uses
temporarily relieves minor aches and pains due to:
headache toothache backache
menstrual cramps the common cold
muscular aches minor pain of arthritis
temporarily reduces fever

Warnings
Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:
hives facial swelling asthma (wheezing)
shock skin reddening rash blisters
If an allergic reaction occurs, stop use and seek medical help right away.
Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:
are age 60 or older
have had stomach ulcers or bleeding problems
take a blood thinning (anticoagulant) or steroid drug
take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
have 3 or more alcoholic drinks every day while using this product
take more or for a longer time than directed

Drug Facts (continued)
Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.
Do not use if you have ever had an allergic reaction to any other pain reliever/fever reducer right before or after heart surgery
Ask a doctor before use if you have problems or serious side effects from taking pain relievers or fever reducers you have a history of stomach problems, such as heartburn you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke you are taking a diuretic
Ask a doctor or pharmacist before use if you are under a doctor's care for any serious condition taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin taking any other drug
When using this product take with food or milk if stomach upset occurs
Stop use and ask a doctor if you experience any of the following signs of stomach bleeding:
feel faint vomit blood have bloody or black stools
have stomach pain that does not get better
you have symptoms of heart problems or stroke:
chest pain trouble breathing
weakness in one part or side of body
slurred speech big swelling
pain gets worse or lasts more than 10 days
fever gets worse or lasts more than 3 days

Drug Facts (continued)
redness or swelling is present in the painful area
any new symptoms appear
If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions do not take more than directed
the smallest effective dose should be used
adults and children 12 years and over: take 1 capsule every 4 to 6 hours while symptoms persist if pain or fever does not respond to 1 capsule, 2 capsules may be used do not exceed 6 capsules in 24 hours, unless directed by a doctor children under 12 years: ask a doctor

Other information each capsule contains: potassium 20 mg
read all warnings and directions before use store at 20-25°C (68-77°F) avoid excessive heat above 40°C (104°F).

Inactive ingredients ammonium hydroxide, FD&C green no.3, gelatin, iron oxide black, medium chain triglycerides, polyethylene glycol, potassium hydroxide, propylene glycol, purified water, shellac, sorbitan monooleate, sorbitol sorbitan

Questions or comments? Call 1-877-290-4008

TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FILM SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING

LOT: 610725
EXP: 8/2025

V000
88670-20401-9
610725
610725
MADE IN INDIA
Varnish Omit Area

IBUPROFEN

ibuprofen capsule, liquid filled

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:68391-610
Route of Administration	ORAL		

Active Ingredient/Active Moiety				
Ingredient Name			Basis of Strength	Strength
IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)			IBUPROFEN	200 mg
Inactive Ingredients				
Ingredient Name				Strength
AMMONIA (UNII: 5138Q19F1X)				
WATER (UNII: 059QF0KO0R)				
SHELLAC (UNII: 46N107B71O)				
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)				
FD&C GREEN NO. 3 (UNII: 3P3ONR6O1S)				
FERROSOFERRIC OXIDE (UNII: XM0M87F357)				
SORBITAN MONOOLEATE (UNII: 06XEA2VD56)				
POTASSIUM HYDROXIDE (UNII: WZH3C48M4T)				
GELATIN, UNSPECIFIED (UNII: 2G86QN327L)				
MEDIUM-CHAIN TRIGLYCERIDES (UNII: C9H2L21V7U)				
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)				
SORBITOL (UNII: 506T60A25R)				
Product Characteristics				
Color	green	Score	no score	
Shape	CAPSULE	Size	19mm	
Flavor		Imprint Code	133	
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:68391-610-99	480 in 1 BOTTLE; Type 0: Not a Combination Product	08/20/2025	
Marketing Information				
Marketing Category	Application Number or Monograph Citation		Marketing Start Date	Marketing End Date
ANDA	ANDA079205		08/20/2025	

Labeler - BJWC (Berkley & Jensen / BJ's) (159082692)

Registrant - TIME CAP LABORATORIES, INC. (037052099)

Establishment

Name	Address	ID/FEI	Business Operations
MARKSANS PHARMA LIMITED		925822975	manufacture(68391-610)

Revised: 8/2025

BJWC (Berkley & Jensen / BJ's)