LOW DOSE ASPIRIN- aspirin tablet, delayed release BJWC (Berkley & Jensen / BJ's)

BJs Low Dose Aspirin 81 mg Tablets

Drug Facts

Active ingredient (in each tablet)

Aspirin 81 mg (NSAID)*

*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever

Uses

- for the temporary relief of minor aches and pains or as recommended by your doctor. Because of its delayed action, this product will not provide fast relief of headaches or other symptoms needing immediate relief.
- ask your doctor about other uses for enteric coated 81 mg Aspirin.

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Do not use

- if you are allergic to aspirin or any other pain reliever/fever reducer
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn

- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma

Ask a doctor or pharmacist before use if you are taking a prescription drug for

- gout
- diabetes
- arthritis

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
- feel faint
- vomit blood
- have bloody or black stools
- have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- ringing in the ears or a loss of hearing occurs

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- drink a full glass of water with each dose
- adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor
- children under 12 years: consult a doctor

Other information

- store at 25°C (77°F) excursions permitted between 15°-30°C (59°-86°F)
- use by expiration date on package
- avoid excessive heat above 40°C (104°F)

Inactive ingredients anhydrous lactose, carnauba wax, colloidal silicon dioxide, croscarmellose sodium, D&C yellow #10 aluminum lake, iron oxide ochre, methacrylic acid and ethyl acrylate copolymer, microcrystalline cellulose, polysorbate 80, simethicone, sodium hydroxide, sodium lauryl sulfate, starch, talc, titanium dioxide, triethyl citrate

Questions or comments? Call **1-877-290-4008**

481R 365ct Label





THE IMPRINTED FOIL SEAI 4, broken or missing PRODUCT IF THE IS CUT, TORN, E NGS AND F COMPLETE WARNI TAMPER EVIDENT: DO NOT USE THIS OVER THE MOUTH OF THE BOTTLE RETAIN (

Drug Facts

*nonsteroidal anti-inflammatory drug

USSE ■ for the temporary relief of minor aches and pains or as recommended
by your doctor. Because of its delayed action, this product will not provide
trast relief of headaches or other symptoms needing immediate relief.

■ ask your doctor about other uses for enteric coated 81 mg Aspirin. ctive ingredient (in each tablet)

Warnings

Reys's syndrome. Children and teenagers who have or are recovering from chicken pox or fill-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reys's syndrome, a rare but senfous lilenss.

Alengy after. Aspirin may cause a severe allergic reaction which may include:

■ hives ■ facial swelling ■ asthma (wheezing) ■ shock

Sumach bleeding warning: This product contains an NSAID, which may cause severe storach beleding. This product contains an NSAID, which may cause severe storach beleding. The chance is higher if you ■ are age 60 or older any card and stomach ulcers or bleeding problems ■ take a blood thinning affanticoagularin or steroid dring. ■ take dher drugs containing prescription or nonprescription NSAIDs (aspirit, hupproten, napproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

If pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Kep our of reach of children.

Directions ■ drink all lights or whater in the control of the cause it may cause problems in the unborn child or complications during delivery.

Kep our of reach of children.

Directions ■ chink a full glass of water with each dose ■ adults and children 12 years and over, take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours units glereited by a doctor ■ children under 12 years; consult a doctor Other information ■ store at 125°.7(7°F) excursions permitted between 15°.30°C (59°-86°F) ■ use by expiration date on package ■ avoid excessive heat above 40°C (104°F) Questions of comments? Call 1-877-290-4008

Questions of comments? Call 1-877-290-4008

Bothorough, MA 01752

800-8JS-CLUB

342613 481R 0725

481R 0725

342613

/arnish Omit Area

LOT:

481R 730ct Carton



LOW DOSE ASPIRIN

aspirin tablet, delayed release

Product Information				
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:68391-481	
Route of Administration	ORAL			

Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	81 mg	

Inactive Ingredients	
Ingredient Name	Strength
POLYSORBATE 80 (UNII: 6OZP39ZG8H)	
DIMETHICONE, UNSPECIFIED (UNII: 92RU3N3Y10)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
FERRIC OXIDE YELLOW (UNII: EX43802MRT)	
D&C YELLOW NO. 10 ALUMINUM LAKE (UNII: CQ3XH3DET6)	
METHACRYLIC ACID AND ETHYL ACRYLATE COPOLYMER (UNII: NX76LV5T8J)	
TALC (UNII: 7SEV7J4R1U)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
SODIUM HYDROXIDE (UNII: 55X04QC32I)	
ANHYDROUS LACTOSE (UNII: 3SY5LH9PMK)	
CARNAUBA WAX (UNII: R12CBM0EIZ)	
CROSCARMELLOSE SODIUM (UNII: M280L1HH48)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
TRIETHYL CITRATE (UNII: 8Z96QXD6UM)	
STARCH, CORN (UNII: 08232NY3SJ)	
SODIUM LAURYL SULFATE (UNII: 368GB5141J)	

Product Characteristics				
Color	yellow	Score	no score	
Shape	ROUND	Size	7mm	
Flavor		Imprint Code	Heart	
Contains				

P	Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date	
1	NDC:68391-481- 93	2 in 1 CARTON	08/25/2025		
1	NDC:68391-481- 65	365 in 1 BOTTLE; Type 0: Not a Combination Product			

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	08/25/2025	

Labeler - BJWC (Berkley & Jensen / BJ's) (159082692)

Registrant - TIME CAP LABORATORIES, INC (037052099)

Establishment				
Name	Address	ID/FEI	Business Operations	
Time Cap Laboratories, Inc.		037052099	manufacture(68391-481)	

Revised: 8/2025 BJWC (Berkley & Jensen / BJ's)