

NORETHINDRONE ACETATE- norethindrone acetate tablet
Ingenus Pharmaceuticals, LLC


Norethindrone acetate tablets, USP

Rx only

DESCRIPTION

Norethindrone acetate tablets, USP - 5 mg oral tablets.

Norethindrone acetate, USP (17-hydroxy-19-nor-17 α -pregn-4-en-20-yn-3-one acetate), a synthetic, orally active progestin, is the acetic acid ester of norethindrone. It is a white to yellowish-white, crystalline powder.



Norethindrone acetate tablets, USP contain the following inactive ingredients: lactose monohydrate, magnesium stearate, and microcrystalline cellulose.

Meets USP dissolution Test 2.

CLINICAL PHARMACOLOGY

Norethindrone acetate induces secretory changes in an estrogen-primed endometrium. On a weight basis, it is twice as potent as norethindrone.


PHARMACOKINETICS

Absorption:

Norethindrone acetate is completely and rapidly deacetylated to norethindrone (NET) after oral administration, and the disposition of norethindrone acetate is indistinguishable from that of orally administered norethindrone. Norethindrone acetate is rapidly absorbed from norethindrone acetate tablets, with maximum plasma concentration of norethindrone generally occurring at about 2 hours post-dose. The pharmacokinetic parameters of norethindrone following single oral administration of norethindrone acetate tablets in 29 healthy female volunteers are summarized in Table 1.

Table 1	
Pharmacokinetic Parameters after a Single Dose of norethindrone acetate tablets in Healthy Women	
Norethindrone acetate tablets (n=29) Arithmetic Mean \pm SD	
Norethindrone (NET)	
AUC(0-inf)(ng/mL*h)	166.90 \pm 56.28
C _{max} (ng/mL)	26.19 \pm 6.19
t _{max} (h)	1.83 \pm 0.58
t _{1/2} (h)	8.51 \pm 2.19
AUC= area under the curve	
C _{max} = maximum plasma concentration	
t _{max} = time at maximum plasma concentration	
t _{1/2} = half-life	
SD = standard deviation	

Figure 1. Mean Plasma Concentration Profile after a Single Dose of 5 mg Administered to 29 Healthy Female Volunteers under Fasting Conditions



Effect of Food:

The effect of food administration on the pharmacokinetics of norethindrone acetate tablets has not been studied.

Distribution:

Norethindrone is 36% bound to sex hormone-binding globulin (SHBG) and 61% bound to albumin. Volume of distribution of norethindrone is about 4 L/kg.

Metabolism:

Norethindrone undergoes extensive biotransformation, primarily via reduction, followed by sulfate and glucuronide conjugation. The majority of metabolites in the circulation are sulfates, with glucuronides accounting for most of the urinary metabolites.

Excretion:

Plasma clearance value for norethindrone is approximately 0.4 L/hr/kg. Norethindrone is excreted in both urine and feces, primarily as metabolites. The mean terminal elimination half-life of norethindrone following a single dose administration of norethindrone acetate tablets is approximately 9 hours.

SPECIAL POPULATIONS

Geriatrics

The effect of age on the pharmacokinetics of norethindrone after norethindrone acetate tablets administration has not been evaluated.

Race

The effect of race on the disposition of norethindrone after norethindrone acetate tablets administration has not been evaluated.

Renal Insufficiency

The effect of renal disease on the disposition of norethindrone after norethindrone acetate tablets administration has not been evaluated. In premenopausal women with chronic renal failure undergoing peritoneal dialysis who received multiple doses of an oral contraceptive containing ethinyl estradiol and norethindrone, plasma norethindrone concentration was unchanged compared to concentrations in premenopausal women with normal renal function.

Hepatic Insufficiency

The effect of hepatic disease on the disposition of norethindrone after norethindrone acetate tablets administration has not been evaluated. However, norethindrone acetate tablets are contraindicated in markedly impaired liver function or liver disease.

Drug Interactions

No pharmacokinetic drug interaction studies investigating any drug-drug interactions with norethindrone acetate tablets have been conducted.

INDICATIONS AND USAGE

Norethindrone acetate tablets are indicated for the treatment of secondary amenorrhea, endometriosis, and abnormal uterine bleeding due to hormonal imbalance in the absence of organic pathology, such as submucous fibroids or uterine cancer.

Norethindrone acetate tablets are not intended, recommended or approved to be used with concomitant estrogen therapy in postmenopausal women for endometrial protection.

CONTRAINDICATIONS

- Known or suspected pregnancy. There is no indication for norethindrone acetate tablets in pregnancy. (See **PRECAUTIONS**.)
- Undiagnosed vaginal bleeding
- Known, suspected or history of cancer of the breast
- Active deep vein thrombosis, pulmonary embolism or history of these conditions
- Active or recent (e.g., within the past year) arterial thromboembolic disease (e.g., stroke, myocardial infarction)
- Impaired liver function or liver disease
- As a diagnostic test for pregnancy
- Hypersensitivity to any of the drug components

WARNINGS

1 Cardiovascular disorders

Patients with risk factors for arterial vascular disease (e.g., hypertension, diabetes mellitus, tobacco use, hypercholesterolemia, and obesity) and/or venous thromboembolism (e.g., personal history or family history of VTE, obesity, and systemic lupus erythematosus) should be managed appropriately.

2 Visual abnormalities

Discontinue medication pending examination if there is a sudden partial or complete loss of vision or if there is sudden onset of proptosis, diplopia, or migraine. If examination reveals papilledema or retinal vascular lesions, medication should be discontinued.

PRECAUTIONS

1 General Precautions

- Because this drug may cause some degree of fluid retention, conditions which might be influenced by this factor, such as epilepsy, migraine, cardiac or renal dysfunctions, require careful observation
- In cases of breakthrough bleeding, and in all cases of irregular bleeding per vagina, nonfunctional causes should be borne in mind. In cases of undiagnosed vaginal bleeding, adequate diagnostic measures are indicated
- Patients who have a history of clinical depression should be carefully observed and the drug discontinued if the depression recurs to a serious degree
- Data suggest that progestin therapy may have adverse effects on lipid and carbohydrate metabolism. The choice of progestin, its dose, and its regimen may be important in minimizing these adverse effects, but these issues will require further study before they are clarified. Women with hyperlipidemias and/or diabetes should be monitored closely during progestin therapy
- The pathologist should be advised of progestin therapy when relevant specimens are submitted

2 Information for the Patient

Healthcare providers are advised to discuss the PATIENT INFORMATION leaflet with patients for whom they prescribe norethindrone acetate tablets.

3 Drug/Laboratory Tests Interactions

The following laboratory test results may be altered by the use of estrogen/progestin combination drugs:

1. Accelerated prothrombin time, partial thromboplastin time, and platelet aggregation time; increased platelet count; increased factors II, VII antigen, VIII antigen, VIII coagulant activity, IX, X, XII, VII-X complex, II-VII-X complex, and beta-thromboglobulin; decreased levels of antifactor X_a and antithrombin III, decreased antithrombin III activity; increased levels of fibrinogen and fibrinogen activity; increased plasminogen antigen and activity.
2. Increased thyroid-binding globulin (TBG) levels leading to increased circulating total thyroid hormone levels as measured by protein-bound iodine (PBI), T₄ levels (by column or by radio immunoassay) or T₃ levels by radio immunoassay. T₃ resin uptake is decreased, reflecting the elevated TBG. Free T₄ and free T₃ concentrations are unaltered. Patients on thyroid replacement therapy may require higher doses of thyroid hormone.
3. Other binding proteins may be elevated in serum (i.e., corticosteroid binding globulin (CBG), sex hormone binding globulin (SHBG)) leading to increased circulating corticosteroid and sex steroids, respectively. Free or biologically active hormone concentrations are unchanged. Other plasma proteins may be increased (angiotensinogen/renin substrate, alpha-1-antitrypsin, ceruloplasmin).
4. Increased plasma HDL and HDL₂ cholesterol subfraction concentrations, reduced LDL cholesterol concentration, increased triglycerides levels.
5. Impaired glucose metabolism.

6. Reduced response to metyrapone test.

4 Carcinogenesis, Mutagenesis, and Impairment of Fertility

Some beagle dogs treated with medroxyprogesterone acetate developed mammary nodules. Although nodules occasionally appeared in control animals, they were intermittent in nature, whereas nodules in treated animals were larger and more numerous, and persisted. There is no general agreement as to whether the nodules are benign or malignant. Their significance with respect to humans has not been established.

5 Pregnancy Category X

Norethindrone acetate is contraindicated during pregnancy as it may cause fetal harm when administered to pregnant women. Several reports suggest an association between intrauterine exposure to progestational drugs in the first trimester of pregnancy and congenital abnormalities in male and female fetuses. Some progestational drugs induce mild virilization of the external genitalia of female fetuses.

6 Nursing Mothers

Detectable amounts of progestins have been identified in the milk of mothers receiving them. Caution should be exercised when progestins are administered to a nursing woman.

7 Pediatric Use

Norethindrone acetate tablets are not indicated in children.

ADVERSE REACTIONS

See **WARNINGS** and **PRECAUTIONS**

The following adverse reactions have been observed in women taking progestins:

- Breakthrough bleeding
- Spotting
- Change in menstrual flow
- Amenorrhea
- Edema
- Changes in weight (decreases, increases)
- Changes in the cervical squamo-columnar junction and cervical secretions
- Cholestatic jaundice
- Rash (allergic) with and without pruritus
- Melasma or chloasma
- Clinical depression
- Acne
- Breast enlargement/tenderness
- Headache/migraine
- Urticaria
- Abnormalities of liver tests (i.e., AST, ALT, Bilirubin)
- Decreased HDL cholesterol and increased LDL/HDL ratio
- Mood swings
- Nausea
- Insomnia
- Anaphylactic/anaphylactoid reactions
- Thrombotic and thromboembolic events (e.g., deep vein thrombosis, pulmonary embolism, retinal vascular thrombosis, cerebral thrombosis and embolism)
- Optic neuritis (which may lead to partial or complete loss of vision)

To report SUSPECTED ADVERSE REACTIONS, please call Ingenus Pharmaceuticals, LLC toll-free at 1-877-748-1970 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

DOSAGE AND ADMINISTRATION

Therapy with norethindrone acetate tablets must be adapted to the specific indications and therapeutic response of the individual patient.

Secondary amenorrhea, abnormal uterine bleeding due to hormonal imbalance in the absence of organic pathology: 2.5 to 10 mg norethindrone acetate tablets may be given daily for 5 to 10 days to produce secretory transformation of an endometrium that has been adequately primed with either endogenous or exogenous estrogen.

Progestin withdrawal bleeding usually occurs within three to seven days after discontinuing norethindrone acetate tablets therapy. Patients with a past history of recurrent episodes of abnormal uterine bleeding may benefit from planned menstrual

cycling with norethindrone acetate tablets.

Endometriosis: Initial daily dosage of 5 mg norethindrone acetate tablets for two weeks. Dosage should be increased by 2.5 mg per day every two weeks until 15 mg per day of norethindrone acetate tablets is reached. Therapy may be held at this level for six to nine months or until annoying breakthrough bleeding demands temporary termination.

HOW SUPPLIED

Norethindrone acetate tablets, USP are available as:

5 mg: White to off-white, oval, flat beveled edged tablet. 'P658' is engraved on one side, and a breakline on the reverse.

Available as follows:

Bottle of 50	NDC	50742-267-50
Bottle of 500	NDC	50742-267-05

Store at 20° to 25°C (68° to 77°F) [See USP Controlled Room Temperature].

Rx Only

Revised: 05/2019

I0093

Rev B

Manufactured for:

Ingenus Pharmaceuticals, LLC
Orlando, FL 32839-6408

Made in China

ingenus

PATIENT INFORMATION

Norethindrone acetate

**(nor-eth-IN-drone as-i-teyt)
tablets, USP**

Read this PATIENT INFORMATION before you start taking norethindrone acetate tablets and read what you get each time you refill norethindrone acetate tablets. There may be new information. This information does not take the place of talking to your healthcare provider about your medical condition.

What is the most important information I should know about norethindrone acetate (A Progestin Hormone) tablets?

- **Do not use norethindrone acetate tablets if you are pregnant, breast-feeding or are trying to conceive.**
- **Do not use norethindrone acetate tablets if you have had a previous blood clot, stroke, or heart attack.**
- **Do not use norethindrone acetate tablets if you are postmenopausal.**

What are norethindrone acetate tablets?

Norethindrone acetate tablets are similar to the progesterone hormones naturally produced by the body.

What are norethindrone acetate tablets used for?

Norethindrone acetate tablets are used for the treatment of secondary amenorrhea (absence of menstrual periods in women who have previously had a menstrual period who are not pregnant), the treatment of endometriosis, and the treatment of irregular menstrual periods due to hormone imbalance.

Who should not take norethindrone acetate tablets?

You should not take norethindrone acetate tablets if you are postmenopausal, pregnant or breastfeeding.

You should not take norethindrone acetate tablets if you have the following conditions:

- **Known or suspected pregnancy.** Norethindrone acetate tablets are not indicated during pregnancy as it may cause fetal harm when administered to pregnant women. There is an increased risk of minor birth defects in children whose mothers take norethindrone acetate tablets during the first 4 months of pregnancy (mild masculinization of the external genitalia of the female fetus, as well as hypospadias in the male fetus). If you take norethindrone acetate tablets and later find out you were pregnant, talk with your healthcare provider right away

- **History of blood clots in the legs, lungs, eyes, brain, or elsewhere, or a past history of these conditions**
- **Liver impairment or disease**
- **Known or suspected cancer of the breast.** If you have or had cancer of the breast, talk with your healthcare provider about whether you should take norethindrone acetate tablets
- **Undiagnosed vaginal bleeding**
- **Hypersensitivity to norethindrone acetate tablets.** See the end of this leaflet for a list of all of the ingredients in norethindrone acetate tablets

What are the risks associated with norethindrone acetate tablets?

- **Risk to the Fetus:**

Norethindrone acetate tablets should not be used if you are pregnant. Norethindrone acetate tablets are contraindicated during pregnancy as it may cause fetal harm when administered to pregnant women. There is an increased risk of minor birth defects in children whose mothers take this drug during the first 4 months of pregnancy. Several reports suggest an association between mothers who take these drugs in the first trimester of pregnancy and congenital abnormalities in male and female babies. Although it is not clear that these events were drug related, you should check with your healthcare provider about the risks to your unborn child of any medication taken during pregnancy.

You should avoid using norethindrone acetate tablets during pregnancy. If you take norethindrone acetate tablets and later find you were pregnant when you took it, be sure to discuss this with your healthcare provider as soon as possible.

- **Abnormal Blood Clotting:**

Use of progestational drugs, such as norethindrone acetate tablets, has been associated with changes in the blood-clotting system. These changes allow the blood to clot more easily, possibly allowing clots to form in the bloodstream. If blood clots do form in your bloodstream, they can cut off the blood supply to vital organs, causing serious problems. These problems may include a stroke (by cutting off blood to part of the brain), a heart attack (by cutting off blood to part of the heart), a pulmonary embolus (by cutting off blood to part of the lungs), visual loss or blindness (by cutting off blood vessels in the eye), or other problems. Any of these conditions may cause death or serious long-term disability. Call your healthcare provider right away if you suspect you have any of these conditions. He or she may advise you to stop using the drug.

- **Eye Abnormalities:**

Discontinue norethindrone acetate tablets and call your healthcare provider right away if you experience sudden partial or complete loss of vision, blurred vision, or sudden onset of bulging eyes, double vision, or migraine.

These are some of the warning signs of serious side effects with progestin therapy:

- Breast lumps
- Dizziness and faintness
- Changes in speech
- Severe headaches
- Chest pain
- Shortness of breath
- Pains in your legs
- Changes in vision

Call your healthcare provider right away if you get any of these warning signs, or any other unusual symptom that concerns you.

Common side effects include:

- Headache
- Breast pain
- Irregular vaginal bleeding or spotting
- Stomach/abdominal cramps/bloating
- Nausea and vomiting
- Hair loss

Other side effects include:

- High blood pressure
- Liver problems
- High blood sugar
- Fluid retention
- Enlargements of benign tumors of the uterus (“fibroids”)

- Vaginal yeast infections
- Mental depression

These are not all the possible side effects of progestin and/or estrogen therapy. For more information, ask your healthcare provider or pharmacist.

What can I do to lower my chances of getting a serious side effect with norethindrone acetate tablets?

- Talk with your healthcare provider regularly about whether you should continue taking norethindrone acetate tablets
- Have a breast exam and mammogram (breast x-ray) every year unless your healthcare provider tells you something else. If members of your family have had breast cancer or if you have ever had breast lumps or an abnormal mammogram, you may need to have breast exams more often
- If you have high blood pressure, high cholesterol (fat in the blood), diabetes, are overweight, or if you use tobacco, you may have higher chances for getting heart disease. Ask your healthcare provider for ways to lower your chances of getting heart attacks

General information about the safe and effective use of norethindrone acetate tablets

Medicines are sometimes prescribed for conditions that are not mentioned in patient information leaflets. Do not take norethindrone acetate tablets for conditions for which it was not prescribed. Do not give norethindrone acetate tablets to other people, even if they have the same symptoms you have. It may harm them.

Keep norethindrone acetate tablets out of the reach of children.

This leaflet provides a summary of the most important information about progestin and/or estrogen therapy. If you would like more information, talk with your healthcare provider or pharmacist. You can ask for information about norethindrone acetate tablets that is written for health professionals.

What are the ingredients in norethindrone acetate tablets?

Norethindrone acetate tablets contain the following inactive ingredients: lactose monohydrate, magnesium stearate, and microcrystalline cellulose

Rx Only

Revised: 05/2019

10094

Rev B

Manufactured for:

Ingenus Pharmaceuticals, LLC
Orlando, FL 32839-6408

Made in China

PACKAGE LABEL.PRINCIPAL DISPLAY PANEL

ingenus

NDC 50742-267-50

Norethindrone Acetate Tablets, USP

5 mg

ORALLY ACTIVE PROGESTIN

PHARMACIST: PLEASE DISPENSE WITH ATTACHED PATIENT INFORMATION LEAFLET

Rx only

50 Tablets

Manufactured for:
Ingenus Pharmaceuticals, LLC
Orlando, FL 32839-6408
Madelin, China.

ingenus

NDC 50742-267-50

Norethindrone Acetate Tablets, USP

5 mg

ORALLY ACTIVE PROGESTIN

PHARMACIST: PLEASE DISPENSE WITH ATTACHED PATIENT INFORMATION LEAFLET

Rx only

267-50 v1/05-2019 50 Tablets Rev. B Iss. 05/2019 P0218

Each tablet contains:
Norethindrone acetate, USP.....5 mg

Usual Dosage: See package Insert for full prescribing Information.

Store at 20° to 25°C (68° to 77°F); [See USP Controlled Room Temperature].

Dispense in a tight, light-resistant container as defined in the USP, with a child-resistant closure (as required).

KEEP THIS AND ALL MEDICATIONS OUT OF THE REACH OF CHILDREN.

P02180519

3 50742-267-50 7

GTIN: 00350742267507

NORETHINDRONE ACETATE

norethindrone acetate tablet

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:50742-267
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
NORETHINDRONE ACETATE (UNII: 9S44LIC7OJ) (NORETHINDRONE - UNII:T18F433X4S)	NORETHINDRONE ACETATE	5 mg

Inactive Ingredients

Ingredient Name	Strength
LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	

Product Characteristics

Color	WHITE (white to off-white)	Score	2 pieces
Shape	OVAL (flat, beveled-edge)	Size	11mm
Flavor		Imprint Code	P658
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:50742-267-50	50 in 1 BOTTLE; Type 0: Not a Combination Product	08/29/2019	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA206490	08/29/2019	

Labeler - Ingenus Pharmaceuticals, LLC (833250017)

Registrant - Novast Laboratories, Ltd. (527695995)

Establishment

Name	Address	ID/FEI	Business Operations
Novast Laboratories Ltd.		527695995	ANALYSIS(50742-267) , label(50742-267) , manufacture(50742-267) , pack(50742-267)

Revised: 12/2023

Ingenus Pharmaceuticals, LLC