# HAN SHIN GAMKONAL EXTRACT GRANULES- dry ginger, glycyrrhiza, peony root granule MISOCOS

Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click here.

-----

## **Drug Facts**

Dry Ginger 1.90g

Glycyrrhiza 1.90g

Peony Root 1.90g

corn starch, Lactose Hydrate

- ■Bronchitis, Bronchial asthma, Coryza, Runny nose, Cough with dilute phlegm, rhinitis keep out of reach of the children
- Take a packet of medicine (one dose) of an adult three times a day before or between meals.

Warnings

When using this product

- ■Do not use in patients with genetic problems such as galactose intolerance, Lapp lactase deficiency, or glucose-galactose malabsorption since this drug contains lactose.
- ■Do not use it for more than 1 month unless the symptoms improve. Long-term use should only be done under medical supervision.
- ■When used in combination with potassium-containing agents, licorice-containing drugs, and diuretics (e.g., furosemide, ethacrynic acid, trichlormethiazide), myopathy is likely to occur, so take it carefully.
- ■When combined with MaHuang or ephedrine-containing agents, MAO inhibitors, thyroid agents(thyroxine, liothyronine), catecholamine agents(epinephrine, isoprenaline), and xanthin agents(theophylline, dipropylene), symptoms such as insomnia, excessive sweat, rapid pulse, frequent urination, palpitations, general weakness, and mental excitement are likely to appear, so take them carefully.

Ask a doctor before use if you have high blood pressure, edema, heart or kidney problems, rash, redness, diabetic, another disease.

Stop use and ask a doctor if the following serious side effects occur. These may be signs of a serious condition.

- ■Gastric aldosteronism (decreased urine, swollen face, hands, feet, eyelids, etc.)
- ■Myopathy (muscle weakness, cramps, paralysis)
- ■Interstitial pneumonia (fever, cough, shortness of breath, lung sound abnormalities)
- ■Liver disorders (elevated liver enzymes, jaundice)

- ■Skin reactions (rash, hives)
- Digestive issues (loss of appetite, stomach discomfort)

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children, In case of overdose, get medical help or contact a poison control center right away

for oral use only

HAN SHIN	GAMK	NAL Fxt	ract Granules

Drug Facts			
Active Ingred	ients (in each dosage unit 3.5g)	Purpose	
Dry Ginger	1.90g	Antiinflammatory activities	
Glycyrrhiza	1.90g	Antitussive effect	
Peony Root	1.90g	Immunity improvement	
Head	(*102	West 7.1	

#### Uses

■Bronchitis, Bronchial asthma, Coryza, Runny nose, Cough with dilute phlegm, rhinitis

### Warnings

### When using this product

- ■Do not use in patients with genetic problems such as galactose intolerance, Lapp lactase deficiency, or glucose-galactose malabsorption since this drug contains lactose.
- ■Do not use it for more than 1 month unless the symptoms improve. Long-term use should only be done under medical supervision.
- ■When used in combination with potassium-containing agents, licorice-containing drugs, and diuretics (e.g., furosemide, ethacrynic acid, trichlormethiazide), myopathy is likely to occur, so take it carefully.
- ■When combined with MaHuang or ephedrine-containing agents, MAO inhibitors, thyroid agents(thyroxine, liothyronine), catecholamine agents(epinephrine, isoprenaline), and xanthin agents(theophylline, dipropylene), symptoms such as insomnia, excessive sweat, rapid pulse, frequent urination, palpitations, general weakness, and mental excitement are likely to appear, so take them carefully.

Ask a doctor before use if you have high blood pressure, edema, heart or kidney problems, rash, redness, diabetic, another disease. Stop use and ask a doctor if the following serious side effects occur. These may be signs of a serious condition.

- ■Gastric aldosteronism (decreased urine, swollen face, hands, feet, eyelids, etc.)
- Myopathy (muscle weakness, cramps, paralysis)
- ■Interstitial pneumonia (fever, cough, shortness of breath, lung sound abnormalities)
- ■Liver disorders (elevated liver enzymes, jaundice)
- ■Skin reactions (rash, hives)

■ Digestive issues (loss of appetite, stomach discomfort)
If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children, In case of overdose, get medical help or contact a poison control center right away

## Directions

■ Take a packet of medicine (one dose) of an adult three times a day before or between meals.

#### Other Information

■ Store at room temperature, 1–30°C (34–86°F)

#### Inactive Ingredient

corn starch, Lactose Hydrate

### Questions or comments ?

Call weekdays from 9 a.m to 5 p.m EST at (201) 669-8405

# HAN SHIN GAMKONAL EXTRACT GRANULES

dry ginger, glycyrrhiza, peony root granule

Dro	duct	Inform	ation
PIU	uuct		alivii

Product Type HUMAN OTC DRUG Item Code (Source) NDC:73466-0033

**Route of Administration** ORAL

Active Ingredient/Active Moiety				
Ingredient Name	Basis of Strength	Strength		
PAEONIA LACTIFLORA ROOT (UNII: 3Z3866YW6P) (PAEONIA LACTIFLORA ROOT - UNII:3Z3866YW6P)	PAEONIA LACTIFLORA ROOT	1.9 g in 3.5 g		
GINGER (UNII: C5529G5JPQ) (GINGER - UNII:C5529G5JPQ)	GINGER	1.9 g in 3.5 g		
LICORICE (UNII: 61ZBX54883) (LICORICE - UNII:61ZBX54883)	LICORICE	1.9 g in 3.5 g		

Inactive Ingredients		
Ingredient Name	Strength	
LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)		

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:73466- 0033-1	120 in 1 PACKAGE	03/01/2025	
1		3.5 g in 1 POUCH; Type 0: Not a Combination Product		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		03/01/2025	

# **Labeler -** MISOCOS (695640173)

# Registrant - MISOCOS (695640173)

Establishment			
Name	Address	ID/FEI	Business Operations
MISOCOS		695640173	manufacture(73466-0033) , label(73466-0033)

Revised: 3/2025 MISOCOS