

UP AND UP ANTI DIARRHEAL- loperamide hcl suspension

Target Corporation

Target Corporation Anti-Diarrheal Drug Facts

Active ingredient (in each 7.5 mL)

Loperamide HCl 1 mg

Purpose

Anti-diarrheal

Use

controls symptoms of diarrhea, including Travelers' Diarrhea

Warnings

Allergy alert: Do not use if you have ever had a rash or other allergic reaction to loperamide HCl

Heart alert: Taking more than directed can cause serious heart problems or death

Do not use

if you have bloody or black stool

Ask a doctor before use if you have

- fever
- mucus in the stool
- a history of liver disease
- a history of abnormal heart rhythm

Ask a doctor or pharmacist before use if you are

taking a prescription drug. Loperamide may interact with certain prescription drugs.

When using this product

tiredness, drowsiness or dizziness may occur. Be careful when driving or operating machinery.

Stop use and ask a doctor if

- symptoms get worse
- diarrhea lasts for more than 2 days
- you get abdominal swelling or bulging. These may be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- **drink plenty of clear fluids to help prevent dehydration caused by diarrhea**
- find right dose on chart. If possible, use weight to dose; otherwise use age.
- shake well before using
- use only enclosed dosing cup specifically designed for use with this product. Do not use any other dosing device.
- mL = milliliter

adults and children 12 years and over	30 mL after the first loose stool; 15 mL after each subsequent loose stool; but no more than 60 mL in 24 hours
children 9-11 years (60-95 lbs)	15 mL after the first loose stool; 7.5 mL after each subsequent loose stool; but no more than 45 mL in 24 hours
children 6-8 years (48-59 lbs)	15 mL after the first loose stool; 7.5 mL after each subsequent loose stool; but no more than 30 mL in 24 hours
children under 6 years (up to 47 lbs)	ask a doctor
children under 2 years (up to 33 lbs)	do not use

Other information

- **each 30 mL contains:** sodium 15 mg
- store between 20-25°C (68-77°F)

Inactive ingredients

anhydrous citric acid, carboxymethylcellulose sodium, D&C yellow no. 10, FD&C blue no. 1, glycerin, microcrystalline cellulose, natural and artificial mint flavor, propylene glycol, purified water, simethicone, sodium benzoate, sucralose, titanium dioxide, xanthan gum

Questions or comments?

1-888-547-7400

Package/Label Principal Display Panel

see new warning and directions

Compare to active ingredient in Imodium® A-D

loperamide hydrochloride oral solution, 1 mg per 7.5 mL

anti-diarrheal

controls the symptoms of diarrhea

anti-diarrheal oral solution

MINT FLAVOR

8 FL OZ (240 mL)

Drug Facts

Active ingredient Purpose (in each 7.5 mL)
Loperamide HCl 1 mg.....Anti-diarrheal

Use controls symptoms of diarrhea, including Travelers' Diarrhea

Warnings

Allergy alert: Do not use if you have ever had a rash or other allergic reaction to loperamide HCl
Heart alert: Taking more than directed can cause serious heart problems or death

Do not use if you have bloody or black stool

Peel open to read complete directions and warnings before purchase

see new warning and directions

Compare to active ingredient in Imodium® A-D*

loperamide hydrochloride oral solution, 1 mg per 7.5 mL

anti-diarrheal

controls the symptoms of diarrhea
anti-diarrheal oral solution



8 FL OZ (240 mL)

NDC 11673-900-34


Do not use if printed plastic neckband is broken or missing.

*This product is not manufactured or distributed by Johnson & Johnson, owner of the registered trademark Imodium®.

100% satisfaction guaranteed or your money back.

GLUTEN FREE

245 05 0173 R02
IDC-000274-01-006
Dist. by Target Corp.
Mpls., MN 55403
©2019
Target Brands, Inc.
Shop Target.com



: 64534 UW F3

Drug Facts (continued)

Ask a doctor before use if you have
 ■ fever ■ mucus in the stool ■ a history of liver disease
 ■ a history of abnormal heart rhythm

Ask a doctor or pharmacist before use if you are taking a prescription drug. Loperamide may interact with certain prescription drugs.

When using this product tiredness, drowsiness or dizziness may occur. Be careful when driving or operating machinery.

Stop use and ask a doctor if ■ symptoms get worse
 ■ diarrhea lasts for more than 2 days
 ■ you get abdominal swelling or bulging.
 These may be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Drug Facts (continued)

Directions

- drink plenty of clear fluids to help prevent dehydration caused by diarrhea
- find right dose on chart. If possible, use weight to dose; otherwise use age.
- shake well before using
- use only enclosed dosing cup specifically designed for use with this product. Do not use any other dosing device.
- mL = milliliter

adults and children 12 years and over	30 mL after the first loose stool; 15 mL after each subsequent loose stool; but no more than 60 mL in 24 hours
---------------------------------------	---

children 9-11 years (60-95 lbs)	15 mL after the first loose stool; 7.5 mL after each subsequent loose stool; but no more than 45 mL in 24 hours
---------------------------------	--

Drug Facts (continued)

children 6-8 years (48-59 lbs)	15 mL after the first loose stool; 7.5 mL after each subsequent loose stool; but no more than 30 mL in 24 hours
--------------------------------	--

children 2-5 years (34 to 47 lbs)	ask a doctor
-----------------------------------	--------------

children under 2 years (up to 33 lbs)	do not use
---------------------------------------	------------

Other information

- each 30 mL contains: sodium 15 mg
- store between 20-25°C (68-77°F)

Drug Facts (continued)

Inactive ingredients anhydrous citric acid, carboxymethylcellulose sodium, D&C yellow no. 10, FD&C blue no. 1, glycerin, microcrystalline cellulose, natural and artificial mint flavor, propylene glycol, purified water, simethicone, sodium benzoate, sucralose, titanium dioxide, xanthan gum

Questions or comments?
1-888-547-7400

UP AND UP ANTI DIARRHEAL

loperamide hcl suspension

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:11673-900
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
LOPERAMIDE HYDROCHLORIDE (UNII: 77TI35393C) (LOPERAMIDE - UNII:6X9OC3H4II)	LOPERAMIDE HYDROCHLORIDE	1 mg in 7.5 mL

Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
CARBOXYMETHYLCELLULOSE SODIUM, UNSPECIFIED FORM (UNII: K679OBS311)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
GLYCERIN (UNII: PDC6A3C0OX)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
XANTHAN GUM (UNII: TTV12P4NEE)	
DIMETHICONE (UNII: 92RU3N3Y1O)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	

Product Characteristics

Color	GREEN (opaque, viscous)	Score	
Shape		Size	
Flavor	MINT	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:11673-900-34	240 mL in 1 BOTTLE; Type 0: Not a Combination Product	02/26/2016	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA091292	02/26/2016	

Labeler - Target Corporation (006961700)

Revised: 1/2020

Target Corporation