

**UP AND UP ANTI DIARRHEAL- loperamide hcl suspension**  
**Target Corporation**

-----  
**Target Corporation Anti-Diarrheal Drug Facts**

**Active ingredient (in each 7.5 mL)**

Loperamide HCl 1 mg

**Purpose**

Anti-diarrheal

**Use**

controls symptoms of diarrhea, including Travelers' Diarrhea

**Warnings**

**Allergy alert:** Do not use if you have ever had a rash or other allergic reaction to loperamide HCl

**Heart alert:** Taking more than directed can cause serious heart problems or death

**Do not use**

if you have bloody or black stool

**Ask a doctor before use if you have**

- fever
- mucus in the stool
- a history of liver disease
- a history of abnormal heart rhythm

**Ask a doctor or pharmacist before use if you are**

taking a prescription drug. Loperamide may interact with certain prescription drugs.

**When using this product**

tiredness, drowsiness or dizziness may occur. Be careful when driving or operating machinery.

**Stop use and ask a doctor if**

- symptoms get worse
- diarrhea lasts for more than 2 days
- you get abdominal swelling or bulging. These may be signs of a serious condition.

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

**Directions**

- **drink plenty of clear fluids to help prevent dehydration caused by diarrhea**
- find right dose on chart. If possible, use weight to dose; otherwise use age.
- shake well before using
- use only enclosed dosing cup specifically designed for use with this product. Do not use any other dosing device.
- mL = milliliter

adults and children 12 years and over	30 mL after the first loose stool; 15 mL after each subsequent loose stool; but no more than 60 mL in 24 hours
children 9-11 years (60-95 lbs)	15 mL after the first loose stool; 7.5 mL after each subsequent loose stool; but no more than 45 mL in 24 hours
children 6-8 years (48-59 lbs)	15 mL after the first loose stool; 7.5 mL after each subsequent loose stool; but no more than 30 mL in 24 hours
children under 6 years (up to 47 lbs)	ask a doctor
children under 2 years (up to 33 lbs)	do not use

**Other information**

- **each 30 mL contains:** sodium 15 mg
- store between 20-25°C (68-77°F)

**Inactive ingredients**

anhydrous citric acid, carboxymethylcellulose sodium, D&C yellow no. 10, FD&C blue no. 1, glycerin, microcrystalline cellulose, natural and artificial mint flavor, propylene glycol, purified water, simethicone, sodium benzoate, sucralose, titanium dioxide, xanthan gum

**Questions or comments?**

**1-888-547-7400**

**Package/Label Principal Display Panel**

Compare to active ingredient in Imodium<sup>®</sup> A-D

Loperamide Hydrochloride Oral Solution, 1 mg per 7.5 mL

Anti-Diarrheal

Controls the symptoms of diarrhea

Anti-Diarrheal Oral Solution

up&up<sup>™</sup>

8 FL OZ (240 mL)

Mint Flavor

## Drug Facts

**Active ingredient Purpose**  
(in each 7.5 mL)

Loperamide HCl 1 mg.....Anti-diarrheal

**Use** controls symptoms of diarrhea, including Travelers' Diarrhea

### Warnings

**Allergy alert:** Do not use if you have ever had a rash or other allergic reaction to loperamide HCl

**Heart alert:** Taking more than directed can cause serious heart problems or death

**Do not use** if you have bloody or black stool

Compare to active ingredient in Imodium® A-D\*

Loperamide Hydrochloride Oral Solution, 1 mg per 7.5 mL

# Anti-Diarrheal



up&up

Controls the symptoms of diarrhea

Anti-Diarrheal Oral Solution

8 FL OZ (240 mL)



Mint Flavor

**DO NOT USE IF PRINTED PLASTIC NECKBAND IS BROKEN OR MISSING.**

\*This product is not manufactured or distributed by Johnson & Johnson Consumer Inc., distributor of Imodium® A-D.

NDC 11673-900-34  
245 05 0173 R00 C-002262-01-073-0000

Dist. by Target Corp.  
Mpls., MN 55403  
TM & ©2024  
Target Brands, Inc.



EXP LOT

: 64534 UW F4

**ADHESIVE  
AREA  
NO COPY**

**NO  
VARNISH  
AREA**

### Drug Facts (continued)

children 6-8 years (48-59 lbs)	15 mL after the first loose stool; 7.5 mL after each subsequent loose stool; but no more than 30 mL in 24 hours
children 2-5 years (34 to 47 lbs)	ask a doctor
children under 2 years (up to 33 lbs)	do not use

### Other information

- each 30 mL contains: sodium 15 mg
- store between 20-25°C (68-77°F)

### Drug Facts (continued)

**Inactive ingredients** anhydrous citric acid, carboxymethylcellulose sodium, D&C yellow no. 10, FD&C blue no. 1, glycerin, microcrystalline cellulose, natural and artificial mint flavor, propylene glycol, purified water, simethicone, sodium benzoate, sucralose, titanium dioxide, xanthan gum

### Questions or comments?

1-888-547-7400

### Drug Facts (continued)

**Ask a doctor before use if you have**

- fever ■ mucus in the stool ■ a history of liver disease
- a history of abnormal heart rhythm

**Ask a doctor or pharmacist before use if you are** taking a prescription drug. Loperamide may interact with certain prescription drugs.

**When using this product** tiredness, drowsiness or dizziness may occur. Be careful when driving or operating machinery.

**Stop use and ask a doctor if** ■ symptoms get worse ■ diarrhea lasts for more than 2 days ■ you get abdominal swelling or bulging. These may be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

### Drug Facts (continued)

#### Directions

■ **drink plenty of clear fluids to help prevent dehydration caused by diarrhea**

■ find right dose on chart. If possible, use weight to dose; otherwise use age.

■ shake well before using

■ use only enclosed dosing cup specifically designed for use with this product. Do not use any other dosing device.

■ mL = milliliter

adults and children 12 years and over

30 mL after the first loose stool; 15 mL after each subsequent loose stool; but no more than 60 mL in 24 hours

children 9-11 years (60-95 lbs)

15 mL after the first loose stool; 7.5 mL after each subsequent loose stool; but no more than 45 mL in 24 hours

**ADHESIVE  
AREA  
NO COPY**

UP AND UP ANTI DIARRHEAL

loperamide hcl suspension

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:11673-900
Route of Administration	ORAL		

Active Ingredient/Active Moiety		
Ingredient Name	Basis of Strength	Strength
LOPERAMIDE HYDROCHLORIDE (UNII: 77TI35393C) (LOPERAMIDE - UNII:6X9OC3H4II)	LOPERAMIDE HYDROCHLORIDE	1 mg in 7.5 mL

Inactive Ingredients	
Ingredient Name	Strength
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
CARBOXYMETHYLCELLULOSE SODIUM, UNSPECIFIED (UNII: K679OBS311)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
GLYCERIN (UNII: PDC6A3C0OX)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
XANTHAN GUM (UNII: TTV12P4NEE)	
DIMETHICONE (UNII: 92RU3N3Y1O)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	

Product Characteristics			
Color	GREEN (opaque, viscous)	Score	
Shape		Size	
Flavor	MINT	Imprint Code	
Contains			

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:11673-900-34	240 mL in 1 BOTTLE; Type 0: Not a Combination Product	02/26/2016	

Marketing Information			
Marketing	Application Number or Monograph	Marketing Start	Marketing End

Category	Citation	Date	Date
ANDA	ANDA091292	02/26/2016	

**Labeler** - Target Corporation (006961700)

Revised: 4/2025

Target Corporation