

**KAOPECTATE- bismuth subsalicylate tablet, chewable**  
**Kramer Laboratories**

-----  
**Drug Facts**

<b>Active ingredient (per chew)</b>	<b>Purposes</b>
Bismuth subsalicylate 262 mg	Anti-diarrheal, Upset stomach reliever

**Uses** relieves:

- traveler's diarrhea
- diarrhea
- upset stomach due to overindulgence in food and drink, including:
  - heartburn
  - indigestion
  - nausea
  - gas
  - belching
  - fullness

**Warnings**

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy Alert: Contains salicylate.**

**Do not take if you are**

- allergic to salicylates (including aspirin)
- taking other salicylate products

**Do not use if you have**

- an ulcer
- a bleeding problem
- bloody or black stool

**Ask a doctor before use if you have**

- fever

- mucus in the stool

**Ask a doctor or pharmacist before use if you are taking any drug for**

- anticoagulation (thinning the blood)
- diabetes
- gout
- arthritis

**When using this product** a temporary, but harmless, darkening of the stool and/or tongue may occur

**Stop use and ask a doctor if**

- symptoms get worse
- ringing in the ears or loss of hearing occurs
- diarrhea lasts more than 2 days

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

***Directions***

- **adults and children 12 years of age and older:** 2 chews (1 dose)
- repeat dose every 1/2 hour to 1 hour as needed
- do not exceed 8 doses in 24 hours
- use until diarrhea stops but not more than 2 days
- **children under 12 years:** ask a doctor
- drink plenty of clear fluids to help prevent dehydration caused by diarrhea

***Other information***

- each soft chew contains: **total salicylates 99 mg**
- **do not use if carton was opened or if printed foil inner seal under bottle cap is torn or missing**
- store at room temperature 20°-25° C (68°-77° F)

***Inactive ingredients***

calcium carbonate, confectioner's sugar, corn syrup, corn syrup solids, FD&C red #40 aluminum lake, glycerin, hydrogenated coconut oil, natural masking flavor, soy lecithin, strawberry flavor, sucrose, water

**PRINCIPAL DISPLAY PANEL**

***GREAT TASTE!***

***FAST & EFFECTIVE RELIEF***

## **Kaopectate®**

*Bismuth Subsalicylate • Anti-Diarrheal • Upset Stomach Reliever*

### ***Soft Chews***

#### **Diarrhea & Upset Stomach**

- ✓ Begins controlling symptoms from the first dose
- ✓ Quickly relieves urgency, gas, and cramping
- ✓ Effective on diarrhea from bacteria, viruses, and other causes

#### **24 Soft Chews**

Individually wrapped

#### ***Strawberry Flavor***



**FAST & EFFECTIVE RELIEF**

**Do not take maximum dose for more than 2 days per week.**

**Kramer Laboratories, Inc.**  
Bridgewater, NJ 08807

1-800-824-4894

502851

# FAST & EFFECTIVE RELIEF

ZIPPER AREA

SIDE SEAL

SIDE SEAL

## Drug Facts

**Active ingredient (per chew)**    **Purposes**  
Bismuth subsalicylate 262 mg.....Anti-diarrheal,  
Upset stomach reliever

**Uses** relieve: • traveler's diarrhea • diarrhea • upset stomach due to overindulgence in food and drink, including: • heartburn • indigestion • nausea • gas • belching • fullness

### Warnings

**Reye's syndrome:** Children and teenagers who have or are recovering from chickenpox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy Alert:** Contains salicylate. Do not take if you are • allergic to salicylates (including aspirin) • taking other salicylate products

Do not use if you have • an ulcer • a bleeding problem • bloody or black stool

Ask a doctor before use if you have • fever • mucus in the stool

Ask a doctor or pharmacist before use if you have or are taking any drug for • anticoagulation (thinning the blood) • diabetes • gout • arthritis

When using this product a temporary, but harmless, darkening of the stool and/or tongue may occur

Stop use and ask a doctor if • symptoms get worse • ringing in the ears or loss of hearing occurs • diarrhea lasts more than 2 days

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. ▶

## Drug Facts (continued)

### Directions

- adults and children 12 years of age and older: 2 chews (1 dose)
- repeat dose every 1/2 hour to 1 hour as needed
- do not exceed 8 doses in 24 hours
- use until diarrhea stops but not more than 2 days
- children under 12 years: ask a doctor
- drink plenty of clear fluids to help prevent dehydration caused by diarrhea

### Other information

- each soft chew contains: total salicylates 99 mg
- do not use if package is opened or damaged
- store at room temperature 20°-25°C (68°-77°F)

### Inactive ingredients

calcium carbonate, confectioners sugar, corn syrup, corn syrup solids, F D & C red #40 aluminum lake, glycerin, hydrogenated coconut oil, natural masking flavor, soy lecithin, strawberry flavor, sucrose, water

Do not take maximum dose for more than 2 days per week.

Kramer Laboratories, Inc.  
Bridgewater, NJ 08807  
1-800-824-4894



502851

# KAOPECTATE

bismuth subsalicylate tablet, chewable

## Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:55505-206
<b>Route of Administration</b>	ORAL		

## Active Ingredient/Active Moiety

<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
<b>Bismuth Subsalyclate</b> (UNII: 62TEY51RR1) (Bismuth Cation - UNII:Z59CD1I8YE)	Bismuth Subsalyclate	262 mg

## Inactive Ingredients

<b>Ingredient Name</b>	<b>Strength</b>
<b>Calcium Carbonate</b> (UNII: H0G9379FGK)	
<b>Sucrose</b> (UNII: C151H8M554)	
<b>Corn Syrup</b> (UNII: 9G5L16BK6N)	
<b>Fd&amp;C Red No. 40</b> (UNII: WZB9127XOA)	
<b>Glycerin</b> (UNII: PDC6A3C00X)	
<b>Hydrogenated Coconut Oil</b> (UNII: JY81OXM1OM)	
<b>Lecithin, Soybean</b> (UNII: 1DI56QDM62)	
<b>Water</b> (UNII: 059QF0KO0R)	

## Product Characteristics

<b>Color</b>	PINK	<b>Score</b>	no score
<b>Shape</b>	RECTANGLE	<b>Size</b>	23mm
<b>Flavor</b>	STRAWBERRY	<b>Imprint Code</b>	K
<b>Contains</b>			

## Packaging

<b>#</b>	<b>Item Code</b>	<b>Package Description</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
1	NDC:55505-206-69	1 in 1 CARTON	02/01/2021	
1		24 in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:55505-206-79	2 in 1 PACKAGE	06/01/2025	
2		1 in 1 CARTON		
2		24 in 1 BOTTLE; Type 0: Not a Combination Product		

## Marketing Information

<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
OTC Monograph Drug	M008	02/01/2021	

**Labeler** - Kramer Laboratories (122720675)

Revised: 12/2025

Kramer Laboratories