

CHERRY SORE THROAT LOZENGES- benzocaine, menthol lozenge
Meijer Distribution Inc.

Menthol 3.6mg
Benzocaine 15mg

Active Ingredients (in each lozenge)

Benzocaine 15mg

Menthol 3.6mg

Purpose

Oral anesthetic

Oral anesthetic

Uses - temporarily relieves these symptoms

occasional minor irritation, pain, sore mouth and sore throat

Warnings

Methemoglobinemia warning:

Use of this product may cause methemoglobinemia, a serious condition that must be treated promptly because it reduces the amount of oxygen carried in blood. This can occur even if you have used this product before. Stop use and seek immediate medical attention if you or a child in your care develops:

- pale, gray, or blue colored skin (cyanosis)
- headache
- rapid heart rate
- shortness of breath
- dizziness or lightheadedness
- fatigue or lack of energy

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, swelling, nausea, or vomiting, consult a doctor promptly. If sore mouth symptoms do not improve in 7 days, or if irritation, pain, or redness persists or worsens, see your dentist or doctor promptly.

Allergy alert: Do not use this product if you have a history of allergy to local anesthetics such as procaine, butacaine, benzocaine, or other "caine" anesthetics.

Do not use

- for teething
- in children under 2 years of age

Stop use and ask a doctor if

- sore throat is severe or irritation, pain or redness persists or worsens
- sore mouth symptoms do not improve in 7 days

If Pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

adults and children 5 years and older - allow 1 lozenge to dissolve slowly in mouth. May be repeated every two hours as necessary or as directed by a doctor

children under 5 years - do not use

Other information

- store at room temperature
- protect from moisture

Inactive Ingredients

Corn starch, FD&C red #40, glucose syrup, glycerin, medium chain triglycerides, natural and artificial flavors, propylene glycol, soybean oil, sucrose, and water.

Questions? Call **1-800-245-2898**, Monday through Friday, 9AM - 4PM EST



Drug Facts

Active ingredients (in each lozenge)

Benzocaine 15 mg.....	Oral anesthetic
Menthol 3.6 mg.....	Oral anesthetic

Purpose

Uses temporarily relieves these symptoms:
 ■ occasional minor irritation, pain, sore mouth and sore throat

Warnings

Methemoglobinemia warning:

Use of this product may cause methemoglobinemia, a serious condition that must be treated promptly because it reduces the amount of oxygen carried in blood. This can occur even if you have used this product before. Stop use and seek immediate medical attention if you or a child in your care develops: ■ pale, gray, or blue colored skin (cyanosis)
 ■ headache ■ rapid heart rate ■ shortness of breath ■ dizziness or lightheadedness ■ fatigue or lack of energy

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, swelling, nausea, or vomiting, consult a doctor promptly. If sore mouth symptoms do not improve in 7 days, or if irritation, pain, or redness persist or worsens, see your dentist or doctor promptly.

Allergy alert: Do not use this product if you have a history of allergy to local anesthetics such as procaine, butacaine, benzocaine, or other "caine" anesthetics.

Do not use

■ for teething ■ in children under 2 years of age

Stop use and ask doctor if

■ sore throat is severe or irritation, pain or redness persists or worsens ■ sore mouth symptoms do not improve in 7 days

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact Poison Control Center right away.

Directions

adults and children 5 years and older	allow 1 lozenge to dissolve slowly in mouth. May be repeated every two hours as necessary or as directed by a doctor
children under 5 years	do not use

Other information ■ store at room temperature ■ protect from moisture

Drug Facts (continued)

Inactive ingredients Artificial flavor, corn starch, corn syrup, FD&C Red #40, glycerin, medium chain triglycerides, propylene glycol, soybean oil, sucrose and water.

Questions? Call 1-800-245-2898, Monday through Friday, 9AM - 4PM EST

*This product is not manufactured or distributed by Reckitt Benckiser owner of the registered trademark Cepacol®.
 WARNING: TAMPER-EVIDENT PACKAGING. DO NOT USE IF OUTER BOX OR BLISTER IS OPEN OR DAMAGED.

Gluten-Free
 DIST. BY MEYER DISTRIBUTION, INC.
 GRAND RAPIDS, MI 49544
 www.meyler.com



PN-60002200
 NDC-41250-069-18

Actual Size

 LOZENGES
 18

Fast Acting

COMPARE TO
 CEPACOL®
 ACTIVE INGREDIENT™



CHERRY

Menthol 3.6 mg | Benzocaine 15 mg
 Oral Anesthetic

FAST, SOOTHING, TEMPORARY RELIEF OF
 MINOR SORE THROAT PAIN AND IRRITATION

meijer
**Sore Throat
 Lozenges**

meijer
**Sore Throat
 Lozenges**



CHERRY SORE THROAT LOZENGES

benzocaine, menthol lozenge

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:41250-068
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BENZOCAINE (UNII: U3RSY48JW5) (BENZOCAINE - UNII:U3RSY48JW5)	BENZOCAINE	15 mg
MENTHOL (UNII: L7T10EIP3A) (MENTHOL - UNII:L7T10EIP3A)	MENTHOL	3.6 mg

Inactive Ingredients

Ingredient Name	Strength
SUCROSE (UNII: C151H8M554)	

Product Characteristics

Color	red	Score	score with uneven pieces
Shape	ROUND	Size	17mm
Flavor	CHERRY	Imprint Code	B
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:41250-068-18	1 in 1 CARTON	04/01/2016	
1		18 in 1 BLISTER PACK; Type 0: Not a Combination Product		



Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	04/01/2016	

Labeler - Meijer Distribution Inc. (006959555)

Revised: 2/2026

Meijer Distribution Inc.