

ASPIRIN LOW DOSE- aspirin tablet, delayed release
L.N.K. International, Inc.

Quality Plus 44-600A

Active ingredient (in each tablet)

Aspirin 81 mg (NSAID)*

*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever

Uses

for the temporary relief of minor aches and pains or as recommended by your doctor. **Because of its delayed action, this product will not provide fast relief of headaches or other symptoms needing immediate relief.**

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction, which may include:

- hives
- facial swelling
- shock
- asthma (wheezing)

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- take more or for a longer time than directed
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- have 3 or more alcoholic drinks every day while using this product
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]

Do not use

- if you are allergic to aspirin or any other pain reliever/fever reducer
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you have asthma
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

taking a prescription drug for

- gout
- diabetes
- arthritis

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
 - vomit blood
 - have bloody or black stools
 - feel faint
 - have stomach pain that does not get better
- ringing in the ears or a loss of hearing occurs
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur

These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- **do not take more than directed**
- drink a full glass of water with each dose
- adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor
- children under 12 years: do not use unless directed by a doctor

Other information

- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- see end flap for expiration date and lot number

Inactive ingredients

corn starch, D&C yellow #10, FD&C yellow #6, hypromellose, methacrylic acid, microcrystalline cellulose, polydextrose, polyethylene glycol, shellac wax, silica, simethicone, sodium bicarbonate, sodium lauryl sulfate, talc, titanium dioxide, triacetin, triethyl citrate

Questions or comments?

1-800-426-9391

Principal display panel

**Quality
+Plus**

NDC 50844-600-32

†Compare to active ingredient in
Bayer® Low Dose Aspirin

**LOW DOSE
ASPIRIN 81 mg
PAIN RELIEVER
(NSAID)**

- **ASPIRIN REGIMEN**
- **SAFETY COATED**

**120
ENTERIC Coated Tablets**

ACTUAL SIZE

Talk to your doctor or other healthcare provider
before using this product for your heart.

**TAMPER EVIDENT: DO NOT USE IF
IMPRINTED SAFETY SEAL UNDER
CAP IS BROKEN OR MISSING**

†This product is not manufactured or distributed by
Bayer AG, owner of the registered trademark Bayer®
Low Dose Aspirin.

50844 REV0122D60032

Distributed by **LNK INTERNATIONAL, INC.**
60 Arkay Drive, Hauppauge, NY 11788
USA

Lot no. & Exp. Date

0150844 56332 2

QUALITY PLUS

LOW DOSE

ASPIRIN 81 mg

PAIN RELIEVER (NSAID)

120 ENTERIC Coated Tablets

NDC 50844-600-32

*Compare to active ingredient in Bayer® Low Dose Aspirin

ASPIRIN REGIMEN

SAFETY COATED

ACTUAL SIZE

Talk to your doctor or other healthcare provider before using this product for your heart.

Drug Facts (continued)

polydextrose, polyethylene glycol, shellac wax, silica, simethicone, sodium bicarbonate, sodium lauryl sulfate, talc, titanium dioxide, triacetin, triethyl citrate

Questions or comments?

1-800-426-9391

*This product is not manufactured or distributed by Bayer AG, owner of the registered trademark Bayer® Low Dose Aspirin. 50844 REV0122D60032

Distributed by LNK INTERNATIONAL, INC.

60 Arkey Drive, Hauppauge, NY 11788 USA

Drug Facts (continued)

■ pain gets worse or lasts more than 10 days

■ fever gets worse or lasts more than 3 days

■ redness or swelling is present

■ new symptoms occur

These could be signs of a serious condition.

If **pregnant or breast-feeding**, ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

■ do not take more than directed

■ drink a full glass of water with each dose

■ adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor

■ children under 12 years: do not use unless directed by a doctor

Other information

■ store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)

■ see end flap for expiration date and lot number

Inactive ingredients corn starch, D&C yellow #10, FD&C yellow #6, hypromellose, methacrylic acid, microcrystalline cellulose,

Drug Facts

Active ingredient (in each tablet)

Aspirin 81 mg (NSAID)*, nonsteroidal anti-inflammatory drug

Uses for the temporary relief of minor aches and pains or as recommended by your doctor. Because of its delayed action, this product will not provide fast relief of headaches or other symptoms needing immediate relief.

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction, which may include:

- hives
- shock
- asthma (wheezing)
- facial swelling

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you take more or for a longer time than directed

- take a blood thinning (anticoagulant) or steroid drug
- are age 60 or older

Drug Facts (continued)

■ you experience any of the following signs of stomach bleeding: ■ vomit blood ■ feel faint ■ have bloody or black stools ■ have stomach pain that does not get better ■ ringing in the ears or a loss of hearing occurs

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.

Ask a doctor or pharmacist before use if you are taking a prescription drug for

- gout
- diabetes
- arthritis

Drug Facts

Active ingredient (in each tablet)

Aspirin 81 mg (NSAID)*, nonsteroidal anti-inflammatory drug

Uses for the temporary relief of minor aches and pains or as recommended by your doctor. Because of its delayed action, this product will not provide fast relief of headaches or other symptoms needing immediate relief.

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction, which may include:

- hives
- shock
- asthma (wheezing)
- facial swelling

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you take more or for a longer time than directed

- take a blood thinning (anticoagulant) or steroid drug
- are age 60 or older

Drug Facts (continued)

■ have had stomach ulcers or bleeding problems

■ have 3 or more alcoholic drinks every day while using this product

■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)

Do not use

- if you are allergic to aspirin or any other pain reliever/fever reducer
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you have asthma
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are taking a prescription drug for

- gout
- diabetes
- arthritis

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.

Ask a doctor or pharmacist before use if you experience any of the following signs of stomach bleeding:

- vomit blood
- feel faint
- have bloody or black stools
- have stomach pain that does not get better
- ringing in the ears or a loss of hearing occurs

Drug Facts (continued)

■ you experience any of the following signs of stomach bleeding: ■ vomit blood ■ feel faint ■ have bloody or black stools ■ have stomach pain that does not get better ■ ringing in the ears or a loss of hearing occurs

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.

Ask a doctor or pharmacist before use if you are taking a prescription drug for

- gout
- diabetes
- arthritis

Quality Plus 44-600A

| | | | |
|--|----------------|---------------------------|---------------|
| ASPIRIN LOW DOSE | | | |
| aspirin tablet, delayed release | | | |
| Product Information | | | |
| Product Type | HUMAN OTC DRUG | Item Code (Source) | NDC:50844-600 |
| Route of Administration | ORAL | | |
| Active Ingredient/Active Moiety | | | |

| Ingredient Name | Basis of Strength | Strength |
|---|-------------------|----------|
| ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E) | ASPIRIN | 81 mg |

Inactive Ingredients

| Ingredient Name | Strength |
|--|----------|
| STARCH, CORN (UNII: O8232NY3SJ) | |
| D&C YELLOW NO. 10 (UNII: 35SW5USQ3G) | |
| FD&C YELLOW NO. 6 (UNII: H77VEI93A8) | |
| HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO) | |
| METHACRYLIC ACID (UNII: 1CS02G8656) | |
| MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U) | |
| POLYDEXTROSE (UNII: VH2XOU12IE) | |
| POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A) | |
| SHELLAC (UNII: 46N107B71O) | |
| SILICON DIOXIDE (UNII: ETJ7Z6XBU4) | |
| DIMETHICONE (UNII: 92RU3N3Y1O) | |
| WATER (UNII: 059QF0KO0R) | |
| SODIUM BICARBONATE (UNII: 8MDF5V39QO) | |
| SODIUM LAURYL SULFATE (UNII: 368GB5141J) | |
| TALC (UNII: 7SEV7J4R1U) | |
| TITANIUM DIOXIDE (UNII: 15FIX9V2JP) | |
| TRIACETIN (UNII: XHX3C3X673) | |
| TRIETHYL CITRATE (UNII: 8Z96QXD6UM) | |

Product Characteristics

| | | | |
|-----------------|--------|---------------------|----------|
| Color | yellow | Score | no score |
| Shape | ROUND | Size | 6mm |
| Flavor | | Imprint Code | L |
| Contains | | | |

Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|---|------------------|---|----------------------|--------------------|
| 1 | NDC:50844-600-32 | 1 in 1 BOX | 05/01/2011 | |
| 1 | | 120 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product | | |
| 2 | NDC:50844-600-95 | 300 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product | 05/01/2011 | |
| 3 | NDC:50844-600-14 | 500 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product | 05/01/2011 | |
| 4 | NDC:50844-600-01 | 1 in 1 BOX | 05/01/2011 | 12/18/2020 |
| 4 | | 30 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product | | |
| 5 | NDC:50844-600-17 | 1 in 1 BOX | 05/01/2011 | 12/18/2020 |
| 5 | | 300 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product | | |

Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|--------------------|--|----------------------|--------------------|
| OTC Monograph Drug | M013 | 05/01/2011 | |

Labeler - L.N.K. International, Inc. (038154464)

Establishment

| Name | Address | ID/FEI | Business Operations |
|-------------------------|---------|-----------|------------------------|
| LNK International, Inc. | | 832867837 | manufacture(50844-600) |

Establishment

| Name | Address | ID/FEI | Business Operations |
|-------------------------|---------|-----------|------------------------|
| LNK International, Inc. | | 832867894 | manufacture(50844-600) |

Establishment

| Name | Address | ID/FEI | Business Operations |
|-------------------------|---------|-----------|--|
| LNK International, Inc. | | 868734088 | manufacture(50844-600) , pack(50844-600) |

Establishment

| Name | Address | ID/FEI | Business Operations |
|-------------------------|---------|-----------|------------------------|
| LNK International, Inc. | | 117025878 | manufacture(50844-600) |

Revised: 6/2025

L.N.K. International, Inc.