

TOPCARE ANTACID PLUS ANTI GAS- aluminum hydroxide, magnesium hydroxide, simethicone suspension

Topco Associates LLC

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Topco Associates LLC. Antacid Plus Anti-Gas Drug Facts

Active ingredients (in each 10 mL)

Aluminum hydroxide (equiv. to dried gel, USP) 800 mg

Magnesium hydroxide 800 mg

Simethicone 80 mg

Purposes

Antacid

Antigas

Uses

relieves

- heartburn
- acid indigestion
- sour stomach
- upset stomach due to these symptoms
- pressure and bloating commonly referred to as gas

Warnings

Ask a doctor before use if you have

- kidney disease
- a magnesium-restricted diet

Ask a doctor or pharmacist before use if you are

presently taking a prescription drug. Antacids may interact with certain prescription drugs.

When using this product

do not exceed 60 mL in a 24-hour period, or use the maximum dosage for more than 2 weeks

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- shake well before use
- measure with dosing cup provided
- adults and children 12 years and over: 10 mL – 20 mL between meals, at bedtime or as directed by a doctor
- do not take more than 60 mL in any 24 hour period
- do not use the maximum dosage for more than 2 weeks
- children under 12 years: ask a doctor
- mL = milliliterr

Other information

- each 10 mL contains: magnesium 350 mg, potassium 20 mg and sodium 3 mg
- does not meet USP requirements for preservative effectiveness
- store at 20-25°C (68-77°F), do not freeze

Inactive ingredients

butylparaben, flavor, hypromellose, microcrystalline cellulose and carboxymethylcellulose sodium, potassium citrate, propylparaben, purified water, simethicone emulsion, sorbitol, sorbitol solution

Questions or comments?

1-888-423-0139

Principal Display Panel

COMPARE TO MAXIMUM STRENGTH MYLANTA® ACTIVE INGREDIENTS

MAXIMUM STRENGTH

Antacid

ANTACID / ANTI-GAS

FAST ACTING

OUR PHARMACISTS RECOMMEND

Tough on Heartburn

Works on Contact

CLASSIC ORIGINAL FLAVOR

12 FL OZ (355 mL)



TOPCARE ANTACID PLUS ANTI GAS

aluminum hydroxide, magnesium hydroxide, simethicone suspension

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:36800-340
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ALUMINUM HYDRO XIDE (UNII: 5QB0T2IUN0) (ALUMINUM HYDRO XIDE - UNII:5QB0T2IUN0)	ALUMINUM HYDRO XIDE	800 mg in 10 mL
MAGNESIUM HYDRO XIDE (UNII: NBZ3QY004S) (HYDRO XIDE ION - UNII:9159UV381P, MAGNESIUM CATION - UNII:T6V3LHY838)	MAGNESIUM HYDRO XIDE	800 mg in 10 mL
DIMETHICONE (UNII: 92RU3N3Y1O) (DIMETHICONE - UNII:92RU3N3Y1O)	DIMETHICONE	80 mg in 10 mL

Inactive Ingredients

Ingredient Name	Strength
SILICON DIO XIDE (UNII: ETJ7Z6XBU4)	
BUTYLPARABEN (UNII: 3QPI1U3FV8)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
CARBOXYMETHYLCELLULOSE SODIUM (UNII: K679OBS311)	
POTASSIUM CITRATE (UNII: EE90ONI6FF)	
PROPYLPARABEN (UNII: Z8IX2SC1OH)	
WATER (UNII: 059QF0K00R)	
SORBITOL (UNII: 506T60A25R)	

Product Characteristics

Color	WHITE (opaque)	Score	
Shape		Size	
Flavor	LEMON	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:36800-340-40	355 mL in 1 BOTTLE; Type 0: Not a Combination Product	07/15/1990	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part331	07/15/1990	

Labeler - Topco Associates LLC (006935977)

Revised: 12/2019

Topco Associates LLC