

**DAYTIME COLD AND FLU- acetaminophen 325mg/ dextromethorphan hbr 10mg/ phenylephrine hcl 5mg capsule**  
**Grocery Outlet**

-----

**Daytime Cold and Flu Relief**

**Active ingredient(s)**

Acetaminophen 325mg

Dextromethorphan HBr 10mg

Phenylephrine HCl 5mg

**Purpose**

Pain reliever/fever reducer

Cough suppressant

Nasal decongestant

**Use(s)**

temporarily relieves common cold and flu symptoms: ■ nasal congestion ■ cough due to minor throat and bronchial irritation ■ sore throat ■ headache ■ minor aches and pains ■ fever

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4 doses in 24 hours, which is the maximum daily amount for this product
- with other drugs containing acetaminophen
- or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, lasts for more than 2 days, occurs with or followed by fever, headache, rash, nausea, or vomiting, see a doctor promptly.

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If

you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drugs. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

### **Ask a doctor before use if you have**

■ liver disease ■ heart disease ■ high blood pressure ■ thyroid disease ■ diabetes ■ trouble urinating due to an enlarged prostate gland ■ persistent or chronic cough such as occurs with smoking, asthma, or emphysema ■ cough that occurs with too much phlegm (mucus)

### **Ask a doctor or pharmacist before use if**

taking the blood thinning drug warfarin.

### **When using this product**

do not use more than directed

### **Stop use and ask a doctor before use if**

- you get nervous, dizzy or sleepless
- symptoms get worse or last more than 5 days (children) or 7 days (adults)
- fever gets worse or lasts more than 3 days and redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with rash or headache that lasts. These could be signs of a serious condition.

### **If pregnant or breastfeeding,**

ask a health professional before use.

### **Keep out of reach of children**

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

### **Directions**

take only as directed

- do not exceed 4 doses per 24 hours
- **Adults and children 12 years and over:** 2 softgels with water every 4 hours
- **Children 4 to under 12 years:** ask a doctor
- **Children under 4 years:** do not use

### **Other information**

- store at room temperature 20-25 °C (68-77°F)

- do not use if blister unit is torn or open

**Inactive ingredients**

FD&C red #40, FD&C yellow #6, gelatin, glycerin, polyethylene glycol, povidone, propylene glycol, purified water, sorbitol sorbitan solution, white edible ink

**Questions/Comments**

Call **1-877-290-4008**

**Principal Display Panel**

**Drug Facts**

**Active Ingredients (in each softgel) Purposes**

Dextromethorphan HBr 10mg.....Cough suppressant  
Phenylephrine HCl 5mg.....Nasal decongestant

**Uses** temporarily relieves common cold and flu symptoms:  
in nasal congestion in cough due to minor throat and bronchial  
irritation in sore throat in headache in minor aches and pains  
in fever

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe  
liver damage may occur if you take in more than 4 doses in 24  
hours, which is the maximum daily amount for this product  
in with other drugs containing acetaminophen in 3 or more  
alcoholic drinks every day while using this product.

**Allergy alert:** Acetaminophen may cause severe skin  
reactions.

Symptoms may include: in skin redness/itching in blisters in rash  
if a skin reaction occurs, stop use and seek medical help right  
away.

**Sore throat warning:** If sore throat is severe, lasts for more than  
2 days, occurs with or followed by fever, headache, rash, nausea,  
or vomiting, see a doctor promptly.

**Do not use** in with any other drug containing acetaminophen  
(prescription or nonprescription). If you are not sure whether a  
drug contains acetaminophen, ask a doctor or pharmacist in if  
you are now taking a prescription monoamine oxidase inhibitor  
(MAOI) (certain drugs for depression, psychiatric or emotional  
conditions, or Parkinson's disease), or for 2 weeks after stopping  
the MAOI drugs. If you do not know if your prescription drug  
contains an MAOI, ask a doctor or pharmacist before taking this  
product.

**Ask a doctor before use** if you have in liver disease in heart  
disease in thyroid disease in diabetes in high blood pressure  
in trouble urinating due to an enlarged prostate gland in cough

**Drug Facts (continued)**

that occurs with too much phlegm (mucus) in persistent or  
chronic cough as occurs with smoking, asthma, or emphysema.

**Ask a doctor or pharmacist before use** if you are taking the  
blood thinning drug warfarin.

**When using this product do not use more than directed.**

**Stop use and ask a doctor if** in you get nervous, dizzy or  
sleepless in symptoms get worse or last more than 5 days  
(children) or 7 days (adults) in fever gets worse or lasts more  
than 3 days in redness or swelling is present in new symptoms  
occur in cough comes back, or occurs with rash or headache that  
lasts. These could be signs of a serious condition.

**If pregnant or breast-feeding, ask a health professional before  
use. Keep out of reach of children. Taking more than directed  
can cause serious health problems. In case of overdose, get  
medical help or contact a Poison Control Center**

(1-800-222-1222) right away. Quick medical attention is critical  
for adults as well as for children, even if you do not notice any  
signs or symptoms.

**Directions** in take only as directed in do not exceed 4 doses  
per 24 hours

**Adults and children 12 years and over** 2 softgels with water  
every 4 hours

**Children 4 to under 12 years** ask a doctor

**Children under 4 years** do not use

**Other information** in store at room temperature 20-25 °C  
(68-77° F)

**Inactive ingredients** FD&C red #40, FD&C yellow #6,  
gelatin, glycerin, polyethylene glycol, povidone, propylene glycol,  
purified water, sorbitol sorbitan solution, white edible ink

**Questions or comments?** Call 1-877-280-4008

**GO** Daytime Cold and Flu Relief

**GO** Relief of:  
• Aches & Pains  
• Sore Throat  
• Nasal Congestion  
• Fever  
• Headache  
• Cough

ACETAMINOPHEN, 325 mg (Pain Reliever / Fever Reducer)  
DEXTROMETHORPHAN HBr, 10 mg (Cough Suppressant)  
PHENYLEPHRINE HCl, 5 mg (Nasal Decongestant)

**Daytime Cold and Flu Relief**

Compare to active ingredients in Vicks® Dayquil™ Cold & Flu Relief Liquorice™

**GO** Daytime Cold and Flu Relief

ACETAMINOPHEN, 325 mg (Pain Reliever / Fever Reducer)  
DEXTROMETHORPHAN HBr, 10 mg (Cough Suppressant)  
PHENYLEPHRINE HCl, 5 mg (Nasal Decongestant)

**GO**

**Blister**

RETAIN CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION  
DO NOT USE IF BLISTER UNIT IS TORN OR OPEN

6 10159 0560 2

**DAYTIME COLD AND FLU**

acetaminophen 325mg/ dextromethorphan hbr 10mg/ phenylephrine hcl 5mg capsule

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:85828-417
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg
<b>PHENYLEPHRINE HYDROCHLORIDE</b> (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg

### Inactive Ingredients

Ingredient Name	Strength
<b>FD&amp;C RED NO. 40</b> (UNII: WZB9127XOA)	
<b>FD&amp;C YELLOW NO. 6</b> (UNII: H77VEI93A8)	
<b>GELATIN</b> (UNII: 2G86QN327L)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>POVIDONE</b> (UNII: FZ989GH94E)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>SORBITOL SOLUTION</b> (UNII: 8KW3E207O2)	

### Product Characteristics

<b>Color</b>	orange	<b>Score</b>	no score
<b>Shape</b>	CAPSULE	<b>Size</b>	21mm
<b>Flavor</b>		<b>Imprint Code</b>	PC9
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:85828-417-93	2 in 1 CARTON	03/25/2026	
1		12 in 1 BLISTER PACK; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	03/25/2026	

**Labeler** - Grocery Outlet (029161585)

Revised: 3/2026

Grocery Outlet