#### CAREONE ACID RELIEF- famotidine tablet, film coated Praxis, LLC

-----

#### American Sales Company Acid Relief Drug Facts

#### Active ingredient (in each tablet)

Famotidine 20 mg

#### Purpose

Acid reducer

#### Uses

- relieves heartburn associated with acid indigestion and sour stomach
- prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain food and beverages

#### Warnings

Allergy alert: Do not use if you are allergic to famotidine or other acid reducers

#### Do not use

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
- with other acid reducers

#### Ask a doctor before use if you have

- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with lightheadedness, sweating, or dizziness
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent chest pain
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain
- kidney disease

# Ask a doctor or pharmacist before use if you are

taking a prescription drug. Acid reducers may interact with certain prescription drugs.

#### Stop use and ask a doctor if

- your heartburn continues or worsens
- you need to take this product for more than 14 days

# If pregnant or breast-feeding,

ask a health professional before use.

#### Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

#### Directions

- adults and children 12 years and over:
- to **relieve**symptoms, swallow 1 tablet with a glass of water. Do not chew.
- to preventsymptoms, swallow 1 tablet with a glass of water at any time from 10 to 60 minutes before ating food or drinking beverages that cause heartburn
- do not use more than 2 tablets in 24 hours
- children under 12 years: ask a doctor

### Other information

- read the directions and warnings before use
- keep the carton. It contains important information.
- store at 20°-25°C (68°-77°F)
- protect from moisture

#### Inactive ingredients

carnauba wax, colloidal silicon dioxide, croscarmellose sodium, lactose monohydrate, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, talc, titanium dioxide7

#### **Questions or comments?**

1-800-719-9260

#### **Principal Display Panel**

CAREone™

mpare to the active ingredient in Maximum Strength Pepcid ®AC

MAXIMUM STRENGTH

ACID RELIEF

Famotidine Tablets, 20mg

Acid Reducer

Just one tablet prevents & relieves heartburn due to acid indigestion

See New Warnings

Gluten Free Actual Size OUR PHARMACISTS RECOMMEND 25 TABLETS



194A7 OF C2	Drug Facts Active ingredient Purpose (in each table t) Famotidine 20 mgAcid reducer	Drug Facts (contin ued) Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid red ucers may interact with certain prescription d u gs. Stop useand ask a doc tor if	he stomach groods and oking
	Uses relieves heartbum associated with acid indigestion and sour stomach prevents heartburn as sociated with acid indigestion and sour stomach brought on by eating or d inking certainfood and beverages	your heartburn continues or worsens you need to take this product for more than 14 days If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Centerright away. (1 + 800-222-1222)	fitting dothing around t Avoid heartburn-causin ohol, and certain fruits : Jose weight • Quit sm
DE AIA	Warrings   Allergy alert: Do not use if you are allergic to famotidine or other acid reducers   Do not use   If you have trouble or pain swallowing food, vomiting with bod, or bbody or black stools. These may be signs of a serious condition. See your doctor.   with other acid reducers   Ask adoctor before use if you have   had heartburn over 3 months. This may be a sign of a more serious condition.   had heartburn with lighthea dedness, sweating, or diziness   c hest painor shoulder pain with shortness of breath; sweatling; pain spreading to arms, neck or should ers; or libtheadedness	Directions adults and children 12 years and over: to relieve symptoms, swallow 1 tablet with a glass of water. Do not chew. to prevent symptoms, swallow 1 tablet with a glass of water at any time from 1 0 to 60 minutes before eating food or drinking beverages that cause heartburn to one use more than 2 tablets in 24 hours th idren under 12 years : ask a doctor	Tips for Managing Heartburn   • Do not le flator bendvoer after eating • Do not eastber bedime - states the head of your bed • Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, cafferine, alcohol, and certain fruits and vegetables   • East sowly and avoid big meals • If overweight, lose weight • Outh smoking
		Other information Fread the directions and warnings before use keep the carton. It contains important information. store at 20°-25°C (68°-77°F) protect from moisture	It aging Heartburn It aging Heartburn thefore beadtime - , spicy, fatty or fried • Eat slowly and av
	a rightheadedress ■ frequent chest pain ■ frequent wheezing, particularly with heartburn ■ unexplained weight bss ■ nausea orvomiting ■ stomachpain ■ kidney disease	Inactive ingredients carnauba wax, colbidal silicon dixxide, croscarmelbse sodium, lactose monohy drate, magnesium steazte, microcry stalline cellulose, polyethy lene glycol, polyvinyl alcohol, talc, titanium dioxide	Tips for Ma • Do not fer • Do not earlish vegetables
	Drug Facts (continued)     Questions or comments?1-800-719-9260		
-	*This product is not manufactured or distributed by Johnson & Johnson Consumer hc., distributer of Maximum Strength Pepcid® AC. DIS TR IBUT EDBY FOODHOLD U.S.A., LLC LAND OVER, MD20 785 -1-877-846-9949 © 2 020 Ahold D elhaize Licen sing Sàrl Guality guaranteed or your money back.	3 41520 31720 0	

# CAREONE ACID RELIEF

famotidine tablet, film coated					
Product Information					
Product Type	HUMAN OTC DRUG Item Code (Source) NDC:59368-29			368-293	
Route of Administration	ORAL				
Active Ingredient/Active Moiety					
Ingredient Name Basis of Strengt			ength	Strength	
FAMOTIDINE (UNII: 5QZ015J2Z8) (FAMOTIDINE - UNII:5QZ015J2Z8)   FAMOTIDINE				20 mg	
Inactive Ingredients					
Ingredient Name				9	Strength
CARNAUBA WAX (UNII: R12CBM0EIZ)					
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)					
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)					

LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POLYVINYL ALCOHOL, UNSPECIFIED (UNII: 532B59J990)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

# Product Characteristics

Color	white	Score	no score
Shape	ROUND	Size	8mm
Flavor		Imprint Code	L194
Contains			

## Packaging

#	ltem Code	Package DescriptionMarketing StartMarketDateDateDate			
1	NDC:59368-293- 01	1 in 1 CARTON	09/22/2016		
1		50 in 1 BOTTLE; Type 0: Not a Combination Product			
Marketing Information					
	Marketing	Application Number or Monograph	Marketing Start	Marketing End	

# CategoryApplication Number of Monograph<br/>CitationMarketing Start<br/>DateMarketing End<br/>DateANDAANDA07735109/22/2016

# Labeler - Praxis, LLC (016329513)

Establishment			
Name	Address	ID/FEI	Business Operations
Praxis, LLC		016329513	manufacture(59368-293) , label(59368-293) , pack(59368-293)

Revised: 1/2023

Praxis, LLC