NATUHEAL WART REMOVER- salicylic acid 17% wart remover liquid Jiangxi Hemei Pharmaceutical Co., Ltd

84010-084

Active Ingredient

Salicylic acid 17%

Purpose

Wart remover

Use

For the removal of common warts and plantar warts. Common warts have a rough "cauliflower-like" surface and are easily recognizable. Plantar warts are located only on the soles of the feet, tender to touch, with an interrupted footprint pattern.

Warnings

For external use only. Flammable. Keep away from fire and flame.

Do not use

on irritated skin or on any area that is infected or reddened On moles, birthmarks, warts with hair growing from them, genital warts, or warts on the face or mucous membranes. If you have diabetes or poor blood circulation.

When Using

Avoid contact with eyes. If product gets into the eye, flush with Water for 15 minutes. Non-edible.

Cap tightly and store at room temperature, away from heat.

Stop Use

Stop use and ask a doctor if discomfort persists.

Ask Doctor

Stop use and ask a doctor if discomfort persists.

Keep Oot Of Reach Of Children

If swallowed, get medical help or contact the Poison Control Center (1-800-222-1222) right away.

Directions

Wash the affected area. May soak wart in warm water for 5 minutes. Dry area thoroughly.

Use a cotton swab to apply a sufficient amount to cover each wart. Let it dry.Repeat this procedure once or twice daily as needed (until wart is removed) for up to 12 weeks.

Other information

Store at room temperature. Avoid excessive heat (37°C/99°F). Skin discoloration may occur during or after use.

Inactive ingredients

Water Polyethylene Glycol Glycerin Borneol Podofilox.

PRINCIPAL DISPLAY PANEL

WART REMOVER

Stop use and ask a doctor if discomfort persists.

Keep out of reach of children. If swallowed, get medical help or contact the Poison Control Center (1-800-222-1222) right away. **Directions**

Wash the affected area.

May soak wart in warm water for 5 minutes. Dry area thoroughly.

Use a cotton swab to apply a sufficient amount to cover each wart. Let it dry. Repeat this procedure once or twice daily as needed (until wart is removed) for up to 12 weeks.

Other information

Store at room temperature. Avoid excessive heat (37°C / 99°F). Skin discoloration may occur during or

Inactive ingredients:

Water52.5%
Polyethylene Glycol · · · · · · · · 16%
Glycerin · · · · · · · · · · · · · · · · · · ·
Borneol
Podofilox 0.5%



NATUHEAL®

WART REMOVER

HELPS REMOVE WARTS LIQUIDEASY TO USE NATURAL FORMULA HELPS WITH WART



- PENETRATES&REMOVES WART
- ADVANCED TECHNOLOGY
 The liquid turns into a mask
 in seconds
- SAFE & HYGIENICD oesn't hurt dermis and prevents spreading

NET: 1 fl oz (30mL)

Manufacturer: Jiangxi Hemei PharmaceuticalCo., Ltd. Address:4th Floor, Building 1956, Phase III, Zhangjiashan Medical Incubation Park, Zhangshu, Yichun City, Jiangxi 331208, China(CHN)

EC REP

SUCCESS COURIER SL.
CALLE RIO TORMES NUM.1,
PLANTA 1,DERECHA,OFICINA
3, Fuenlabrada,Madrid,28947
Spain+34 910602659
successservice2@hotmail.com

UK REP

EVATOST CONSULTING LTD. Office 10132 Threadneedle Street, London, United Kingdom, EC2R 8AY +442076494690 contact@evatost.com







MADE IN CHINA
SHELF LIFE: 3 years

Drug Facts

Active ingredient:

Purpose

Salicylic acid 17%······Wart remover

Uses

For the removal of common warts and plantar warts.

Common warts have a rough "cauliflower -like" surface and are easily recognizable. Plantar warts are located only on the soles of the feet, tender to touch, with an interrupted footprint pattern.

Warnings

For external use only. Flammable. Keep away from fire and flame.

Do not use

on irritated skin or on any area that is infected or reddened.On moles, birthmarks, warts with hair growing from them, genital warts, or warts on the face or mucous membranes.If you have diabetes or poor blood circulation.

When using this product

Avoid contact with eyes. If product gets into the eye, flush with Water for 15 minutes.Non-edible.

Cap tightly and store at room temperature, away from heat.

NATUHEAL WART REMOVER

salicylic acid 17% wart remover liquid

Product Information

Product Type HUMAN OTC DRUG Item Code (Source) NDC:84010-084

Route of Administration TOPICAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
SALICYLIC ACID (UNII: O414PZ4LPZ) (SALICYLIC ACID - UNII:O414PZ4LPZ)	SALICYLIC ACID	17 g in 100 mL

Inactive Ingredients	
Ingredient Name	Strength
WATER (UNII: 059QF0KO0R)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
PODOFILOX (UNII: L36H50F353)	
GLYCERIN (UNII: PDC6A3C0OX)	
BORNEOL (UNII: M89NIB437X)	

Packaging				
# Ite	m Code	Package Description	Marketing Start Date	Marketing End Date
1 NDC:8	84010-084-	30 mL in 1 BOTTLE; Type 0: Not a Combination Product	03/07/2025	

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M028	03/07/2025	

Labeler - Jiangxi Hemei Pharmaceutical Co., Ltd (724892056)

Establishment					
Name	Address	ID/FEI	Business Operations		
Jiangxi Hemei Pharmaceutical Co., Ltd		724892056	manufacture(84010-084)		

Revised: 3/2025 Jiangxi Hemei Pharmaceutical Co., Ltd