

EQUALACTIN LAXATIVE- calcium polycarbophil tablet
NUMARK BRANDS, INC

Drug Facts

Active Ingredients

Calcium polycarbophil 625 mg
(equivalent to 500 mg polycarbophil)

Purpose

Bulk for mixing Laxative

Use

for relief of occasional constipation and irregularity. This product generally produces bowel movement in 12 - 72 hours.

Warnings

Choking: Taking this product without adequate fluid may cause it to swell and block your throat or esophagus and may cause choking. Do not take this product if you have difficulty in swallowing. If you experience chest pain, vomiting, or difficulty in swallowing or breathing after taking this product, seek immediate medical attention.

Do not use

this product for more than 1 week unless directed by a doctor

Ask a doctor before use if you have

- abdominal pain
- nausea
- vomiting
- a sudden change in bowel habits over a period of 2 weeks

Ask a doctor or pharmacist before use if you are

- taking an antibiotic containing tetracycline as this product contains calcium
- taking any other drug. Take this product 2 or more hours before or after other drugs. All laxatives may affect how other drugs work.

Stop use and ask a doctor if

- rectal bleeding occurs
- you fail to have bowel movement after use of laxative. These may indicate a serious

condition

As with any drug, if you are pregnant or nursing a baby, seek the advice of a health professional before using this product

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- drink a full glass (8 fl. oz.) of liquid with each dose
- chew tablets thoroughly
- do not swallow tablets whole

adults and children 12 years of age and over	up to 8 tablets daily in divided doses of 2 tablets per dose
children 6 to under 12 years of age	up to 4 tablets daily in divided doses of 1 tablet per dose
children 2 to under 6 years of age	up to 2 tablets daily in divided doses of 1 tablet per dose
children under 2 years of age	consult a doctor

Other information

- each tablet contains calcium 125 mg
- store at room temperature
- avoid excessive humidity

Inactive ingredients citric acid, crospovidone, dextrose, flavor, magnesium stearate, maltodextrin, microcrystalline cellulose, sucralose

Questions? 1-800-331-0221

PRINCIPAL DISPLAY PANEL

**Equalactin®
Bulk Forming Laxative**

Relieves Symptoms of Nervous Stomach

- Irregularity
- Abdominal Discomfort
- Loose Bowels
- Constipation

48 CITRUS FLAVORED CHEWABLE TABLETS

Nervous Stomach (similar to Irritable Bowel), with its symptoms of loose bowels, irregularity, abdominal discomfort and constipation, is caused by water imbalance in the bowel, and afflicts millions. Equalactin provides relief by equalizing the water balance to restore normal bowel movements. When you feel the onset of nervous stomach symptoms, take Equalactin to provide relief and prevent the episode from becoming severe.

Tamper evident: Do not use if carton is open or if printed seal on blister is broken or missing.

Distributed by: Numark Brands
138 East Main Street, Suite 201
Kent, OH 44240
Rev 7/25 10001C

EQ-48FC

LOT:

EXP:

Equalactin®

Equalactin®

Equalactin®

Bulk Forming Laxative

Relieves Symptoms of Nervous Stomach



- Irregularity
- Loose Bowels
- Abdominal Discomfort
- Constipation

48 CITRUS FLAVORED CHEWABLE TABLETS



Nervous Stomach (similar to irritable bowel), with its symptoms of loose bowels, irregularity, abdominal discomfort and constipation, is caused by water imbalance in the bowel, and afflicts millions. Equalactin provides relief by equalizing the water balance to restore normal bowel movements. When you feel the onset of nervous stomach symptoms, take Equalactin to provide relief and prevent the episode from becoming severe.



Tamper evident: Do not use if carton is open or if printed seal on blister is broken or missing.

Distributed by: Numark Brands
138 East Main Street, Suite 201
Kent, OH 44240
Rev 7/25 10001C

EQ-48FC

LOT:

EXP:

Drug Facts	
Active ingredients	Calcium polycarphyl 625 mg (equivalent to 500 mg polycarphyl)
Purpose	Bulk forming laxative
Use For relief of occasional constipation and irregularity. This product generally produces bowel movement in 12 - 72 hours.	
Warnings	
Choking: Taking this product without adequate fluid may cause it to swell and block your throat or esophagus and may cause choking. Do not take this product if you have difficulty in swallowing. If you experience chest pain, vomiting, or difficulty in swallowing or breathing after taking this product, seek immediate medical attention.	
Do not use this product for more than 1 week unless directed by a doctor.	
Ask a doctor before use if you have ■ abdominal pain ■ nausea ■ vomiting ■ a sudden change in bowel habits over a period of 2 weeks	
Ask a doctor or pharmacist before use if you are ■ taking an antibiotic containing tetracycline as this product contains calcium ■ taking any other drug. Take this product 2 or more hours before or after other drugs. All laxatives may affect how other drugs work.	
Stop use and ask a doctor if ■ rectal bleeding occurs ■ you fail to have bowel movement after use of laxative. These may indicate a serious condition.	
As with any drug, if you are pregnant or nursing a baby, ask the advice of a health professional before using this product.	
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.	
Directions ■ Drink a full glass (8 oz.) of fluid with each dose ■ Chew tablets thoroughly ■ Do not swallow tablets whole	
adults and children 12 years of age and over	up to 8 tablets daily in divided doses of 2 tablets per dose
children 6 to under 12 years of age	up to 4 tablets daily in divided doses of 1 tablet per dose
children 2 to under 6 years of age	up to 2 tablets daily in divided doses of 1 tablet per dose
children under 2 years of age	consult a doctor
Other information ■ Each tablet contains calcium 125 mg ■ Store at room temperature ■ Avoid excessive humidity	
Inactive ingredients cholic acid, croscollon, dextrose, flavor, magnesium stearate, methylcellulose, microcrystalline cellulose, sucralose	
Questions? 1-800-214-2379	

EQUALACTIN LAXATIVE

calcium polycarbophil tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:69846-000
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
Calcium Polycarbophil (UNII: 8F049NKY49) (Polycarbophil - UNII:W25LM17A4W)	Calcium Polycarbophil	625 mg

Inactive Ingredients

Ingredient Name	Strength
Citric Acid Monohydrate (UNII: 2968PHW8QP)	
Crospovidone (UNII: 2S7830E561)	
Dextrose, Unspecified Form (UNII: IY9XDZ35W2)	
Magnesium Stearate (UNII: 70097M6I30)	
Maltodextrin (UNII: 7CVR7L4A2D)	
Cellulose, Microcrystalline (UNII: OP1R32D61U)	
Sucralose (UNII: 96K6UQ3ZD4)	

Product Characteristics

Color	WHITE	Score	no score
Shape	ROUND	Size	16mm
Flavor	LEMON	Imprint Code	E
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:69846-000-26	2 in 1 CARTON	11/21/1986	03/03/2023
1		12 in 1 BLISTER PACK; Type 0: Not a Combination Product		
2	NDC:69846-000-27	4 in 1 CARTON	11/21/1986	
2		12 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M007	11/21/1986	

