

**NIGHT-TIME COLD AND FLU RELIEF- acetaminophen, doxylamine succinate, dextromethorphan hbr capsule, liquid filled  
HealthLife of USA**

-----  
**Night-Time Cold & Flu Relief**

**Active ingredients (in each Softgels)**

Acetaminophen 325 mg

Dextromethorphan HBr 15 mg

Doxylamine succinate 6.25 mg

**Purpose**

Pain reliever/Fever reducer

Cough suppressant

Antihistamine

**Uses**

temporarily relieves common cold/flu symptoms:

- cough due to minor throat and bronchial irritation
- sore throat
- headache
- minor aches and pains
- fever
- runny nose and sneezing

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4 doses in 24 hours, which is the maximum daily amount for this product
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

**Allergy Alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- Skin reddening
- Blisters
- Rash

If a skin reaction occurs, stop use and seek medical help right away.

**Sore Throat warning:** If sore throat is severe, lasts for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a

doctor promptly.

### **Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription).
- if you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

### **Ask a doctor before use if you have**

- liver disease
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis or emphysema
- trouble urinating due to enlarged prostate gland

### **Ask a doctor or pharmacist before use if you are**

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

### **When using this product**

- excitability may occur, especially in children
- marked drowsiness may occur
- avoid alcoholic drink
- be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives, and tranquilizers may increase drowsiness

### **Stop use and ask a doctor if**

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts.

These could be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use.

### **Keep out of reach of children**

In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

### **Direction**

- take only as directed
- do not exceed 4 doses per 24 hours

---

adults & children 12 yrs & over	2 softgels with water every 6 hrs
children 4 to under 12 yrs	ask a doctor
children under 4 yrs	do not use

---

### **Other information**

- store at room temperature.
- TAMPER EVIDENT: This package is safety sealed & child resistant. Use only if blisters are intact. If difficult to open, use scissors.
- KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION

### **Inactive ingredients**

D&C Yellow No.10, FD&C Blue No. 1, gelatin, glycerin, polyethylene glycol, povidone, propylene glycol, purified water, sorbitol sorbitan solution, titanium dioxide, methyl paraben, propyl paraben.

### **Questions or comments?**

1-844-832-1138

M-F 9am-5pm EST

### **Distributed by:**

Health life of USA LLC  
Rahway, NJ 07065  
www.healthlifeus.com

\*This product is not manufactured or distributed by Procter & Gamble, Distributor of Vicks Nyquil<sup>®</sup>

### **PACKAGE LABEL**

**NDC 69517-161-10**

**10 Softgels**

### **Night-Time Cold & Flu Relief**

Acetaminophen, Doxylamine Succinate, Dextromethorphan HBr

- Headache, Fever, Sore Throat, Minor Aches & Pains
- Sneezing, Runny Nose
- Cough

**\*Compare to the active ingredients in Vicks Nyquil<sup>®</sup>**



# Night-Time

Cold & FLU

NDC 69517-161-10

## POWERFUL Nighttime RELIEF



# Night-Time

## COLD & FLU

### Nighttime Relief

#### Acetaminophen, Doxylamine Succinate, Dextromethorphan HBr

- Headache, Fever, Sore Throat, Minor Aches & Pains
- Sneezing, Runny Nose
- Cough



### 10 Softgels

*\*Compare to the active ingredients in Vicks Nyquil®*

**TAMPER EVIDENT:** This package is easily sealed & child resistant. Use only if blister are intact. If difficult to open, use scissors.

**Parents:**  
Learn about teen medicine abuse  
[www.stopmedicineabuse.org](http://www.stopmedicineabuse.org)

**Drug Facts (Continued)**  
Questions? 1-844-832-1138  
Distributed by:  
Health Life of USA, LLC  
Rahway, NJ 07065  
[www.healthlifeusa.com](http://www.healthlifeusa.com)

NO VARNISH  
AREA

Lot. No.:  
Exp. Date:

Questions or comments?  
1-844-832-1138 M-F 9am-5pm EST  
**KEEP OUTER CARTON FOR COMPLETE  
WARNINGS AND PRODUCT INFORMATION**  
\*This product is not manufactured or distributed by  
Procter & Gamble, Distributor of Vicks Nyquil®

#### Drug Facts (continued)

##### Directions

- Take only as directed
  - do not exceed 4 doses per 24 hours
- |                                 |                                   |
|---------------------------------|-----------------------------------|
| adults & children 12 yrs & over | 2 softgels with water every 6 hrs |
| children 4 to under 12 yrs      | ask a doctor                      |
| children under 4 yrs            | do not use                        |

#### Drug Facts (continued)

##### Other information

- store at room temperature

Inactive ingredients: D&C Yellow No. 10, FD&C Blue No. 1, gelatin, glycerin, polyethylene glycol, polydioxane, propylene glycol, purified water, sorbitol sorbitan solution, titanium dioxide, methyl paraben, propyl paraben

#### Drug Facts

#### Purpose

##### Active Ingredients (In each Softgel)

Acetaminophen 325 mg	Pain reliever/Fever reducer
Dextromethorphan HBr 15 mg	Cough suppressant
Doxylamine succinate 6.25 mg	Antihistamine

##### Uses temporarily relieves common cold/flu symptoms:

- cough due to minor throat and bronchial irritation
- sore throat
- headache
- minor aches and pains
- fever
- runny nose and sneezing

##### Warnings

**Liver warning:** This product contains acetaminophen.

Severe liver damage may occur if you take

- more than 4 doses in 24 hrs, which is the maximum daily amount for this product
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

**Allergy Alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- Skin reddening
- Blisters
- Rash

If a skin reaction occurs, stop use and seek medical help right away.

**Sore Throat warning:** If sore throat is severe, lasts for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use:** • with any other drug containing acetaminophen (prescription or nonprescription).

If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

• If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have:** • liver disease • glaucoma

- cough that occurs with too much phlegm (mucus)
- a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis or emphysema
- trouble urinating due to enlarged prostate gland

**Ask a doctor or pharmacist before use if you are:**

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

##### When using this product

- excitability may occur, especially in children
- marked drowsiness may occur
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives, & tranquilizers may increase drowsiness

**Stop use and ask a doctor if:** • pain or cough gets worse or lasts more than 7 days

- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts.

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.





## NIGHT-TIME COLD AND FLU RELIEF

acetaminophen, doxylamine succinate, dextromethorphan hbr capsule, liquid filled

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:69517-161
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	15 mg
<b>DOXYLAMINE SUCCINATE</b> (UNII: V9B19B5Y12) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	6.25 mg

### Inactive Ingredients

Ingredient Name	Strength
<b>FD&amp;C BLUE NO. 1</b> (UNII: H3R47K3TBD)	
<b>D&amp;C YELLOW NO. 10</b> (UNII: 35SW5USQ3G)	
<b>GELATIN</b> (UNII: 2G86QN327L)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>POLYETHYLENE GLYCOL 400</b> (UNII: B697894SGQ)	
<b>POVIDONE K30</b> (UNII: U725QWY32X)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>SORBITOL</b> (UNII: 506T60A25R)	
<b>METHYLPARABEN</b> (UNII: A2I8C7HI9T)	
<b>PROPYLPARABEN</b> (UNII: Z8IX2SC1OH)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	

### Product Characteristics

<b>Color</b>	green	<b>Score</b>	no score
<b>Shape</b>	CAPSULE	<b>Size</b>	21mm
<b>Flavor</b>		<b>Imprint Code</b>	SN3
<b>Contains</b>			

### Packaging

Marketing Start      Marketing End

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:69517-161-10	1 in 1 CARTON	04/04/2023	
1		10 in 1 BLISTER PACK; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	04/04/2023	

**Labeler** - HealthLife of USA (079656178)

**Registrant** - HealthLife of USA (079656178)

Revised: 10/2023

HealthLife of USA