

**HISTEX-DM NEW FORMULATION- dextromethorphan hbr, pseudoephedrine hci, and triprolidine hci syrup**  
**Allegis Pharmaceuticals, LLC**

-----  
**HISTEX™-DM Syrup New Formulation**

***Drug Facts***

<b><i>Active ingredients (in each 5 mL teaspoonful)</i></b>	<b><i>Purpose</i></b>
Dextromethorphan HBr 20 mg	Cough Suppressant
Pseudoephedrine HCl 30 mg	Nasal Decongestant
Tripolidine HCl 2.5 mg	Antihistamine

**Uses**

temporarily relieves these symptoms due to common cold, hay fever (allergic rhinitis) or other upper respiratory allergies:

- cough due to minor throat or bronchial irritation
- runny nose
- sneezing
- itching of the nose or throat
- itchy, watery eyes
- nasal congestion
- reduces swelling of nasal passages

**Warnings**

**Do not exceed recommended dosage.**

**Do not use this product**

- If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before giving this product.

**Ask a doctor before use if you have**

- a breathing problem such as emphysema or chronic bronchitis
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- glaucoma
- a persistent or chronic cough such occurs with smoking, asthma, chronic bronchitis, or emphysema



- a cough that occurs with too much phlegm (mucus)
- trouble urinating due to an enlarged prostate gland

**Ask a doctor before use if you are taking sedatives or tranquilizers**

### **When using this product**

- excitability may occur, especially in children
- may cause drowsiness
- alcohol, sedatives and tranquilizers may increase the drowsiness effect
- avoid alcoholic beverages
- use caution when driving a motor vehicle or operating machinery

### **Stop use and ask a doctor if**

- nervousness, dizziness, or sleeplessness occur
- cough or nasal congestion persists for more than 1 week, tends to recur or is accompanied by a fever, rash or persistent headache.
- new symptoms occur

These could be signs of a serious condition.

**If pregnant or breast-feeding**, ask a health professional before use.

**Keep out of the reach of children.**

**In case of overdose, get medical help or contact a Poison Control Center right away.**

### **Directions**

**Do not exceed recommended dosage.**

<b>AGE</b>	<b>DOSE</b>
Adults and Children 12 years of age and older:	1 teaspoonful (5 mL) every 4 hours, not to exceed 4 teaspoonfuls (20 mL) in 24 hours, or as directed by a doctor
Children 6 to under 12 years of age	½ teaspoonful (2.5 mL) every 4 hours, not to exceed 2 teaspoonfuls (10 mL) in 24 hours, or as directed by a doctor.
Children under 6 years of age	Consult a doctor

### **Other Information**

Store at 15°-30° C (59°-86° F).

Tamper evident by foil seal under cap. Do not use if foil seal is missing or broken.



**Inactive ingredients**

Citric Acid, Glycerin, Grape Flavor, Propylene Glycol, Purified Water, Sodium Benzoate, Sodium Citrate, Sodium Saccharin, Sorbitol Solution.

**Questions? Comments?**

**Call 1-866-633-9033.**

**Manufactured for: Allegis Pharmaceuticals, Inc.**

**Canton MS 39046**

**[www.allegispharma.com](http://www.allegispharma.com)**

**Rev. 01/2025**

**PRINCIPAL DISPLAY PANEL - 473 mL Bottle Label**

**NDC 28595-310-16**

**Antihistamine - Nasal Decongestant - Cough Suppressant**

**HISTEX™-DM New Formulation  
Syrup**

**Each teaspoonful (5 mL)**

**contains:**

Dextromethorphan HBr 20 mg

Pseudoephedrine HCl 30mg

Tripolidine HCl 2.5 mg

**Sugar-Free - Dye Free - Alcohol Free**

**Grape Flavor**

**16 fl oz (473 mL)**



NDC 28595-310-16

Antihistamine • Nasal Decongestant  
Cough Suppressant

NEW FORMULATION

HISTEX-DM  
Syrup

Each teaspoonful (5 mL)  
contains:

Dextromethorphan HBr ... 20 mg  
Pseudoephedrine HCl ..... 30 mg  
Triprolidine HCl ..... 2.5 mg

Sugar-Free • Dye Free  
Alcohol Free

Grape Flavor

Tamper evident by foil seal under cap.  
Do not use if foil seal is missing or broken.

3

2859531016

9

16 fl oz (473 mL)

Drug Facts

Active ingredients

(in each 5 mL teaspoonful)

Dextromethorphan HBr 20 mg.....Cough Suppressant  
Pseudoephedrine HCl 30 mg.....Nasal Decongestant  
Triprolidine HCl 2.5 mg.....Antihistamine

Uses

temporarily relieves these symptoms due to common cold,  
hay fever (allergic rhinitis) or other upper respiratory allergies:

■ cough due to minor throat and bronchial irritation  
■ runny nose ■ sneezing ■ itching of the nose or throat  
■ itchy, watery eyes ■ nasal congestion  
■ reduces swelling of nasal passages

Warnings

Do not exceed recommended dosage.

Do not use this product

■ If you are now taking a prescription monoamine oxidase  
inhibitor (MAOI) (certain drugs for depression, psychiatric,  
or emotional conditions, or Parkinson's disease), or for 2  
weeks after stopping the MAOI drug. If you do not know if  
your prescription drug contains an MAOI, ask a doctor or  
pharmacist before giving this product.

Ask a doctor before use if you have

■ a breathing problem such as emphysema or chronic bronchitis  
■ heart disease ■ high blood pressure  
■ thyroid disease ■ diabetes ■ glaucoma  
■ a persistent or chronic cough such occurs with smoking,  
asthma, chronic bronchitis, or emphysema  
■ a cough that occurs with too much phlegm (mucus)  
■ trouble urinating due to an enlarged prostate gland

Ask a doctor before use if you are taking sedatives or tranquilizers

When using this product

■ excitability may occur, especially in children  
■ may cause marked drowsiness  
■ alcohol, sedatives and tranquilizers may increase the  
drowsiness effect  
■ avoid alcoholic beverages

Drug Facts (continued)

■ use caution when driving a motor vehicle or operating  
machinery

Stop use and ask a doctor if

■ nervousness, dizziness, or sleeplessness occur  
■ cough or nasal congestion persists for more than 1 week,  
tends to recur or is accompanied by a fever, rash or persistent  
headache.  
■ new symptoms occur  
These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional  
before use.

Keep out of the reach of children.

In case of overdose, get medical help or contact a  
Poison Control Center right away.

Directions Do not exceed recommended dosage.

AGE	DOSE
Adults and Children 12 years of age and older	1 teaspoonful (5 mL) every 4 hours, not to exceed 4 teaspoonfuls (20 mL) in 24 hours, or as directed by a doctor
Children 6 to under 12 years of age	½ teaspoonful (2.5 mL) every 4 hours, not to exceed 2 teaspoonfuls (10 mL) in 24 hours, or as directed by a doctor.
Children under 6 years of age	Consult a doctor

Other Information

Store at 15°-30° C (59°-86° F).  
Tamper evident by foil seal under cap. Do not use if foil  
seal is missing or broken.

Inactive ingredients Citric acid, glycerin, grape flavor,  
propylene glycol, purified water, sodium benzoate, sodium citrate, sodium  
saccharin, sorbitol solution

Questions? Comments? Call 1-866-633-9033.

Manufactured for: Allegis Pharmaceuticals, Inc.  
www.allegispharma.com Canton, MS 39046 Rev. 01/2025

HISTEX-DM NEW FORMULATION

dextromethorphan hbr, pseudoephedrine hci, and triprolidine hci syrup

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:28595-310
Route of Administration	ORAL		
Active Ingredient/Active Moiety			
Ingredient Name		Basis of Strength	Strength
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)		DEXTROMETHORPHAN HYDROBROMIDE	20 mg in 5 mL
PSEUDOEPHEDRINE HYDROCHLORIDE (UNII: 6V9V2RYJ8N) (PSEUDOEPHEDRINE - UNII:7CUC9DDI9F)		PSEUDOEPHEDRINE HYDROCHLORIDE	30 mg in 5 mL
TRIPROLIDINE HYDROCHLORIDE (UNII: YAN7R5L890) (TRIPROLIDINE - UNII:2L8T9S52QM)		TRIPROLIDINE HYDROCHLORIDE	2.5 mg in 5 mL
Inactive Ingredients			
Ingredient Name			Strength
CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)			
GLYCERIN (UNII: PDC6A3C0OX)			
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)			
WATER (UNII: 059QF0KO0R)			
SODIUM CITRATE, UNSPECIFIED FORM (UNII: 1Q73Q2JULR)			
SACCHARIN SODIUM (UNII: SB8ZUX40TY)			



<b>SORBITOL</b> (UNII: 506T60A25R)				
<b>SODIUM BENZOATE</b> (UNII: OJ245FE5EU)				
<b>GRAPE</b> (UNII: 6X543N684K)				
<b>Product Characteristics</b>				
<b>Color</b>			<b>Score</b>	
<b>Shape</b>			<b>Size</b>	
<b>Flavor</b>		GRAPE	<b>Imprint Code</b>	
<b>Contains</b>				
<b>Packaging</b>				
<b>#</b>	<b>Item Code</b>	<b>Package Description</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
1	NDC:28595-310-16	473 mL in 1 BOTTLE; Type 0: Not a Combination Product	03/10/2025	
<b>Marketing Information</b>				
<b>Marketing Category</b>		<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
OTC Monograph Drug		M012	03/10/2025	

**Labeler -**
Allegis Pharmaceuticals, LLC (792272861)