

RUGBY CHILDRENS ACETAMINOPHEN- acetaminophen suspension

Rugby Laboratories

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Rugby Laboratories Children's Acetaminophen Drug Facts

Active ingredient (in each 5 mL)

Acetaminophen 160 mg

Purpose

Pain reliever/fever reducer

Uses

temporarily:

- reduces fever
- relieves minor aches and pains due to:
- the common cold
- flu
- headache
- sore throat
- toothache

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if your child takes

- more than 5 doses in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

- if your child has ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if your child has

liver disease

Ask a doctor or pharmacist before use if your child is

taking the blood thinning drug warfarin

When using this product

do not exceed recommended dose (see overdose warning)

Stop use and ask a doctor if

- pain gets worse or lasts more than 5 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical even if you do not notice any signs or symptoms.

Directions

- **this product does not contain directions or complete warnings for adult use**
- **do not give more than directed (see overdose warning)**
- **shake well before using**
- mL = milliliter
- find right dose on chart below. If possible, use weight to dose; otherwise, use age.
- remove the child protective cap and squeeze your child's dose into the dosing cup
- repeat dose every 4 hours while symptoms last
- do not give more than 5 times in 24 hours

Weight (lb)	Age (yr)	Dose (mL)*
under 24	under 2 years	ask a doctor
24-35	2-3 years	5 mL
36-47	4-5 years	7.5 mL
48-59	6-8 years	10 mL
60-71	9-10 years	12.5 mL
72-95	11 years	15 mL

*or as directed by a doctor

Attention: use only enclosed dosing cup specifically designed for use with this product. Do not use any other dosing device.

Other information

- **each 5 mL contains :** sodium 3 mg
- store at 20-25°C (68-77°F)
- **do not use if printed neckband is broken or missing**

Inactive ingredients

anhydrous citric acid, butylparaben, calcium sulfate, carrageenan, D&C red #33, FD&C blue #1, flavor, glycerin, high fructose corn syrup, hydroxyethyl cellulose, microcrystalline cellulose and carboxymethylcellulose sodium, propylene glycol, purified water, sodium benzoate, sorbitol solution, tribasic sodium phosphate

Questions or comments?

1-800-645-2158

Principal Display Panel

Compare to the active ingredient in Children's Tylenol® Oral Suspension

Children's

Acetaminophen

160 mg per 5 mL

Oral Suspension

Pain Reliever/Fever Reducer

Fast, effective relief of children's pain and fever

Ibuprofen Free


Alcohol Free

Aspirin Free

Grape Flavor

For Ages 2 to 11 Years


4 FL OZ (118 mL)



Rugby®

Children's
Acetaminophen
160 mg per 5 mL
Oral Suspension

Pain Reliever/Fever Reducer



Drug Facts

Active ingredient (in each 5 mL) **Purpose**
 Acetaminophen 160 mg.....Pain reliever/fever reducer

Uses temporarily:
 ■ reduces fever
 ■ relieves minor aches and pains due to:
 ■ the common cold ■ flu ■ headache
 ■ sore throat ■ toothache

Warnings
Liver warnings: This product contains acetaminophen. Severe liver damage may occur if your child takes more than 5 doses in 24 hours, which is the maximum daily amount.
 ■ with other drugs containing acetaminophen
Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:
 ■ skin reddening ■ blisters ■ rash
 If a skin reaction occurs, stop use and seek medical help right away.
Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use
 ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
 ■ if your child has ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if your child has liver disease
Ask a doctor or pharmacist before use if your child is taking the blood thinning drug warfarin

When using this product do not exceed recommended dose (see overdose warning)

Stop use and ask a doctor if
 ■ pain gets worse or lasts more than 5 days
 ■ fever gets worse or lasts more than 3 days
 ■ new symptoms occur
 ■ redness or swelling is present
 These could be signs of a serious condition.

Keep out of reach of children.
Overdose warnings: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical even if you do not notice any signs or symptoms.

Drug Facts (continued)

Directions
 ■ this product does not contain directions or complete warnings for adult use
 ■ do not give more than directed (see overdose warning)
 ■ shake well before using
 ■ mL = milliliter
 ■ find right dose on chart below. If possible, use weight to dose; otherwise, use age.
 ■ remove the child protective cap and squeeze your child's dose into the dosing cup
 ■ repeat dose every 4 hours while symptoms last
 ■ do not give more than 5 times in 24 hours

Weight (lb)	Age (yr)	Dose (mL)*
under 24	under 2 years	ask a doctor
24-35	2-3 years	5 mL
36-47	4-5 years	7.5 mL
48-59	6-8 years	10 mL
60-71	9-10 years	12.5 mL
72-95	11 years	15 mL

*or as directed by a doctor
Attention: use only enclosed dosing cup specifically designed for use with this product. Do not use any other dosing device.

Other information
 ■ each 5 mL contains: sodium 3 mg
 ■ store at 20-25°C (68-77°F)
 ■ do not use if printed neckband is broken or missing

Inactive ingredients
 anhydrous citric acid, butylparaben, calcium sulfate, carrageenan, D&C red #33, FD&C blue #1, flavor, glycerin, high fructose corn syrup, hydroxyethyl cellulose, microcrystalline cellulose and carboxymethylcellulose sodium, propylene glycol, purified water, sodium benzoate, sorbitol solution, tribasic sodium phosphate

Questions or comments?
 1-800-645-2158

**This product is not manufactured or distributed by Johnson & Johnson Consumer Inc., distributor of Children's Tylenol® Oral Suspension.
 Distributed by:
 RUGBY® LABORATORIES
 17177 N Laurel Park Drive, Suite 233
 Livonia, MI 48152
 www.rugbylaboratories.com

Rugby® NDC 0536-1321-97
 Compare to the active ingredient in Children's Tylenol® Oral Suspension**

Children's
Acetaminophen
160 mg per 5 mL
Oral Suspension

Pain Reliever/Fever Reducer

Fast, effective relief of children's pain and fever

Ibuprofen Free
Alcohol Free
Aspirin Free



Grape Flavor
For Ages 2 to 11 Years

4 FL OZ (118 mL)

Rugby®

Children's
Acetaminophen
160 mg per 5 mL
Oral Suspension

Pain Reliever/Fever Reducer

DIRECTIONS:

STEP 1. Shake well before using



STEP 2. Remove child protective cap




STEP 3. Squeeze your child's dose into the dosing cup



IMPORTANT:
 Please see dosing chart or ask a doctor for your child's correct dose.

Keep carton, it contains important information.




Rev. 09/20 R-05 Re-order No. 371066



CODE AREA

13026 8T C 1



RUGBY CHILDRENS ACETAMINOPHEN
 acetaminophen suspension

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0536-1321
Route of Administration	ORAL		
Active Ingredient/Active Moiety			

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	160 mg in 5 mL

Inactive Ingredients	
Ingredient Name	Strength
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
BUTYLPARABEN (UNII: 3QP1IU3FV8)	
CALCIUM SULFATE, UNSPECIFIED FORM (UNII: WAT0DDB505)	
CARRAGEENAN (UNII: 5C69YCD2YJ)	
D&C RED NO. 33 (UNII: 9DBA0SBB0L)	
FD&C BLUE NO. 1 (UNII: HBR47K3TBD)	
GLYCERIN (UNII: PDC6A3C0OX)	
HIGH FRUCTOSE CORN SYRUP (UNII: XY6UN3QB6S)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
CARBOXYMETHYLCELLULOSE SODIUM, UNSPECIFIED FORM (UNII: K679OBS311)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SORBITOL (UNII: 506T60A25R)	
SODIUM PHOSPHATE, TRIBASIC (UNII: A752Q30A6X)	

Product Characteristics			
Color	PURPLE (viscous)	Score	
Shape		Size	
Flavor	GRAPE	Imprint Code	
Contains			

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0536-1321-97	1 in 1 CARTON	10/21/2020	
1		118 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part343	10/21/2020	

Labeler - Rugby Laboratories (079246066)