

SIGNATURE SELECT EFFERVESCENT PAIN RELIEF- aspirin tablet, effervescent
Safeway Inc.

Signature Select Effervescent Pain Relief

Active ingredient

(in each effervescent tablet)

Buffered aspirin 325 mg **(NSAID)***

*Nonsteroidal anti-inflammatory drug

Purpose

(in each effervescent tablet)

Buffered aspirin 325 mg **(NSAID)***....Pain reliever/fever reducer

*Nonsteroidal anti-inflammatory drug

Uses

■ for the temporary relief of minor aches and pains due to:

■ headache ■ muscle pain ■ backache

■ toothache ■ menstrual pain ■ colds

■ minor arthritis pain

■ temporarily reduces fever

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction which may include:

■ hives ■ facial swelling ■ asthma (wheezing) ■ shock

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic

drinks every day while using this product ■ take more or for a longer time than directed

Do not use

Do not use ■ if you are allergic to aspirin or any other pain reliever/fever reducer ■ if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if

■ stomach bleeding warning applies to you ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease ■ you are taking a diuretic ■ you have asthma ■ you have a sodium-restricted diet

Ask a doctor or pharmacist before use if you are

Ask a doctor or pharmacist before use if you are taking a prescription drug for ■ diabetes ■ gout ■ arthritis

When using this product

When using this product do not exceed recommended dosage

Stop use and ask a doctor if

Stop use and ask a doctor if ■ you experience any of the following signs of stomach bleeding

■ feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better

■ an allergic reaction occurs. Seek medical help right away.

■ pain lasts for more than 10 days or gets worse

■ fever gets worse or lasts more than 3 days

■ new symptoms occur

■ redness or swelling is present

■ ringing in the ears or a loss of hearing occurs

If pregnant or breast-feeding

If pregnant or breast-feeding, ask a health professional before use. **It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.**

Keep out of reach of children.

In case of overdose

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not exceed recommended dosage
- to avoid serious injury, completely dissolve tablets in 4 oz. of water before taking

adults and children 12 years and over: 2 tablets every 4 hours, or as directed by a doctor - do not exceed 8 tablets in 24 hours

60 years and over: 2 tablets every 4 hours, or as directed by a doctor - do not exceed 4 tablets in 24 hours

children under 12 years: consult a doctor

Take this product any time - morning, noon or night - when you need relief from headaches, backaches, body aches or muscle pain.

Place two (2) tablets in 4 oz. of water. Dissolve tablets completely.

You do not have to drink any of the residue that may be on the bottom of the glass. The medicines are in the water.

Other information

- **each tablet contains:** sodium 524 mg
- store at room temperature (59°-86°F). Avoid excessive heat.

Inactive ingredients

anhydrous citric acid, sodium bicarbonate

Questions or comments?

Questions or comments? 1-888-723-3929

1 3 5 9 8 8 1 2 4



ORIGINAL FLAVOR

EFFERVESCENT PAIN RELIEF



Drug Facts (continued)

■ you are taking a diuretic ■ you have asthma ■ you have a sodium salt/acidifier dye
 ■ diabetes ■ gout ■ arthritis
 When using this product do not exceed recommended dosage
 Stop use and ask a doctor if you experience any of the following signs of stomach bleeding
 ■ feel faint ■ vomit blood ■ have bloody or black stools
 ■ have stomach pain that does not get better
 ■ feel sick for more than 2 days or more than 1 day if you are a child
 ■ feel weak or tired for more than 3 days
 ■ fever gets worse or lasts more than 3 days
 ■ new symptoms occur
 ■ ringing in the ears or a loss of hearing occurs
 If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children. In case of overdose, get medical help or contact a poison control center right away.

Directions

■ do not exceed recommended dosage
 ■ adults and children 12 years and over 2 tablets every 4 hours, or as directed by a doctor
 ■ do not exceed 8 tablets in 24 hours
 ■ 60 years and over 2 tablets every 4 hours, or as directed by a doctor
 ■ do not exceed 4 tablets in 24 hours
 ■ children under 12 years consult a doctor

Other Information

■ each tablet contains sodium 524 mg
 ■ store at room temperature (59°-86°F). Avoid excessive heat.

Inactive Ingredients

anhydrous citric acid, sodium bicarbonate

Questions or comments? 1-888-723-3929

DO NOT USE IF PRINTED PACKETS ARE TORN OR PUNCTURED

USES

Take this product any time – morning, noon or night – when you need relief from headaches, backaches, body aches or muscle pain.

MORNINGS NOONTIME EVENINGS



Place two (2) tablets in 4 oz. of water. Dissolve tablets completely.

You do not have to drink any of the residue that may be on the bottom of the glass. The medicines are in the water.

PK: 42188 SS CI
 R: 02/24

COMPLETE
 Also See "Active Ingredient"

NDC 21130-295-36



EFFERVESCENT PAIN RELIEF
 ORIGINAL FLAVOR

36 EFFERVESCENT TABLETS

BUFFERED ASPIRIN (NSAID)

- Pain Reliever/Fever Reducer
- Headache
 - Body Ache
 - Backache
 - Muscle Pain

DISTRIBUTED BY:
 BETTER LIVING BRANDS LLC
 P.O. BOX 99
 FLEMINGTON, NJ 08520-0099
 609-791-3200

*This product is not manufactured, distributed or licensed by the maker of Advil-Schwarz®



53748

RD 24156

3 21130 78184 4

4 2108866 1

Drug Facts

Active Ingredient

(In each effervescent tablet)
 Buffered aspirin 255 mg (NSAID), Pain reliever/fever reducer
 Water-soluble anti-inflammatory drug

Purpose
 ■ for the temporary relief of minor aches and pains due to:
 ■ headache ■ muscle pain ■ backache
 ■ minor arthritis pain ■ colds
 ■ temporarily reduces fever

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if you experience vomiting, persistent nausea, or other signs of Reye's syndrome, a rare but serious illness, stop use immediately and get medical help right away.

Stomach bleeding warning

This product contains an NSAID, which may cause severe stomach bleeding. The risk of stomach bleeding is higher if you are also taking aspirin, other NSAIDs, or blood thinners. You may be at a higher risk if you are taking aspirin, other NSAIDs, or blood thinners along with this product. Do not use for a longer time than directed.

Do not use if you are allergic to aspirin or any other pain reliever/fever reducer. If you have ever had an allergic reaction to this product or any of the ingredients, do not use.

Ask a doctor before use if you have a stomach bleeding warning sign by you, you have a history of stomach bleeding, you are taking aspirin, you have high blood pressure, heart disease, liver

SIGNATURE SELECT EFFERVESCENT PAIN RELIEF

aspirin tablet, effervescent

Product Information

Product Type

HUMAN OTC DRUG

Item Code (Source)

NDC:21130-295

Route of Administration ORAL

Active Ingredient/Active Moiety

| Ingredient Name | Basis of Strength | Strength |
|--|-------------------|----------|
| ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E) | ASPIRIN | 325 mg |

Inactive Ingredients

| Ingredient Name | Strength |
|--|----------|
| ANHYDROUS CITRIC ACID (UNII: XF417D3PSL) | |
| SODIUM BICARBONATE (UNII: 8MDF5V39QO) | |

Product Characteristics

| | | | |
|----------|-------|--------------|----------|
| Color | white | Score | no score |
| Shape | ROUND | Size | 25mm |
| Flavor | | Imprint Code | L421 |
| Contains | | | |

Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|---|------------------|---|----------------------|--------------------|
| 1 | NDC:21130-295-36 | 18 in 1 CARTON | 07/02/2024 | |
| 1 | | 2 in 1 POUCH; Type 0: Not a Combination Product | | |

Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|--------------------|--|----------------------|--------------------|
| OTC Monograph Drug | M013 | 07/02/2024 | |

Labeler - Safeway Inc. (009137209)

Revised: 3/2025

Safeway Inc.