

ACTICON PEDIATRIC- dextbrompheniramine maleate, pseudoephedrine hydrochloride solution
ACTIPHARMA, LLC

ACTICON® PEDIATRIC Cold & Allergy

Drug Facts

Active ingredients (in each 5 mL tsp)

Dextbrompheniramine Maleate, USP 1 mg

Pseudoephedrine HCl, USP 30 mg

Purpose

Antihistamine

Nasal Decongestant

Uses

- Temporarily relieves these symptoms due to the common cold, hay fever (allergic rhinitis) or other respiratory allergies:
- relieves sinus congestion and pressure, helps decongest sinus openings and passages
- restores freer breathing through the nose
- runny nose • sneezing • itching of the nose or throat
- itchy, watery eyes • nasal congestion

Warnings

Do not exceed recommended dosage

Do not use this product

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- a breathing problem such as emphysema or chronic bronchitis • glaucoma • heart disease • high blood pressure • thyroid disease • diabetes
- difficulty in urination due to enlargement of the prostate gland

Do not take this product if you are taking sedatives or tranquilizers, without first consulting your doctor.

When using this product

- excitability may occur, especially in children • may cause drowsiness • alcohol,

sedatives and tranquilizers may increase drowsiness effect

- avoid alcoholic beverages
- use caution when driving a motor vehicle or operating machinery

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- symptoms do not improve within 7 days or are accompanied by fever
- new symptoms occur

If pregnant or breast feeding, ask a health professional before use.

Keep out of reach of children. In case of accidental overdose, seek professional help or contact a Poison Control Center immediately.

Directions

Do not exceed more than 4 doses in 24 hours, or as directed by a doctor.

AGE	DOSE
Adults and children 12 years of age and over	2 teaspoonfuls (10 mL) every 4-6 hours
Children 6 to under 12 years of age	1 teaspoonful (5 mL) every 4-6 hours
Children under 6 years of age	ask a doctor

Other information

- Tamper Evident Feature: Do not use if inner seal is torn, cut, or opened.
- Store at controlled room temperature 15°- 30°C (59°-86°F).

Inactive ingredients

citric acid, flavor, methylparaben, potassium citrate, propylene glycol, propylparaben, purified water, sorbitol, sucralose.

Questions & Comments?

call weekdays from 8 AM to 4 PM AST at **1.787.608.0882.**

Contains the same active ingredients as Conex® Pediatric*

Alcohol FREE

Dye FREE

Sugar FREE

Cherry Flavor

ActiPharma

COMMITTED TO HEALTH AND WELL-BEING

WWW.ACTIPHARMA.NET

Manufactured in the USA with imported ingredients for ActiPharma, LLC. San Juan, PR

00917. *Conex[®] is a registered trademark of Llorens Pharmaceutical Corp. This product is not manufactured, distributed or marketed by Llorens Pharmaceutical Corp.

Packaging

NDC 63102-109-16

ACTICON[®]

PEDIATRIC

Cold & Allergy

Contains the same active ingredients as Conex[®] Pediatric*

ANTIHISTAMINE

NASAL DECONGESTANT

Alcohol FREE

Dye FREE

Sugar FREE

Cherry Flavor


ActiPharma
COMMITTED TO HEALTH AND WELL-BEING
WWW.ACTIPHARMA.NET

16 Fl.oz. (473 mL)

Drug Facts

Active ingredients
(in each 5 mL tsp)
Dexbrompheniramine Maleate, USP 1 mg.....Antihistamine
Pseudoephedrine HCl, USP 30 mg.....Nasal Decongestant

Purpose

Uses

- Temporarily relieves these symptoms due to the common cold, hay fever (allergic rhinitis) or other upper respiratory allergies:
- relieves sinus congestion and pressure, helps decongest sinus openings and passages
- restores freer breathing through the nose
- runny nose ■ sneezing ■ itching of the nose or throat
- itchy, watery eyes ■ nasal congestion

Warnings

Do not exceed recommended dosage

Do not use this product

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- a breathing problem such as emphysema or chronic bronchitis
- glaucoma ■ heart disease ■ high blood pressure ■ thyroid disease
- diabetes ■ difficulty in urination due to enlargement of the prostate gland

Do not take this product if you are taking sedatives or tranquilizers, without first consulting your doctor.

When using this product

- excitability may occur, especially in children ■ may cause drowsiness
- alcohol, sedatives and tranquilizers may increase drowsiness effect
- avoid alcoholic beverages ■ use caution when driving a motor vehicle or operating machinery

Drug Facts (continued)

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur ■ symptoms do not improve within 7 days or are accompanied by fever
- new symptoms occur

If pregnant or breast feeding, ask a health professional before use. Keep out of reach of children. In case of accidental overdose, seek professional help or contact a Poison Control Center immediately.

Directions

Do not exceed more than 4 doses in 24 hours, or as directed by a doctor.

AGE	DOSE
Adults and children 12 years of age and over	2 teaspoonfuls (10 mL) every 4-6 hours
Children 6 to under 12 years of age	1 teaspoonful (5 mL) every 4-6 hours
Children under 6 years of age	ask a doctor

Other information

- Tamper Evident Feature: Do not use if inner seal is torn, cut, or opened.
- Store at controlled room temperature 15° - 30°C (59° - 86°F).

Inactive ingredients citric acid, flavor, methylparaben, potassium citrate, propylene glycol, propylparaben, purified water, sorbitol, sucralose.

Questions & Comments? call weekdays from 8 AM to 4 PM AST at 1.787.608.0882.

Manufactured in the USA with imported ingredients for ActiPharma, LLC. San Juan, PR 00917. *Conex[®] is a registered trademark of Llorens Pharmaceutical Corp. This product is not manufactured, distributed or marketed by Llorens Pharmaceutical Corp.

Rev. 04/25


3 63102 10916 4

ACTICON PEDIATRIC

dexbrompheniramine maleate, pseudoephedrine hydrochloride solution

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:63102-109
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
DEXBROMPHENIRAMINE MALEATE (UNII: BPA9UT29BS) (DEXBROMPHENIRAMINE - UNII: 75T64B71RP)	DEXBROMPHENIRAMINE MALEATE	1 mg in 5 mL
PSEUDOEPHEDRINE HYDROCHLORIDE (UNII: 6V9V2RYJ8N) (PSEUDOEPHEDRINE - UNII: 7CUC9DDI9F)	PSEUDOEPHEDRINE HYDROCHLORIDE	30 mg in 5 mL

Inactive Ingredients

Ingredient Name	Strength
CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)	
METHYLPARABEN (UNII: A2I8C7HI9T)	
POTASSIUM CITRATE (UNII: EE90ONI6FF)	

PROPYLENE GLYCOL (UNII: 6DC9Q167V3)				
PROPYLPARABEN (UNII: Z8IX2SC1OH)				
WATER (UNII: 059QF0KO0R)				
SORBITOL (UNII: 506T60A25R)				
SUCRALOSE (UNII: 96K6UQ3ZD4)				
Product Characteristics				
Color			Score	
Shape			Size	
Flavor		CHERRY	Imprint Code	
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:63102-109-16	473 mL in 1 BOTTLE; Type 0: Not a Combination Product	07/17/2025	
Marketing Information				
Marketing Category		Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug		M012	07/17/2025	

Labeler - ACTIPHARMA, LLC (079340948)

Revised: 7/2025

ACTIPHARMA, LLC