

SUDAFED SINUS 12 HOUR PRESSURE PLUS PAIN- naproxen sodium and pseudoephedrine hydrochloride tablet, multilayer, extended release
Kenvue Brands LLC

Sudafed Sinus 12 Hour Pressure + Pain

Drug Facts

<i>Active ingredients (in each caplet)</i>	<i>Purposes</i>
Naproxen sodium 220 mg (naproxen 200 mg) (NSAID) *	Pain reliever/fever reducer
Pseudoephedrine HCl 120 mg, extended-release	Nasal decongestant

* nonsteroidal anti-inflammatory drug

Uses

temporarily relieves these cold, sinus, and flu symptoms:

- sinus pressure
- minor body aches and pains
- headache
- nasal and sinus congestion (promotes sinus drainage and restores freer breathing through the nose)
- fever

Warnings

Allergy alert

Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning

This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older

- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning

NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- in children under 12 years of age

Ask a doctor before use if

- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, thyroid disease, diabetes, have trouble urinating due to an enlarged prostate gland, or had a stroke
- you are taking a diuretic
- you have problems or serious side effects from taking pain relievers or fever reducers

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because naproxen may decrease this benefit of aspirin
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing

- weakness in one part or side of body
- slurred speech
- leg swelling
- redness or swelling is present in the painful area
- any new symptoms appear
- fever gets worse or lasts more than 3 days
- you have difficulty swallowing or the caplet feels stuck in your throat
- you get nervous, dizzy, or sleepless
- nasal congestion lasts more than 7 days

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use naproxen sodium at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- **do not take more than directed**
- **the smallest effective dose should be used**
- **swallow whole;** do not crush or chew
- **drink a full glass of water with each dose**

adults and children 12 years and older	<ul style="list-style-type: none"> ▪ 1 caplet every 12 hours ▪ do not take more than 2 caplets in 24 hours
children under 12 years	<ul style="list-style-type: none"> ▪ do not use

Other information

- **each caplet contains:** sodium 21 mg
- FDA approved dissolution specification differs from the USP dissolution specification
- store at 20-25°C (68-77°F)
- **do not use if blister unit is torn or broken**
- store in a dry place

Inactive ingredients

colloidal silicon dioxide, hypromellose, lactose monohydrate, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polysorbate 80, povidone, talc, titanium dioxide

Questions or comments?

call **1-888-217-2117** (toll-free) or **215-273-8755** (collect)

PRINCIPAL DISPLAY PANEL

NDC 50580-434-01

SUDAFED®

SINUS

12 HOUR PRESSURE + PAIN

Naproxen Sodium 220 mg (NSAID),
Pseudoephedrine Hydrochloride 120 mg
Extended Release Tablets,
Pain Reliever/Fever Reducer, Nasal Decongestant

12 HOUR MULTI-SYMPTOM RELIEF OF:

- SINUS PRESSURE • BODY ACHES
- NASAL CONGESTION • SINUS CONGESTION
- HEADACHE


actual size

16 CAPLETS*

*CAPSULE-SHAPED TABLETS

NON-DROWSY

1 CAPLET / 12 HOURS



The makers of the SUDAFED® family of products are dedicated to helping you and your family reduce head congestion and sinus pressure.

NDC 50580-434-01

SUDAFED®

SINUS

12 HOUR PRESSURE + PAIN

Naproxen Sodium 220 mg (NSAID),
Pseudoephedrine Hydrochloride 120 mg
Extended Release Tablets,
Pain Reliever/Fever Reducer, Nasal Decongestant



- 12 HOUR MULTI-SYMP TOM RELIEF OF:
- SINUS PRESSURE
 - BODY ACHES
 - NASAL CONGESTION
 - SINUS CONGESTION
 - HEADACHE

16 CAPLETS*
*CAPSULE-SHAPED TABLETS

NON-DROWSY

1 CAPLET / 12 HOURS

SUDAFED®

Distributed by:
JOHNSON & JOHNSON CONSUMER INC.
McNeil Consumer Healthcare Division
Fort Washington, PA 19034 USA
©J&JCI 2024
30056021



5F773MNC11

Questions or comments?

call 1-888-217-2117 (toll-free) or 215-273-8755 (collect)

Drug Facts (continued)

Other information ■ each caplet contains: sodium 21 mg ■ FDA approved dissolution specification differs from the USP dissolution specification ■ store at 20-25°C (68-77°F) ■ do not use if blister unit is torn or broken ■ store in a dry place

Inactive ingredients colloidal silicon dioxide, hypromellose, lactose monohydrate, magnesium stearate, microcrystalline cellulose, polyethylene glycol, poly sorbate 80, povidone, talc, titanium dioxide

Directions

■ do not take more than directed
■ swallow whole; do not crush or chew
■ drink a full glass of water with each dose
■ the smallest effective dose should be used

adults and children 12 years and older ■ 1 caplet every 12 hours ■ do not take more than 2 caplets in 24 hours
children under 12 years ■ do not use


Stop use and ask a doctor if
■ you experience any of the following signs of stomach bleeding:
■ feel faint ■ vomit blood ■ have bloody or black stools
■ you have symptoms of heart problems or stroke:
■ chest pain ■ trouble breathing ■ weakness in one part or side of body ■ slurred speech ■ leg swelling
■ redds or swelling is present in the painful area ■ any new symptoms appear
■ fever gets worse or lasts more than 3 days ■ you have difficulty swallowing or the caplet feels stuck in your throat
■ you get nervous, dizzy, or sleepless
■ nasal congestion lasts more than 7 days

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use naproxen sodium at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

When using this product ■ take with food or milk if stomach upset occurs
■ under a doctor's care for any serious condition ■ taking aspirin for heart attack or stroke, because naproxen may decrease this benefit of aspirin ■ taking any other drug

Ask a doctor or pharmacist before use if you are
■ you are taking a diuretic ■ you have problems or serious side effects from taking pain relievers or fever reducers



5F773MNC11



5F773MNC11

SUDAFED SINUS 12 HOUR PRESSURE PLUS PAIN

naproxen sodium and pseudoephedrine hydrochloride tablet, multilayer, extended release

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:50580-434
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
NAPROXEN SODIUM (UNII: 9TN87S3A3C) (NAPROXEN - UNII:57Y76R9ATQ)	NAPROXEN SODIUM	220 mg
PSEUDOEPHEDRINE HYDROCHLORIDE (UNII: 6V9V2RYJ8N) (PSEUDOEPHEDRINE - UNII:7CUC9DDI9F)	PSEUDOEPHEDRINE HYDROCHLORIDE	120 mg

Inactive Ingredients

Ingredient Name	Strength
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ05DW1A)	
POLYSORBATE 80 (UNII: 6OZP39ZG8H)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
TALC (UNII: 7SEV7J4R1U)	

Product Characteristics

Color	white	Score	no score
Shape	OVAL	Size	19mm
Flavor		Imprint Code	SUDAFED
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:50580-434-01	2 in 1 CARTON	06/17/2019	
1		8 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA076518	06/17/2019	

Labeler - Kenvue Brands LLC (118772437)

Revised: 11/2024

Kenvue Brands LLC