MIXED BERRY SORE THROAT AND COUGH LOZENGES- benzocaine lozenge HEB

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Benzocaine 7.5mg Dextromethorphan Hydrobromide 5mg

Active Ingredients (in each lozenge)

Benzocaine 7.5mg

Dextromethorphan Hydrobromide 5mg

Purpose

Oral Pain Reliever

Cough Suppressant

Uses temporarily relieves these symptoms:

occasiomal minor mouth irritation, sore mouth and sore throat

cough due to minor throat and bronchial irritation as may occur with the common cold

Warnings

Methemoglobinemia warning:

Use of this product may cause methemoglobinemia, a serious condition that must be treated promptly because it reduces the amount of oxygen carried in blood. This can occur even if you have used this product before. Stop use and seek immediate medical attention if you or a child in your care develops:

- pale, gray, or blue colored skin (cyanosis)
- headache
- rapid heart rate
- shortness of breath
- dizziness or lightheadedness
- fatigue or lack of energy

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, swelling, nausea or vomiting, consult a doctor promptly. If sore mouth symptoms do not improve in 7 days, or if irritation, pain, or redness persists or worsens, see your dentist or doctor promptly.

Allergy alert: Do not use this product if you have a history of allergy to local anesthetics such as procaine, butacaine, benzocaine, or other "caine" anesthetics.

Do not use is you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stoping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

- for teething
- in children under 2 years of age

Ask a doctor before use if you have:

- persistant or chronic cough such as occurs with smoking, asthma, or emphysema
- cough that occurs with too much phlegm (mucus)

Stop use and ask a doctor if

- sore mouth symptoms do not improve in 7 days
- irritation, pain, or redness persists or worsens
- swelling develops
- cough lasts for more than 7 days, comes back, or occurs with fever, rash, or persistent headache. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact Poison Control Center right away.

Directions

adults and children 12 years and older: allow 2 lozenges to dissolve slowly in mouth, one at a time; may be repeated every 4 hours, not to exceed 12 logenges in 24 hours, or as directed by a doctor

children 6 to 12 years of age: take 1 lozenge and allow it to dissolve slowly in the mouth; may be repeated every 4 hours, not to exceed 6 lozenges in 24 hours, or as directed by a doctor

children under 6 years of age; do not use

- store at room temperature
- protect from moisture

Inactive ingredients: Acesulfame potassium, corn starch, corn syrup, FD&C blue 1, FD&C red 40, glycerin, medium chain triglycerires, natural and artificial flavors, propylene glycol, soybean oil, sucrose, and water.

Questions? Or to report an adverse event call **1-800-245-2898**, Monday through Friday, 9AM - 4PM EST







inspect year proof(s) very curebody as we dischain horselfor any responsibility for errors or emissions. Our schedule will be edjusted to a later data by the number of days of approval delay.

Costoner represents and warrants to MFS that (q) Costoner within overs the copyrights or has obtained reproduction; and

(h) rights of any third party. Custoner should refer to the Standard Torms for additional terms governing its agreement with AIPS. AIPS reproduction of authorities requested by Custoner will not violate or furtings upon the
copyrights, publicity rights or other intrillactual property.

MPS PROOF COMPARED V12.2 BUILD 219 April 19, 2016 18:25:32 320123_J148880_MPS.pdf18:25:31April 19, 2016 sgenslak

Please enable overprint preview in Acrobat if you see an X in any of the below boxes. This will ensure that you are viewing your proof correctly.

When correctly rendered in compliance with PDFX standards no $oldsymbol{x}$ should appear in any b



MIXED BERRY SORE THROAT AND COUGH LOZENGES

benzocaine lozenge

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:37808-062
Route of Administration	ORAL		

Active Ingredient/Active Moiety				
Ingredient Name	Basis of Strength	Strength		
BENZOCAINE (UNII: U3RSY48JW5) (BENZOCAINE - UNII:U3RSY48JW5)	BENZOCAINE	7.5 mg		
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	5 mg		

Inactive Ingredients		
	Ingredient Name	Strength
WATER (UNII: 059QF0KO0R)		

Product Characteristics			
Color	red (Purplish Red)	Score	no score
Shape	ROUND	Size	17mm
Flavor	BERRY	Imprint Code	В
Contains			

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:37808-062- 18	18 in 1 CARTON; Type 0: Not a Combination Product	04/13/2016	



Marketing Information			
Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
part356	04/13/2016		
	Application Number or Monograph Citation	Application Number or Monograph Marketing Start Citation Date	

Labeler - HEB (007924756)

Registrant - Bestco Inc. (002149136)

Establishment				
Name	Address	ID/FEI	Business Operations	
Bestco Inc.		002149136	manufacture(37808-062)	

Revised: 10/2023 HEB