

# **TOPCARE BISMUTH STOMACH RELIEF- bismuth tablet, chewable**

## **TopCo**

-----  
**Bismuth subsalicylate 262 mg**

### **Active ingredient (in each chewable tablet)**

Bismuth subsalicylate 262 mg

### **Purpose**

Upset stomach reliever and antidiarrheal

### **Uses** relieves

- Traveler's diarrhea
- Diarrhea
- Upset stomach due to overindulgence in food or drink, including
- Heartburn
- Indigestion
- Nausea
- Gas
- Belching
- Fullness

### **Warnings**

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use the product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

### **Allergy alert: Contains salicylate. Do not take if you are**

- Allergic to salicylates (including aspirin)
- taking other salicylate products

### **Do not use if you have**

An ulcer

A bleeding problem

Bloody or black stool

### **Ask a doctor before use if you have**

- Fever
- Mucus in the stool

### **Ask a doctor or pharmacist before use if you are** taking any drug for

- Anticoagulation (thinning the blood)
- Diabetes
- Gout
- Arthritis

**When using this product** a temporary, but harmless, darkening of the stool and/or

tongue may occur

**Stop use and ask a doctor if**

- Symptoms get worse or last more than 2 days
- Ringing in the ears or loss of hearing occurs
- Diarrhea lasts more than 2 days

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

- Fully chew before swallowing
- Adults and children 12 years and over:
  - 2 chewable tablets every 1/2 hour or 4 chewable tablets every hour as needed for diarrhea
  - 2 chewable tablets every 1/2 hour to 1 hour as needed for overindulgence (upset stomach, heartburn, indigestion, nausea).
- Do not exceed 8 doses (16 chewable tablets) in 24 hours.
- Use until diarrhea stops but not more than 2 days
- Children under 12 years: ask a doctor.
- Drink plenty of clear fluids to help prevent dehydration caused by diarrhea.

**Inactive ingredients:** Beeswax, calcium carbonate, carmine, carnauba wax, citric acid, corn starch, corn syrup, dextrin, ethyl acetate, FD&C blue no 1 aluminum lake, FD&C blue no. 2 aluminum lake, FD&C red no. 40 aluminum lake, FD&C yellow no. 6 aluminum lake, glycerin, gum arabic, hydrogenated coconut oil, maltodextrin, medium chain triglycerides, natural and artificial flavors, phosphoric acid, propylene glycol, shellac, sodium benzoate, sorbic acid, sorbitol, soy lecithin, soybean oil, sucrose, titanium dioxide, vegetable oil (coconut and/or canola oil), water.

**Other information**

- **Each chewable tablet contains:** Calcium 50 mg
- Salicylate 123 mg
- Contains soy
- Store between 20° to 25°C (68° to 77°F) in a dry place. Avoid excessive heat (over 40°C).

**Questions?** Call **1-800-245-2989** Mon - Fri, 9AM - 4PM EST

NEVER EAT OR DRINK IF THE PRINTED SEAL ON BOTTLE IS BROKEN OR MISSING.  
DISTRIBUTED BY TOPCARE ASSOCIATES LLC  
DASCA, L 60143 ©TOPCO BOSTON  
QUESTIONS? 1-800-421-0139  
topcare@topco.com www.topcaremed.com

QUALITY GUARANTEED

2D

Scan here for more information or call 1-800-421-0139

TOPCARE ASSOCIATES LLC  
REGULAR STRENGTH  
**Stomach Relief**  
Chews  
BISMUTH SUBSALICATE 252 mg  
IPRED STOMACH RELIEVER • ANTI-DIARRHEAL

**SYMPTOM RELIEF**

- Upset stomach
- Indigestion
- Nausea
- Heartburn
- Diarrhea

24 CHEWABLE TABLETS

MINT FLAVOR

COMPARE TO PEPTO-BISMOL ACTIVE INGREDIENT\*

**Drug Facts**

**Active ingredient (in each chewable tablet)** Bismuth subsalicylate 252 mg

**Purpose** Upset stomach relief and anti-diarrheal

**Uses** relieves:
 

- Traveler's diarrhea
- Diarrhea
- Upset stomach due to overindulgence in food and drink, including:
  - Heartburn
  - Indigestion
- Nausea
- Gas
- Belching
- Fullness

**Warnings**  
**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

PN 60002096 PEEL BACK FOR DRUG FACTS

**Drug Facts (continued)**

**Allergy alert:** Contains salicylates. Do not take if you are:
 

- Allergic to salicylates (including aspirin)
- Taking other salicylate products

**Do not use if you have:**

- An ulcer
- A bleeding problem
- Blood or black stool

**Ask a doctor before use if you have:**

- Fever
- Mucus in the stool

**Ask a doctor or pharmacist before use if you are taking any drug for:**

- Anticoagulation (thinning the blood)
- Diabetes
- Gout
- Arthritis

**When using this product** a temporary, but harmless, darkening of the stool and/or tongue may occur.

**Stop use and ask a doctor if:**

- Symptoms get worse or last more than 2 days
- Ringing in the ears or loss of hearing occurs
- Diarrhea lasts more than 2 days

**For parent or babysitter:** ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Drug Facts (continued)**

**Directions**

- Fully chew before swallowing
- Adults and children 12 year and over:
  - 2 chewable tablets every 15 hour or 4 chewable tablets every hour as needed for diarrhea.
  - 2 chewable tablets every 15 hour to 1 hour as needed for overindulgence (upset stomach, heartburn, indigestion, nausea).
- Do not exceed 8 doses (16 chewable tablets) in 24 hours.
- Use until diarrhea stops but not more than 2 days.
- Children under 12 years: ask a doctor.
- Drink plenty of clear fluids to help prevent dehydration caused by diarrhea.

**Inactive ingredients** Sennoside, calcium carbonate, carnine, carnauba wax, citric acid, corn starch, croscarmellose, dextrin, ethyl acetate, FD&C blue no. 1 aluminum lake, FD&C blue no. 2 aluminum lake, FD&C red no. 40 aluminum lake, FD&C yellow no. 6 aluminum lake, glycerin, gum arabic, hydrogenated coconut oil, maltodextrin, medium chain triglycerides, natural and artificial flavors, phosphoric acid, propylene glycol, shellac, sodium benzoate, sorbic acid, sorbitol.

HINGE AREA - NO TEXT

**Drug Facts (continued)**

pey lecithin, soybean oil, sucrose, titanium dioxide, vegetable oil (coconut and/or canola oil), water.

**Other information**

- Each chewable tablet contains: Calcium 50 mg
- Salicylate 125 mg
- Contains pey
- Store between 20° to 25°C (68° to 77°F) in a dry place. Avoid excessive heat (over 40°C).

**Questions?** Call 1-800-240-2096 Mon - Fri 9AM - 4PM EST

This product is not manufactured or distributed by the owner of the registered trademark PeptoBismol®.

HINGE AREA - NO TEXT

bismuth tablet, chewable

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:76162-406
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
<b>BISMUTH SUBSALICYLATE</b> (UNII: 62TEY51RR1) (BISMUTH CATION - UNII:ZS9CD1I8YE)	BISMUTH SUBSALICYLATE	262 mg

### Inactive Ingredients

<b>Ingredient Name</b>	<b>Strength</b>
<b>SUCROSE</b> (UNII: C151H8M554)	

### Product Characteristics

<b>Color</b>	pink	<b>Score</b>	no score
<b>Shape</b>	ROUND	<b>Size</b>	19mm
<b>Flavor</b>	MINT	<b>Imprint Code</b>	SR
<b>Contains</b>			

### Packaging

<b>#</b>	<b>Item Code</b>	<b>Package Description</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
1	NDC:76162-406-24	24 in 1 BOTTLE; Type 0: Not a Combination Product	03/17/2025	



## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M008	03/17/2025	

**Labeler** - TopCo (006935977)

**Registrant** - SCNBestco (002149136)

## Establishment

Name	Address	ID/FEI	Business Operations
Betco		002149136	manufacture(76162-406)

