

MAXIMUM STRENGTH ANTACID AND ANTIGAS- aluminum hydroxide, magnesium hydroxide, simethicone suspension
Meijer, Inc.

mei antacid max cherry

Active ingredients (in each 10 mL teaspoonful)

Aluminum hydroxide (equivalent to dried gel, USP) 800 mg

Magnesium hydroxide 800 mg

Simethicone 80 mg

Purposes

Antacid

Antigas

Uses

relieves

- heartburn
- sour stomach
- acid indigestion
- the symptoms referred to as gas

Warnings

Ask a doctor before use if you have

- kidney disease
- a magnesium-restricted diet

Ask a doctor or pharmacist before use if you are presently taking a prescription drug.

Antacids may interact with certain prescription drugs.

Stop use and ask a doctor if symptoms last more than 2 weeks

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Directions

- shake well before use
- adults and children 12 years and older: take 10 mL to 20 mL (1 to 2 doses) between

meals, at bedtime, or as directed by a doctor

- do not take more than 60 mL (6 doses) in 24 hours
- do not use the maximum dosage for more than 2 weeks
- children under 12 years of age: ask a doctor
- measure with dosing cup provided
- mL = milliliter

Other information

- **each 10 mL teaspoonful contains:** magnesium 340 mg, sodium 10 mg
- protect from freezing
- store at room temperature tightly closed

Inactive ingredients

benzyl alcohol, butylparaben, carboxymethylcellulose sodium, flavor, hypromellose, microcrystalline cellulose, propylparaben, purified water, saccharin sodium, sorbitol solution

Questions or Comments?

1-800-540-3765

package label

meijer.

NDC 79481-0638-1

MAXIMUM STRENGTH
**Antacid &
Antigas**

RELIEVES HEARTBURN, SOUR STOMACH
ACID INDIGESTION, PRESSURE AND BLOATING

CHERRY

Alumina, Magnesia, and
Simethicone Oral Suspension



12 fl oz (355 mL)

225-11322-1
REV 0825

Drug Facts

**TAMPER-EVIDENT: DO NOT USE IF
IMPRINTED NECKBAND IS MISSING
OR BROKEN.**

**Active ingredients
(in each 10 mL dose)**

Aluminum hydroxide 800 mgAntacid
(equivalent to dried gel USP)
Magnesium hydroxide 800 mgAntacid
Simethicone 80 mgAntigas

Purposes

Uses relieves • heartburn • sour stomach
• acid indigestion • the symptoms referred to as gas

Warnings

Ask a doctor before use if you have

• kidney disease • a magnesium-restricted diet

Ask a doctor or pharmacist before use if you are presently
taking a prescription drug. Antacids may interact with certain
prescription drugs.

Stop use and ask a doctor if symptoms last more than 2 weeks

If pregnant or breast-feeding, ask a health professional
before use.

Keep out of reach of children.

Directions

• shake well before each use
• adults and children 12 years and older: take 10 mL to 20 mL
(1 to 2 doses) between meals, at bedtime, or as directed
by a doctor
• do not take more than 60 mL (6 doses) in 24 hours
• do not use the maximum dosage for more than 2 weeks
• children under 12 years of age; ask a doctor
• measure with dosing cup provided • mL = milliliter

Other information

• each 10 mL dose contains:
magnesium 340 mg, sodium 10 mg • protect from freezing
• store at room temperature tightly closed

Inactive ingredients

benzyl alcohol, butylparaben,
carboxymethylcellulose sodium, flavor, hypromellose,
microcrystalline cellulose, propylparaben, purified water,
saccharin sodium, sorbitol solution

Questions or comments? 1-800-540-3765

DIST. BY MEIJER DISTRIBUTION, INC. PID 3103042
GRAND RAPIDS, MI 49544
www.meijer.com



925-11322-1
REV 6C0725

MAXIMUM STRENGTH ANTACID AND ANTIGAS

aluminum hydroxide, magnesium hydroxide, simethicone suspension

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:79481-0638
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ALUMINUM HYDROXIDE (UNII: 5QB0T2IUN0) (ALUMINUM HYDROXIDE - UNII:5QB0T2IUN0)	ALUMINUM HYDROXIDE	800 mg in 10 mL
MAGNESIUM HYDROXIDE (UNII: NBZ3QY004S) (MAGNESIUM CATION - UNII:T6V3LHY838, HYDROXIDE ION - UNII:9159UV381P)	MAGNESIUM HYDROXIDE	800 mg in 10 mL
DIMETHICONE (UNII: 92RU3N3Y1O) (DIMETHICONE - UNII:92RU3N3Y1O)	DIMETHICONE	80 mg in 10 mL

Inactive Ingredients				
Ingredient Name				Strength
BENZYL ALCOHOL (UNII: LKG8494WBH)				
BUTYLPARABEN (UNII: 3QPI1U3FV8)				
CARBOXYMETHYLCELLULOSE SODIUM (UNII: K679OBS311)				
HYPROMELLOSES (UNII: 3NXW29V3WO)				
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)				
PROPYLPARABEN (UNII: Z8IX2SC1OH)				
WATER (UNII: 059QF0KO0R)				
SACCHARIN SODIUM (UNII: SB8ZUX40TY)				
SORBITOL SOLUTION (UNII: 8KW3E207O2)				

Product Characteristics			
Color		Score	
Shape		Size	
Flavor	CHERRY	Imprint Code	
Contains			

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:79481-0638-1	355 mL in 1 BOTTLE; Type 0: Not a Combination Product	08/01/2025	

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M001	08/01/2025	

Labeler - Meijer, Inc. (006959555)

Registrant - GCP Laboratories (965480861)

Establishment			
Name	Address	ID/FEI	Business Operations
GCP Laboratories		965480861	manufacture(79481-0638)