

THERABLOAT DRENCH- poloxalene concentrate
Zoetis Inc.

Therabloat®

Therabloat®
(poloxalene)

Drench Concentrate

For treatment of legume (alfalfa, clover) bloat in cattle

For veterinary use only

2 fl oz (59.1 mL)

Approved by FDA under NADA # 039-729

Active Ingredient: poloxalene (each fl oz contains 25 grams)

Inactive Ingredients: methylparaben, propylparaben, propylene glycol

Indications

Therabloat Drench Concentrate will relieve legume bloat in cattle within minutes when used as directed. For best results administer at the earliest sign of bloat. If bloat has progressed in its severity to the degree that the animal is down, other means of treatment also are recommended (rumenotomy-rumen puncture). For oral use only— not for injection. May be used in lactating animals.

Dosage

For animals up to 500 lb, use 1 fl oz of Drench Concentrate. For animals over 500 lb, use 2 fl oz of Drench Concentrate.

Directions for Preparing Drench

Add the proper amount of concentrate to 1 pint of water. Mix well and administer using a drenching bottle. If a stomach tube is to be used, add concentrate to 1 gallon of water.

Caution

Keep out of reach of children.

Storage

Store at 20° to 25°C (68° to 77°F), with excursions permitted between 15° and 37°C (59° and 99°F).

For Animal Use Only

Not for Human Use

zoetis

Distributed by:
Zoetis Inc.
Kalamazoo, MI 49007

40039268

PRINCIPAL DISPLAY PANEL - 2 fl oz (59.1 mL) label

Therabloat®
(poloxalene)
Drench Concentrate
For treatment of legume (alfalfa, clover) bloat in cattle
For veterinary use only
2 fl oz (59.1 mL)
Approved by FDA under NADA # 039-729

zoetis

Active Ingredient: poloxalene (each fl oz contains 25 grams)
Inactive Ingredients: methylparaben, propylparaben, propylene glycol
Indications: Therabloat Drench Concentrate will relieve legume bloat in cattle within minutes when used as directed. For best results administer at the earliest sign of bloat. If bloat has progressed in its severity to the degree that the animal is down, other means of treatment also are recommended (rumenotomy-rumen puncture). For oral use only—not for injection. May be used in lactating animals.
Dosage: For animals up to 500 lb, use 1 fl oz of Drench Concentrate. For animals over 500 lb, use 2 fl oz of Drench Concentrate.
Directions for Preparing Drench: Add the proper amount of concentrate to 1 pint of water. Mix well and administer using a drenching bottle. If a stomach tube is to be used, add concentrate to 1 gallon of water.
Caution: Keep out of reach of children.
Store at 20° to 25°C (68° to 77°F), with excursions permitted between 15° and 37°C (59° and 99°F).
For Animal Use Only
Not for Human Use

zoetis
Distributed by:
Zoetis Inc.,
Kalamazoo, MI 49007

40051846

THERABLOAT DRENCH			
poloxalene concentrate			
Product Information			
Product Type	OTC ANIMAL DRUG	Item Code (Source)	NDC:54771-8620
Route of Administration	ORAL		
Active Ingredient/Active Moiety			
Ingredient Name		Basis of Strength	Strength

POLOXALENE (UNII: V8B3K56SW0) (POLOXALENE - UNII:V8B3K56SW0) | POLOXALENE | 25 g in 30 mL

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:54771-8620-1	59.1 mL in 1 BOTTLE		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NADA	NADA039729	05/12/1970	

Labeler - Zoetis Inc. (828851555)

Revised: 5/2025

Zoetis Inc.