

**REVASOL PEPTIC RELIEF- bismuth subsalicylate liquid**  
**Rnv LLC**

-----  
**REVASOL PEPTIC RELIEF**

**Active ingredient (in each 30 mL dose) Purpose**

Bismuth subsalicylate 525 mg.....Upset stomach reliever and antidiarrheal  
Upset stomach reliever and antidiarrheal

**Uses**

relieves • travelers' diarrhea • diarrhea • upset stomach due to overindulgence in food and drink, including:  
• heartburn • indigestion • nausea • gas • belching • fullness

**Warnings**

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert: Contains salicylate. Do not take if you are**

• allergic to salicylates (including aspirin) • taking other salicylate products

**Do not use if you have** • an ulcer • a bleeding problem • bloody or black stool

**Ask a doctor before use if you have** • fever • mucus in the stool

**Ask a doctor or pharmacist before use** if you are taking any drug for

• anticoagulation (thinning the blood) • diabetes • gout • arthritis

**When using this product** a temporary, but harmless, darkening of the stool and/or tongue may occur

**Stop use and ask a doctor if** • symptoms get worse or last more than 2 days

• ringing in the ears or loss of hearing occurs • diarrhea lasts more than 2 days

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of the reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

• shake well before use • only with dosing cup provided

• **adults and children 12 years and over:**

• 30 mL (1 dose) every 1/2 hour or 60 mL (2 doses) every hour as needed for diarrhea/traveler's diarrhea

• 30 mL (1 dose) every 1/2 hour as needed for overindulgence (upset stomach, heartburn, indigestion, nausea)

• do not exceed 8 doses (240 mL) in 24 hours

- use until diarrhea stops but no more than 2 days
- **children under 12 years:** ask a doctor
- drink plenty of clear fluids to help prevent dehydration caused by diarrhea

### Other information

- **each 30 mL dose contains:** sodium 22 mg
- salicylate 249 mg • protect from freezing • avoid excessive heat (over 104°F or 40°C)
- **TAMPER-EVIDENT FEATURE:** Do not use if printed safety seal is torn, broken or missing.

### Inactive ingredients

anhydrous citric acid, benzoic acid, D&C Red #22, D&C Red #28, flavor, purified water, salicylic acid, sodium citrate, sodium salicylate, sorbic acid, sucralose, xanthan gum.

### Questions or comments 1-888-446-4753

Monday - Friday (9 AM - 5 PM EST)

**Drug Facts**

**Active ingredient (in each 30 mL dose)** Bismuth subsalicylate 525 mg..... **Purpose** Upset stomach reliever and antidiarrheal

**Uses** relieves • travelers' diarrhea • diarrhea • upset stomach due to overindulgence in food and drink, including:  
• heartburn • indigestion • nausea • gas • belching • fullness

**Warnings**  
**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert: Contains salicylate. Do not take if you are**  
• allergic to salicylates (including aspirin) • taking other salicylate products

**Do not use if you have** • an ulcer • a bleeding problem • bloody or black stool

**Ask a doctor before use if you have** • fever • mucus in the stool

**Ask a doctor or pharmacist before use if you are** taking any drug for  
• anticoagulation (thinning the blood) • diabetes • gout • arthritis

**When using this product** a temporary, but harmless, darkening of the stool and/or tongue may occur

**Stop use and ask a doctor if** • symptoms get worse or last more than 2 days • ringing in the ears or loss of hearing occurs • diarrhea lasts more than 2 days

**If pregnant or breast-feeding,** ask a health professional before use.  
**Keep out of the reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

\*This product is not manufactured or distributed by The Procter and Gamble Company. Peto-Bismol® is a registered trademark of The Procter and Gamble Company.

**Drug Facts (continued)**

**Directions** • shake well before use • only with dosing cup provided  
• **adults and children 12 years and over:**  
• 30 mL (1 dose) every 1/2 hour or 60 mL (2 doses) every hour as needed for diarrhea/traveler's diarrhea  
• 30 mL (1 dose) every 1/2 hour as needed for overindulgence (upset stomach, heartburn, indigestion, nausea)  
• do not exceed 8 doses (240 mL) in 24 hours  
• use until diarrhea stops but no more than 2 days  
• **children under 12 years:** ask a doctor  
• drink plenty of clear fluids to help prevent dehydration caused by diarrhea

**Other information**  
• each 30 mL dose contains: sodium 22 mg  
• salicylate 249 mg • protect from freezing • avoid excessive heat (over 104°F or 40°C)  
• **TAMPER-EVIDENT FEATURE:** Do not use if printed safety seal is torn, broken or missing.

**Inactive ingredients** anhydrous citric acid, benzoic acid, D&C Red #22, D&C Red #28, flavor, purified water, salicylic acid, sodium citrate, sodium salicylate, sorbic acid, sucralose, xanthan gum.

**Questions or comments 1-888-446-4753**  
Monday - Friday (9 AM - 5 PM EST)

**Made in USA**  
Distributed by:  
RNV LLC  
14501 NW 60 Ave,  
Miami Lakes, FL 33014  
Rev. 1

8 60013 43590 5  
Lot:  
Exp:

## REVASOL PEPTIC RELIEF

bismuth subsalicylate liquid

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:84379-283
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BISMUTH SUBSALICYLATE (UNII: 62TEY51RR1) (SALICYLIC ACID - UNII:O414PZ4LPZ,	BISMUTH	525 mg

BISMUTH CATION - UNII:ZS9CD1I8YE) SUBSALICYLATE in 30 mL

### Inactive Ingredients

Ingredient Name	Strength
<b>ANHYDROUS CITRIC ACID</b> (UNII: XF417D3PSL)	
<b>SALICYLIC ACID</b> (UNII: O414PZ4LPZ)	
<b>XANTHAN GUM</b> (UNII: TTV12P4NEE)	
<b>METHYL SALICYLATE</b> (UNII: LAV5U5022Y)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>SODIUM SALICYLATE</b> (UNII: WQ1H85SYP)	
<b>SUCRALOSE</b> (UNII: 96K6UQ3ZD4)	
<b>SODIUM CITRATE</b> (UNII: 1Q73Q2JULR)	
<b>SORBIC ACID</b> (UNII: X045WJ989B)	
<b>D&amp;C RED NO. 22</b> (UNII: 1678RKX8RT)	
<b>D&amp;C RED NO. 28</b> (UNII: 767IP0Y5NH)	
<b>BENZOIC ACID</b> (UNII: 8SKN0B0MIM)	

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:84379-283-04	118 mL in 1 BOTTLE; Type 0: Not a Combination Product	02/03/2025	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M008	02/03/2025	

**Labeler** - Rnv LLC (118917568)

**Registrant** - Rnv LLC (118917568)

### Establishment

Name	Address	ID/FEI	Business Operations
Rnv LLC		118917568	manufacture(84379-283)

Revised: 3/2025

Rnv LLC