

GREEN GUARD MAXIMUM NON-ASPIRIN- acetaminophen tablet, film coated
Unifirst First Aid Corporation

Green Guard® Maximum Strength Non-Aspirin

Drug Facts

Active ingredient (in each tablet)

Acetaminophen 500 mg

Purpose

Pain reliever/fever reducer

Uses

temporarily relieves minor aches and pains due to:

- headache
- muscular aches
- minor pain of arthritis
- backache
- the common cold
- toothache
- premenstrual and menstrual cramps

temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if you have

- liver disease

Ask a doctor or pharmacist before use if

- you are taking the blood thinning drug warfarin

Stop use and ask a doctor if

- pain gets worse or lasts more than for more than 10 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose Warning:

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not use more than directed (see overdose warning)**

Adults and children: (12 years and over) ■ take 2 tablets every 6 hours while symptoms last.

■ do not take more than 6 tablets in 24 hours, unless directed by a doctor

■ do not use for more than 10 days unless directed by a doctor

Children under 12 years: Ask a doctor

Other information

- store at room temperature 59°-86°F (15°-30°C)
- tamper-evident sealed packets
- do not use any opened or torn packets

Inactive ingredients

corn starch*, hypromellose*, polyethylene glycol*, povidone (K-30)*, pregelatinized starch*, purified water*, sodium starch glycolate*, stearic acid*, titanium dioxide*

* may contain

Questions or comments? 1-800-869-6970

Green Guard Max Non-Aspirin Label

Maximum Strength Non-Aspirin relief™

Acetaminophen 500 mg/Acetaminofeno 500 mg

Pain Reliever/Fever Reducer

Alivia el Dolor/Reduce la Fiebre

Compare Active Ingredient To Extra Strength Tylenol®

Green Guard®

Pull Out

Tire Para Abrir

Tylenol® is a Registered Trademark of McNeil Consumer Products

50 Packets of 2 Tablets

100 Tablets

Order #2224

Distributed by Green Guard® St Louis. MO 63045

GREEN GUARD
Pain Reliever/Fever Reducer
Alivia el Dolor/Reduce la Fiebre
Acetaminophen 500 mg
Non-Aspirin
MAXIMUM STRENGTH
100 Tablets

Información del medicamento (continuación)
Dijo de tomarlo y consulte al médico si:
■ el dolor empeora y perdura más de 10 días
■ la fiebre empeora o perdura más de 3 días
■ aparecen nuevos síntomas
■ aparece hinchazón o enrojecimiento
■ Esto pueden ser señales de una enfermedad grave.
Si está usted embarazada o en periodo de lactancia, consulte la opinión de un médico antes de consumir este producto.
Mantenga este y todos los medicamentos fuera del alcance de los niños.
Advertencia de sobredosis: En caso de ocurrir una sobredosis, comuníquese de inmediato con un médico o llame a un centro de control de envenenamiento (1-800-222-1222). La información de advertencia es crítica tanto para los adultos como para los niños sin importar la edad ni los síntomas.
Instrucciones
■ No consuma más de lo que se indica (ver advertencia de sobredosis).
Adultos y niños (de 12 años y mayores):
■ Tome 2 tabletas cada 6 horas mientras los síntomas perduren
■ no tomar más de 6 tabletas en 24 horas, a menos que lo indique el médico.
■ no administrar por más de 10 días a menos que lo indique el médico.
Niños menores de 12 años:
■ consultar al médico.
Otra información
■ almacenamiento: conservarse a temperatura ambiente 59°-86°F (15°-30°C)
■ empaquetado con sellado evidente.
■ no use ningún plástico adherido o otro.
Ingredientes inactivos
amidon de maíz*, hidroximetilcelulosa polietileno*, povidona K30®, almidón pregelatinizado*, agua purificada*, ácido glicólico sódico, ácido estearico, dióxido de titanio*
*fuente: conifer.
¿Tiene alguna pregunta o comentario? Llame al 1-800-989-6370.
Rev. 11/62 Mtg. 1-2A-H

Drug Facts
Active ingredient (in each tablet) Purpose
acetaminophen 500 mg Pain reliever/fever reducer
Uses
temporarily relieves minor aches and pains due to:
■ headache ■ muscle aches ■ minor pain of arthritis
■ backache ■ the common cold ■ toothache
■ rheumatism and menstrual cramps
temporarily reduces fever.
Warnings
Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:
■ more than 4,000 mg in 24 hours, which is the maximum daily amount.
■ with other drugs containing acetaminophen.
■ for more alcoholic drinks every day while using this product.
Other warnings: Acetaminophen may cause severe skin reactions. Symptoms may include:
■ skin rash/ hives ■ blisters ■ rash
If a skin reaction occurs, stop use and seek medical help right away.
Do not use
■ with any other drug containing acetaminophen (prescription or over-the-counter). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
■ if you are allergic to acetaminophen or any of the inactive ingredients in this product.
Ask a doctor before use if you have ■ liver disease
■ you are taking the blood thinning drug warfarin
Stop use and ask a doctor if
■ liver gets worse or lasts for more than 10 days
■ new symptoms occur
■ redness or swelling is present
These could be signs of a serious condition.
If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children.

Información del medicamento
ingrediente activo (en cada tableta) Propiedad
acetaminofeno 500 mg Alivia el dolor/Reduce la fiebre
Usos
para aliviar temporalmente las fiebres y molestias asociadas con:
■ dolor de cabeza ■ dolores musculares
■ dolores de las articulaciones ■ dolor de espalda
■ resaca ■ dolor de dientes
■ dolores premenstruales y menstruales
Advertencias
Advertencia con respecto al hígado: Este producto contiene acetaminofeno. Daño grave al hígado puede ocurrir si usted consume:
■ más de 4,000 mg en 24 horas, lo cual es la máxima dosis recomendada.
■ este producto con cualquier otro medicamento que contenga acetaminofeno.
■ 3 o más bebidas alcohólicas diariamente mientras toma este producto.
Información del medicamento (continuación)
Alerta a las alergias: El acetaminofeno puede causar severas reacciones alérgicas en la piel. Las cuales pueden incluir:
■ el enrojecimiento de la piel ■ ampollas ■ erupción.
Si se presenta una reacción alérgica en la piel, suspenda su consumo de inmediato y obtenga atención médica.
No se administre
■ con ningún otro producto que contenga acetaminofeno (sobre-dosis o sin receta) si no está seguro si su medicamento contiene acetaminofeno consulte con un médico o farmacéutico.
■ si usted es alérgico al acetaminofeno o a cualquiera de los ingredientes inactivos en este producto.
Consulte a un médico antes de administrarle al usted (niño) ■ la enfermedad del hígado.
Consulte con un médico o farmacéutico antes de administrarlo si:
■ toma el medicamento warfarin para el adelgazamiento de la sangre.
(continúa en la página inferior de la caja)
CONSERVE LA CAJILLA PARA OBTENER LA INFORMACIÓN COMPLETA DEL PRODUCTO

Drug Facts (continued)
Directions
Do not use more than directed (see overdose warning).
Adults and Children:
■ take 2 tablets every 6 hours while symptoms last.
■ do not take more than 6 tablets in 24 hours unless directed by a doctor.
■ do not use for more than 10 days unless directed by a doctor.
Children under 12 years:
■ Ask a doctor.
Drug Facts (continued)
Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.
Other information
■ store at room temperature 59°-86°F (15°-30°C)
■ use paper-lined sealed packets
■ do not use any opened or torn packets
Inactive ingredients
corn starch*, hydroxyethylcellulose*, polyethylene glycol*, povidone (K-30)*, pregelatinized starch*, purified water*, sodium starch glycolate, stearic acid, titanium dioxide*
*may contain
Questions or comments?
1-800-989-6370

GREEN GUARD
Pain Reliever/Fever Reducer
Alivia el Dolor/Reduce la Fiebre
Acetaminophen 500 mg/Acetaminofeno 500 mg
50 Packets of 2 Tablets
100 Tablets
Order #2224
Distributed by Green Guard®
St Louis, MO 63045

PMS 528
PMS 286

RETAIN CARTON FOR COMPLETE PRODUCT INFORMATION

GREEN GUARD MAXIMUM NON-ASPIRIN

acetaminophen tablet, film coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:47682-009
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	500 mg

Inactive Ingredients

Ingredient Name	Strength
HYPROMELLOSES (UNII: 3NXW29V3WO)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ05DW1A)	
POVIDONE (UNII: FZ989GH94E)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	
STARCH, CORN (UNII: O8232NY3SJ)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	white (white)	Score	no score
Shape	ROUND (ROUND)	Size	12mm
Flavor		Imprint Code	AZ;235
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:47682-009-33	50 in 1 BOX	12/30/2008	
1	NDC:47682-009-99	2 in 1 PACKET; Type 0: Not a Combination Product		
2	NDC:47682-009-48	125 in 1 BOX	12/30/2008	
2	NDC:47682-009-99	2 in 1 PACKET; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	12/30/2008	

Labeler - Unifirst First Aid Corporation (832947092)

Revised: 2/2024

Unifirst First Aid Corporation