

**TOPCARE NIGHTTIME SEVERE COLD AND COUGH HONEY LEMON INFUSED WITH CHAMOMILE AND WHITE TEA FLAVORS- acetaminophen, diphenhydramine hcl and phenylephrine hcl granule, for solution**  
**TOP CARE (Topco Associates LLC)**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

**Topcare Nighttime Severe Cold and Cough Honey lemon infused with Chamomile and White tea flavors**

Active ingredients (in each packet)	Purpose
Acetaminophen 650 mg .....	Pain reliever/Fever reducer
Diphenhydramine hydrochloride 25 mg .....	Antihistamine/Cough suppressant
Phenylephrine hydrochloride 10 mg.....	Nasal decongestant.

**Uses**

- temporarily relieves:
  - minor aches and pains - headache - minor sore throat pain
  - nasal and sinus congestion - cough due to minor throat and bronchial irritation
  - runny nose - sneezing - itchy, watery eyes due to hay fever
  - itchy nose and throat
- temporarily reduces fever.

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take - more than 6 packets in 24 hours, which is the maximum daily amount - with other drugs containing acetaminophen - 3 or more alcoholic drinks every day while using this product.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use**

- with any other drug containing acetaminophen (prescription and nonprescription). Ask a doctor or pharmacist before using with other drugs if you are not sure.
- with any other product containing diphenhydramine, even one used on the skin
- If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains a MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have**

- liver disease - heart disease - high blood pressure - thyroid disease - glaucoma - diabetes - trouble urinating due to an enlarged prostate gland - a breathing problem such as emphysema, asthma, or chronic bronchitis - cough that occurs with too much phlegm (mucus) - cough that lasts or is chronic such as occurs with smoking, asthma or emphysema.

**Ask a doctor or pharmacist before use if you are**

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin.

**When using this product**

- do not exceed recommended dosage
- avoid alcoholic drinks
- marked drowsiness may occur
- alcohol, sedatives and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery
- excitability may occur, especially in children.

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occurs
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- symptoms do not get better or worsen
- pain, cough or nasal congestion gets worse or lasts more than 7 days
- cough comes back or occurs with fever, rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health care professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

#### Directions

- do not use more than directed
- take every 4 hours; not to exceed 6 packets in 24 hours or as directed by a doctor.
- adults and children 12 years of age and over: dissolve contents of one packet into 8 oz. hot water; sip while hot. Consume entire drink within 10-15 minutes.
- children under 12 years of age: consult a doctor.
- if using a microwave, add contents of one packet to 8 oz. of cool water: stir briskly before and after heating. Do not overheat.

#### Other information

- each packet contains: potassium 7 mg, sodium 46 mg
- phenylketonurics: contains phenylalanine 13 mg per packet
- store at controlled room temperature 20-25C (68-77F). Protect from excessive heat and moisture.

Acesulfame K, Aspartame, Caramel Color, Citric Acid, D and C Yellow 10, Maltodextrin, Natural Flavors, Sodium Citrate, Starch and Sugar.

**Drug Facts**

**Active ingredients (in each packet)** **Purpose**

Acetaminophen 650 mg ..... Pain reliever/Fever reducer

Diphenhydramine hydrochloride 25 mg ..... Antihistamine/Cough suppressant

Phenylephrine hydrochloride 10 mg ..... Nasal decongestant

**Uses**

- Temporarily relieves:
  - minor aches and pains
  - headache
  - minor sore throat pain
  - nasal and sinus congestion
  - cough due to minor throat and bronchial irritation
  - runny nose
  - sneezing
  - itchy, watery eyes due to hay fever
  - itchy nose and throat
- Temporarily reduces fever

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take more than 6 packets in 24 hours, which is the maximum daily amount, with other drugs containing acetaminophen, or 3 or more alcoholic drinks every day while using this product. **Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use**

- with any other drug containing acetaminophen (prescription and nonprescription).
- Ask a doctor or pharmacist before using with other drugs if you are not sure.
- with any other product containing diphenhydramine, even one used on the skin.
- If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains a MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have**

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- glaucoma
- diabetes
- trouble urinating due to an enlarged prostate gland
- a breathing problem such as emphysema, asthma, or chronic bronchitis
- cough that occurs with too much phlegm (mucus)
- cough that lasts or is chronic such as occurs with smoking, asthma or emphysema.

**Ask a doctor or pharmacist before use if you are**

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

**When using this product**

- do not exceed recommended dosage
- avoid alcoholic drinks
- marked drowsiness may occur
- alcohol, sedatives and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery
- excitability may occur, especially in children.

**Stop use and ask a doctor if**

- nervousness, dizziness, or sleeplessness occurs
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- symptoms do not get better or worsen
- pain, cough or nasal congestion gets worse or lasts more than 7 days
- cough comes back or occurs with fever, rash or headache that lasts. These could be signs of a serious condition.

**Drug Facts (continued)**

**If pregnant or breast-feeding,** ask a health care professional before use.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

- take every 4 hours; not to exceed 6 packets in 24 hours or as directed by a doctor.
- adults and children 12 years of age and over: dissolve contents of one packet into 8 oz. hot water; sip while hot. Consume entire drink within 10-15 minutes.
- children under 12 years of age: consult a doctor.
- if using a microwave, add contents of one packet to 8 oz. of cool water; stir briskly before and after heating. Do not overheat.

**Other information**

- each packet contains: potassium 7 mg, sodium 46 mg
- phenylketonurics: contains phenylalanine 13 mg per packet
- store at controlled room temperature 20-25°C (68-77°F). Protect from excessive heat and moisture.

**Inactive ingredients:** Acesulfame K, aspartame, caramel color, citric acid, D&C Yellow 10, maltodextrin, natural flavors, sodium citrate, starch and sugar.

**DO NOT USE IF SEALED PACKET IS TORN OR BROKEN.**

DISTRIBUTED BY TOPCO ASSOCIATES LLC  
ELK GROVE VILLAGE, IL 60007  
1-888-423-0139 ©TOPCO QHP313  
topcare@topco.com  
MADE IN CANADA

**TopCare®**

FOR NIGHTTIME RELIEF

**Nighttime Severe Cold & Cough**

ACETAMINOPHEN  
Pain Reliever • Fever Reducer  
DIPHENHYDRAMINE HCl  
Antihistamine • Cough Suppressant  
PHENYLEPHRINE HCl  
Nasal Decongestant

For relief of

- Nasal Congestion • Cough
- Runny Nose • Sneezing
- Body Ache
- Sore Throat Pain
- Headache • Fever

Honey Lemon Infused with Chamomile and White Tea flavors



\*COMPARE TO THERAFLU® NIGHTTIME SEVERE COLD & COUGH

See New Warnings Information

**6 PACKETS**

**TopCare®**

FOR NIGHTTIME RELIEF

**Nighttime Severe Cold & Cough**

ACETAMINOPHEN  
Pain Reliever • Fever Reducer  
DIPHENHYDRAMINE HCl  
Antihistamine  
Cough Suppressant  
PHENYLEPHRINE HCl  
Nasal Decongestant

For relief of

- Nasal Congestion • Cough
- Runny Nose • Sneezing
- Body Ache
- Sore Throat Pain
- Headache • Fever

Honey Lemon Infused with Chamomile and White Tea flavors



**6 PACKETS**

\*This product is not affiliated with Novartis Consumer Health, Inc., owner of the registered trademark, Theraflu®.

192

C-3109-30508-6B-4





## TOPCARE NIGHTTIME SEVERE COLD AND COUGH HONEY LEMON INFUSED WITH CHAMOMILE AND WHITE TEA FLAVORS

acetaminophen, diphenhydramine hcl and phenylephrine hcl granule, for solution

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:36800-800
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>Acetaminophen</b> (UNII: 362O9ITL9D) (Acetaminophen - UNII:362O9ITL9D)	Acetaminophen	650 mg
<b>Diphenhydramine Hydrochloride</b> (UNII: TC2D6JAD40) (Diphenhydramine - UNII:8GTS82S83M)	Diphenhydramine Hydrochloride	25 mg
<b>Phenylephrine Hydrochloride</b> (UNII: 04JA59TNSJ) (Phenylephrine - UNII:1WS297W6MV)	Phenylephrine Hydrochloride	10 mg

### Inactive Ingredients

Ingredient Name	Strength
<b>Acesulfame</b> (UNII: MA3UYZ6K1H)	
<b>Aspartame</b> (UNII: Z0H242BBR1)	
<b>Citric acid monohydrate</b> (UNII: 2968PHW8QP)	
<b>Maltodextrin</b> (UNII: 7CVR7L4A2D)	
<b>Sodium Citrate</b> (UNII: 1Q73Q2JULR)	
<b>Sucrose</b> (UNII: C151H8M554)	

### Product Characteristics

<b>Color</b>	yellow (Caramel Color) , yellow (D & C Yellow 10)	<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>	HONEY (Natural Honey and Lemon Flavor) , LEMON (Natural Honey and Lemon Flavor)	<b>Imprint Code</b>	
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:36800-800-06	6 in 1 BOX		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	03/18/2014	

**Labeler** - TOP CARE (Topco Associates LLC) (006935977)

Revised: 5/2014

TOP CARE (Topco Associates LLC)