

## **MAGNESIUM CITRATE- magnesium citrate solution**

### **H E BUTT**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

### **Cherry Magnesium Citrate**

#### **164.000/164AA rev 2-AB**

### **TEP**

Do Not Use If Tamper Evident Twist-off Cap Is Missing, Broken or Separated From Neckring

### **Active ingredient**

Active ingredient in each fl oz

Magnesium citrate 1.745 g

### **Purpose**

Saline Laxative

### **Uses**

- for relief of occasional constipation (irregularity).
- Generally produces bowel movement in 1/2 to 6 hours

### **Warnings**

Ask a doctor before use if you have

- kidney disease
- a magnesium restricted diet
- abdominal pain, nausea, or vomiting
- noticed a sudden change in bowel habits that persists over a period of 2 weeks
- already used a laxative for a period longer than 1 week

### **Ask a doctor or pharmacist before use if you are**

- taking any other drug.
- Take this product 2 or more hours before or after other drugs. Laxatives may affect how other drugs work.

### **Stop use and ask a doctor if**

You have rectal bleeding or failure to have a bowel movement after use. These could be signs of a serious condition.

### **If pregnant**

**If pregnant or breast-feeding**, ask a health professional before use.

### **Keep out of reach of children**

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Store at temperatures between 46° and 86°F (8° and 30°C)

### **Directions**

- shake well before using
- drink a full glass (8 ounces) of liquid each dose
- may be taken as a single dose or in a divided doses

adults and children 12 years of age and over - 6.5 - 10 fl oz maximum 10 fl oz in 24 hours

children 6 to under 12 years of age - 3 to 7 fl oz maximum 7 fl oz in 24 hours

children 2 to under 6 years of age - 2 to 3 fl oz maximum 3 fl oz in 24 hours

children under 2 years of age- ask a doctor

### **other information**

- each fl oz contains: magnesium 290 mg
- each fl oz contains: sodium 1 mg

### **Inactive ingredients**

benzoic acid, citric acid, disodium EDTA, flavor, sucralose, water

### **Questions**

Call? 1-888-593-0593

### **Adverse reactions**

MADE WITH PRIDE & CARE FOR H-E-B

SAN ANTONIO, TX 78204

### **principal display panel**

H-E-B

Magnesium Citrate

Oral Solution

Saline Laxative

- Relief of: Occasional Constipation

Sugar & Dye Free

Gluten Free

Cherry Flavor

10 FL OZ (296 mL)



## **MAGNESIUM CITRATE**

magnesium citrate solution

**Product Information**

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:37808-673
<b>Route of Administration</b>	ORAL		

**Active Ingredient/Active Moiety**

<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
<b>MAGNESIUM CITRATE</b> (UNII: RHO26O1T9V) (MAGNESIUM CATION - UNII:T6V3LHY838)	MAGNESIUM CITRATE	0.059 g in 1 mL

**Inactive Ingredients**

<b>Ingredient Name</b>	<b>Strength</b>
<b>BENZOIC ACID</b> (UNII: 8SKN0B0MIM)	
<b>CITRIC ACID MONOHYDRATE</b> (UNII: 2968PHW8QP)	
<b>EDETATE DISODIUM ANHYDROUS</b> (UNII: 8NLQ36F6MM)	
<b>SUCRALOSE</b> (UNII: 96K6UQ3ZD4)	
<b>water</b> (UNII: 059QF0KO0R)	

**Packaging**

<b>#</b>	<b>Item Code</b>	<b>Package Description</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
1	NDC:37808-673-38	296 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	08/17/2021	

**Marketing Information**

<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
OTC monograph not final	part334	08/17/2021	

**Labeler** - H E BUTT (007924756)**Registrant** - Vi-Jon, LLC (790752542)**Establishment**

<b>Name</b>	<b>Address</b>	<b>ID/FEI</b>	<b>Business Operations</b>
Vi-Jon, LLC		790752542	manufacture(37808-673)

**Establishment**

<b>Name</b>	<b>Address</b>	<b>ID/FEI</b>	<b>Business Operations</b>
Vi-Jon, LLC		088520668	manufacture(37808-673)

