

**GOODYS COOL ORANGE HEADACHE- acetaminophen and aspirin and caffeine powder
Medtech Products Inc.**

Goodys Cool Orange Headache Powder

Drug Facts

Active ingredients (in each powder)

Purposes

Acetaminophen 325 mg.....	Pain reliever
Aspirin (NSAID*) 500 mg.....	Pain reliever
Caffeine 65 mg.....	Pain reliever aid

*nonsteroidal anti-inflammatory drug

Uses

Temporarily relieves minor aches and pains due to:

- headache
- backache
- toothache
- menstrual cramps
- muscular aches
- a cold
- minor arthritis pain

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- shock
- asthma (wheezing)

Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash. If a skin reaction occurs, stop use and seek medical help right away.

Liver warning: This product contains acetaminophen.

Severe liver damage may occur if you take

- more than 4 packs in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Caffeine warning: This product contains caffeine. Limit the use of caffeine-containing medications, foods, or beverages because too much caffeine may cause nervousness, irritability, sleeplessness, and occasionally, rapid heart beat.

Do not use

- if you have ever had an allergic reaction to aspirin or any other pain reliever/fever reducer
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

Ask a doctor before use if

- you have liver disease
- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma

Ask a doctor or pharmacist before use if you are taking

- a prescription drug for diabetes, gout or arthritis

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away
- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- any new symptoms appear
- ringing in the ears or a loss of hearing occurs

These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose can cause serious health problems. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not take more than directed** (see overdose warning)
- **adults and children 12 years of age and over:** open 1 pack, place powder on tongue every 6 hours, while symptoms persist. Drink a full glass of water with each dose, or may stir powder into glass of water or other liquid.
- do not take more than 4 packs in 24 hours unless directed by a doctor.
- **children under 12 years of age:** ask a doctor

Other information

Store below 25°C (77°F)

Inactive ingredients

colloidal silicon dioxide, flavors, guar gum, mannitol, microcrystalline cellulose, sucralose

Questions ?

1-866-255-5197

PRINCIPAL DISPLAY PANEL

Goody's ®

COOL ORANGE POWDER

ACETAMINOPHEN/ASPIRIN (NSAID)/ CAFFEINE

PAIN RELIEVER / PAIN RELIEVER AID

24 POWDERS



PRINCIPAL DISPLAY PANEL

Goody's ®

COOL ORANGE POWDER

ACETAMINOPHEN/ASPIRIN (NSAID)/ CAFFEINE

PAIN RELIEVER / PAIN RELIEVER AID

4 POWDERS

NO COPY

Drug Facts (continued) 3

• **See other drug contents:** This product contains acetaminophen, aspirin, and caffeine. **Do not use** if you have ever had an allergic reaction to aspirin or any other non-steroidal anti-inflammatory drug (NSAID), including aspirin, ibuprofen, naproxen, and acetaminophen, or if you have ever had an allergic reaction to caffeine or any other substance.

• **Contains warnings:** This product contains caffeine. Limit the use of caffeine-containing medications, foods, or beverages. **Do not use** if you have ever had an allergic reaction to aspirin or any other non-steroidal anti-inflammatory drug (NSAID), including aspirin, ibuprofen, naproxen, and acetaminophen, or if you have ever had an allergic reaction to caffeine or any other substance.

• **Do not use** if you have ever had an allergic reaction to aspirin or any other non-steroidal anti-inflammatory drug (NSAID), including aspirin, ibuprofen, naproxen, and acetaminophen, or if you have ever had an allergic reaction to caffeine or any other substance.

• **Do not use** if you have ever had an allergic reaction to aspirin or any other non-steroidal anti-inflammatory drug (NSAID), including aspirin, ibuprofen, naproxen, and acetaminophen, or if you have ever had an allergic reaction to caffeine or any other substance.

Drug Facts (continued) 4

• **See other drug contents:** This product contains acetaminophen, aspirin, and caffeine. **Do not use** if you have ever had an allergic reaction to aspirin or any other non-steroidal anti-inflammatory drug (NSAID), including aspirin, ibuprofen, naproxen, and acetaminophen, or if you have ever had an allergic reaction to caffeine or any other substance.

• **Contains warnings:** This product contains caffeine. Limit the use of caffeine-containing medications, foods, or beverages. **Do not use** if you have ever had an allergic reaction to aspirin or any other non-steroidal anti-inflammatory drug (NSAID), including aspirin, ibuprofen, naproxen, and acetaminophen, or if you have ever had an allergic reaction to caffeine or any other substance.

• **Do not use** if you have ever had an allergic reaction to aspirin or any other non-steroidal anti-inflammatory drug (NSAID), including aspirin, ibuprofen, naproxen, and acetaminophen, or if you have ever had an allergic reaction to caffeine or any other substance.

• **Do not use** if you have ever had an allergic reaction to aspirin or any other non-steroidal anti-inflammatory drug (NSAID), including aspirin, ibuprofen, naproxen, and acetaminophen, or if you have ever had an allergic reaction to caffeine or any other substance.

Drug Facts (continued) 5

• **See other drug contents:** This product contains acetaminophen, aspirin, and caffeine. **Do not use** if you have ever had an allergic reaction to aspirin or any other non-steroidal anti-inflammatory drug (NSAID), including aspirin, ibuprofen, naproxen, and acetaminophen, or if you have ever had an allergic reaction to caffeine or any other substance.

• **Contains warnings:** This product contains caffeine. Limit the use of caffeine-containing medications, foods, or beverages. **Do not use** if you have ever had an allergic reaction to aspirin or any other non-steroidal anti-inflammatory drug (NSAID), including aspirin, ibuprofen, naproxen, and acetaminophen, or if you have ever had an allergic reaction to caffeine or any other substance.

• **Do not use** if you have ever had an allergic reaction to aspirin or any other non-steroidal anti-inflammatory drug (NSAID), including aspirin, ibuprofen, naproxen, and acetaminophen, or if you have ever had an allergic reaction to caffeine or any other substance.

• **Do not use** if you have ever had an allergic reaction to aspirin or any other non-steroidal anti-inflammatory drug (NSAID), including aspirin, ibuprofen, naproxen, and acetaminophen, or if you have ever had an allergic reaction to caffeine or any other substance.

Goody's
COOL ORANGE

FAST PAIN RELIEF

Goody's
COOL ORANGE
POWDER

ACETAMINOPHEN
ASPIRIN (NSAID) / CAFFEINE
PAIN RELIEVER / PAIN RELIEVER AID

4 PACKS

Goody's

© 2013 Watson Pharmaceuticals, Inc. All rights reserved. Watson Pharmaceuticals, Inc. is a subsidiary of Johnson & Johnson.

TO OPEN CHILD-RESISTANT PACKAGE

1. Slide pack bottom.

2. Pushed tear at other corner.

8 42037 00060 0

852265

Drug Facts

Warnings

• **Do not use** if you have ever had an allergic reaction to aspirin or any other non-steroidal anti-inflammatory drug (NSAID), including aspirin, ibuprofen, naproxen, and acetaminophen, or if you have ever had an allergic reaction to caffeine or any other substance.

• **Do not use** if you have ever had an allergic reaction to aspirin or any other non-steroidal anti-inflammatory drug (NSAID), including aspirin, ibuprofen, naproxen, and acetaminophen, or if you have ever had an allergic reaction to caffeine or any other substance.

• **Do not use** if you have ever had an allergic reaction to aspirin or any other non-steroidal anti-inflammatory drug (NSAID), including aspirin, ibuprofen, naproxen, and acetaminophen, or if you have ever had an allergic reaction to caffeine or any other substance.

• **Do not use** if you have ever had an allergic reaction to aspirin or any other non-steroidal anti-inflammatory drug (NSAID), including aspirin, ibuprofen, naproxen, and acetaminophen, or if you have ever had an allergic reaction to caffeine or any other substance.

Directions

• **Adults and teenagers:** Take 1 packet every 4 to 6 hours, which is the same as 2 tablets every 4 to 6 hours. Do not take more than 4 packets in 24 hours, which is the same as 8 tablets in 24 hours. Do not take more than 4 packets in 24 hours, which is the same as 8 tablets in 24 hours. Do not take more than 4 packets in 24 hours, which is the same as 8 tablets in 24 hours.

• **Children:** See the dosing chart for children's use. Do not give to children under 12 years of age.

• **Elderly and frail patients:** Use with caution. Do not take more than 4 packets in 24 hours, which is the same as 8 tablets in 24 hours.

Drug Facts (continued) 2

• **See other drug contents:** This product contains acetaminophen, aspirin, and caffeine. **Do not use** if you have ever had an allergic reaction to aspirin or any other non-steroidal anti-inflammatory drug (NSAID), including aspirin, ibuprofen, naproxen, and acetaminophen, or if you have ever had an allergic reaction to caffeine or any other substance.

• **Contains warnings:** This product contains caffeine. Limit the use of caffeine-containing medications, foods, or beverages. **Do not use** if you have ever had an allergic reaction to aspirin or any other non-steroidal anti-inflammatory drug (NSAID), including aspirin, ibuprofen, naproxen, and acetaminophen, or if you have ever had an allergic reaction to caffeine or any other substance.

• **Do not use** if you have ever had an allergic reaction to aspirin or any other non-steroidal anti-inflammatory drug (NSAID), including aspirin, ibuprofen, naproxen, and acetaminophen, or if you have ever had an allergic reaction to caffeine or any other substance.

• **Do not use** if you have ever had an allergic reaction to aspirin or any other non-steroidal anti-inflammatory drug (NSAID), including aspirin, ibuprofen, naproxen, and acetaminophen, or if you have ever had an allergic reaction to caffeine or any other substance.

Drug Facts (continued) 5	Drug Facts (continued) 6
<ul style="list-style-type: none"> ringing in the ears or a loss of hearing occurs. These could be signs of a serious condition. 	Directions <ul style="list-style-type: none"> do not take more than directed (see overdose warning). adults and children 12 years and over: open 1 pack, place powder on tongue every 6 hours, while symptoms persist. Drink a full glass of water with each dose, or stir powder into a glass of water or other liquid. do not take more than 4 packs in 24 hours unless directed by a doctor. children under 12 years: ask a doctor.
<p>If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.</p> <p>Keep out of reach of children.</p>	Other information Store below 25°C (77°F)
<p>Overdose warning: Taking more than the recommended dose can cause serious health problems. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.</p>	Inactive ingredients colloidal silicon dioxide, flavors, guar gum, mannitol, microcrystalline cellulose, sucralose
	Questions? 1-866-255-6197 goodyspowder.com

GOODYS COOL ORANGE HEADACHE

acetaminophen and aspirin and caffeine powder

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:63029-663
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	500 mg
CAFFEINE (UNII: 3G6A5W338E) (CAFFEINE - UNII:3G6A5W338E)	CAFFEINE	65 mg

Inactive Ingredients

Ingredient Name	Strength
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
GUAR GUM (UNII: E89I1637KE)	
MANNITOL (UNII: 3OWL53L36A)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	

Product Characteristics

Color		Score	
Shape		Size	
Flavor	ORANGE	Imprint Code	

Contains**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:63029-663-01	24 in 1 BOX; Type 0: Not a Combination Product	06/01/2012	
2	NDC:63029-663-04	4 in 1 BOX; Type 0: Not a Combination Product	06/01/2012	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	06/01/2012	

Labeler - Medtech Products Inc. (122715688)

Revised: 10/2025

Medtech Products Inc.