REGENCY POST MILKING TEAT DIP- post-milking phenol teat dip solution CLSolutions

Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click here.

Regency Post-Milking Teat Dip

ACTIVE INGREDIENTS:

Active Ingredients	Purpose
PHENOL (1.56% w/v)	Disinfectant
PHENOLATE SODIUM (0.06% w/v)	Disinfectant

PURPOSE:

Active Ingredients	Purpose
Phenol (1.56% w/v)	Disinfectant
Phenolate Sodium (0.06% w/v)	Disinfectant

Keep Out of Reach of Children

If swallowed, get medical help or contact Poison Control Center right away.

USES:

FOR ANIMAL USE ONLY

Aids in controlling mastitis organisms with a softening effect on teats.

Directions for Use:

Before milking, wash udder and teats with damp cloth. Immediately after milking, dip each teat 1 to 2 inches into REGENCY Teat Dip.

Allow each teat to air dry. Do not wipe.

During the winter months, keep cows inside until the teats have completely dried to prevent freezing and chapping.

When cow is being dried out, dip each teat in REGENCY 1X Post-Milking Teat Dip once a day for 3 to 4 days after the last milking.

CAUTION:

Avoid eye contact. Do not inhale fumes. For external use only.

FIRST AID:

For Ingestion - Immediately give castor oil or vegetable oil (15 -30cc). Get medical attention immediately.

Be ready tp induce vomiting at the advice of a physician or poison control center.

For Eyes - Immediately flush eyes with water for at least 15 minutes. Wash affected skin with water.

When Using this product:

Do not Store in Direct Sunlight.

Avoid Excessive Heat.

USER SAFETY WARNING:

Do Not Allow Product To Freeze. If frozen, allow product to completely than in a warm room.

Shake and thoroughly mix product before using.

RInse teat dip cup with water after each milking and invert to dry.

Never pour used teat dip into original container.

Keep cap on container when not in use.

INACTIVE INGREDIENTS:

Water, Glycerin, Linear Alkylbenzene Sulfonate, Sodium Borate, Fragrance, FD&C Blue # 1



FIRST AID: For Ingestion - Immediately give castor oil or vegetable oil (15 - 30cc). Get medical attention immediately. Be ready to induce vomiting at the advice of a physician or poison control center. For Eyes - Immediately flush eyes with water for at least 15 minutes. Wash affected skin with water.

CAUTION: Avoid eye contact. For external use only

DIRECTIONS: Before milking, wash udder and teats with damp cloth. Immediately after milking, dip each teat 1 to 2 inches into Regency 1X Post-Milking Teat Dip.

Allow each teat to air dry. Do not wipe. During the winter months, keep cows inside until the teats have completely dried to prevent freezing and chapping. If the solution in the dip cup becomes contaminated, discard and use fresh solution. Rinse teat dip cup with water after each milking and invert to dry. Never pour used teat dip into the original container. Keep cap on container when not in use

When a cow is being dried out, dip each teat in Regency 1X Post-Milking Teat Dip once a day for 3 to 4 days after the last milking. Do not allow product to freeze. If frozen, allow the product to completely thaw in a warm room. Shake and thoroughly mix product before using.

INGREDIENTS: Water, Glycerin, Phenol, LAS (Linear Alkylbenzene Sulfonate), Sodium Borate, Fragrance, FD&C Blue #1.

CLSolutions, LLC PO BOX 837 Fulton, TX 78358 (361) 729-3993 www.CoastalLivestock.com Sporicidin* is a registered trademark of Contec, Inc. United Sates Patent # US6,586,477 B1 CLS-1X-1GAL



Regency Masticide "Ready To USe Teat Dip" with Sporicidin Technology

1X Post-Milking Teat Dip

Aids in controlling mastitis organisms with a softening effect on the teats

1.56 % % (w/w) Phenol with 0.06 % (w/v) Sodium Phenate

Teat Dip Will Not Burn, Dry, Irritate or Chap Teats. Contains a special blend of emollients and phenol.

NET CONTENTS: 1 U.S. Gallon (3.785 L)



Aids in controlling mastitis organisms with a softening effect on the teats.

1.56% (w/v) Phenol with 0.06% (w/v) Sodium Phenate Teat Dip Will Not Burn, Dry, Irritate Or Chap Teats. Contains a special blend of emollients and phenol.

NOTE: Store at room temperature away from direct sunlight.

NET CONTENTS: 5 U.S. Gallons (18.927 L)

FOR ANIMAL USE ONLY KEEP OUT OF REACH OF CHILDREN DO NOT STORE IN DIRECT SUNLIGHT AVOID EXCESSIVE HEAT



FIRST AID: For Ingestion - Immediately give castor oil or vegetable oil (15 - 30cc). Get medical attention immediately. Be ready to induce vomiting at the advice of a physician or poison control center. For Eyes - Immediately flush eyes with water for at least 15 minutes. Wash affected skin with water.

CAUTION: Avoid eve contact. For external use only.

DIRECTIONS: Before milking, wash udder and teats with damp cloth. Immediately after milking, dip each teat 1 to 2 inches into Regency 1X Post-Milking Teat Dip.

Allow each teat to air dry. Do not wipe. During the winter months, keep cows inside until the teats have completely dried to prevent freezing and chapping. If the solution in the dip cup becomes contaminated discard and use fresh solution. Rinse teat dip cup with water after each milking and invert to dry. Never pour used teat dip into the original container. Keep cap on container when not in use.

When a cow is being dried out, dip each teat in Regency 1X Post-Milking Teat Dip once a day for 3 to 4 days after the last milking. Do not allow product to freeze. If frozen, allow the product to completely thaw in a warm room. Shake and thoroughly mix product before using.

INGREDIENTS: Water, Glycerin, Phenol, LAS (Linear Alkylbenzene Sulfonate), Sodium Borate, Fragrance, FD&C Blue #1.

Distributed Exclusively By: CLSolutions, LLC PO BOX 837 Fulton, TX 78358 (361) 729-3993 www.CoastalLivestock.com

Sporicidin* is a registered trademark of Cor United Sates Patent # US6,586,477 B1 CLS-1X-5GAL



Regency Masticide "Ready To USe Teat Dip" with Sporicidin Technology

1X Post-Milking Teat Dip

Aids in controlling mastitis organisms with a softening effect on the teats

1.56% % (w/w) Phenol with 0.06% (w/v) Sodium Phenate

Teat Dip Will Not Burn, Dry, Irritate or Chap Teats. Contains a special blend of emollients and phenol.

NET CONTENTS: 5 U.S. Gallon (18.927 L)



KEEP OUT OF REACH OF CHILDREN DO NOT STORE IN DIRECT SUNLIGHT AVOID EXCESSIVE HEAT



FIRST AID: For Ingestion - Immediately give castor oil or vegetable oil (15 - 30cc). Get medical attention immediately. Be ready to induce vomiting at the advice of a physician or poison control center. For Eyes - Immediately flush eyes with water for at least 15 minutes. Wash affected skin with water.

CAUTION: Avoid eye contact. For external use only.

DIRECTIONS: Before milking, wash udder and teats with damp cloth. Immediately after milking, dip each teat 1 to 2 inches into Regency 1X Post-Milking Teat Dip.


Allow each teat to air dry. Do not wipe. During the winter months, keep cows inside until the teats have completely dried to prevent freezing and chapping. If the solution in the dip cup becomes contaminated, discard and use fresh solution. Rinse teat dip cup with water after each milking and invert to dry. Never pour used teat dip into the original container. Keep cap on container when not in use.

When a cow is being dried out, dip each teat in Regency 1X Post-Milking Teat Dip once a day for 3 to 4 days after the last milking. Do not allow product to freeze. If frozen, allow the product to completely thaw in a warm room. Shake and thoroughly mix product before using.

INGREDIENTS: Water, Glycerin, Phenol, LAS (Linear Alkylbenzene Sulfonate), Sodium Borate, Xanthan Gum, Fragrance, FD&C Blue #1.

Distributed Exclusively By: CLSolutions, LLCTM 7031 North 16th St. Omaha, NE 68112 (402) 502-3928

Sporicidin* is a registered trademark of Contec, Inc. United Sates Patent # US6,586,477 B1



Regency Masticide "Ready To USe Teat Dip" with Sporicidin Technology

1X Post-Milking Teat Dip

special blend of emollients and phenol

NOTE: Store at room temperature away from direct sunlight.

NET CONTENTS: 55 U.S. Gallons (208.2 L)

Aids in controlling mastitis organisms with a softening effect on the teats

1.56%% (w/w) Phenol with 0.06% (w/v) Sodium Phenate

Teat Dip Will Not Burn, Dry, Irritate or Chap Teats. Contains a special blend of emollients and phenol.

NET CONTENTS: 55 U.S. Gallon (208.2 L)



Aids in controlling mastitis organisms with a softening effect on the teats.

1.56% (w/v) Phenol with 0.06% (w/v) Sodium Phenate Teat Dip Will Not Burn, Dry, Irritate Or Chap Teats. Contains a special blend of emollients and phenol.

NOTE: Store at room temperature away from direct sunlight.

NET CONTENTS: 270 U.S. Gallons

FIRST AID: For Ingestion - Immediately give castor oil or vegetable oil (16 - 30cc). Get medical attention immediately. Be ready to induce vomiting at the advice of a physician or poison control center. For Eyes - Immediately flush eyes with water for at least 15 minutes. Wash affected skin with water.

CAUTION: Avoid eye contact. For external use only.

DIRECTIONS: Before milking, wash udder and teats with damp cloth. Immediately after milking, dip teat entirely into Regency® Masticide® 1X Post-Milking Teat Dip.

Allow each teat to air dry. Do not wipe. During the winter months, keep cows inside until the teats have completely dried to prevent freezing and chapping. If the solution in the dip cup becomes contaminated, discard and use fresh solution. Rinse teat dip cup with water after each milking and invert to dry. Never pour used teat dip into the original container. Keep cap on container when not in use.

When a cow is being dried out, dip each teat in Regency® Masticide® 1X Post-Milking Teat Dip once a day for 3 to 4 days after the last milking.

Do not allow product to freeze. If frozen, allow the product to completely thaw in a warm room. Shake and thoroughly mix product before using.

INGREDIENTS: Water, Glycerin, Phenol, LAS (Linear Alkylbenzene Sulfonate), Sodium Borate, Fragrance, FD&C Blue #1.

Distributed Exclusively By: CLSolutions, LLCTM 7031 North 16th St. Omaha, NE 68112 (402) 502-3928 www.clsag.com CLS-1X-REV-8 Sporicidin* is a registered trademark of Contec, Inc. United Sates Patent # US6,586,477 B1

LOT NO.

Regency Masticide "Ready To USe Teat Dip" with Sporicidin Technology

1X Post-Milking Teat Dip

Aids in controlling mastitis organisms with a softening effect on the teats

1.56 % % (w/w) Phenol with 0.06 % (w/v) Sodium Phenate

Teat Dip Will Not Burn, Dry, Irritate or Chap Teats. Contains a special blend of emollients and phenol.

Technology

FOR ANIMAL USE ONLY

KEEP OUT OF REACH OF CHILDREN

DO NOT STORE IN DIRECT SUNLIGHT

AVOID EXCESSIVE HEAT

OTC

NET CONTENTS: 270 U.S. Gallons

REGENCY POST MILKING TEAT DIP

post-milking phenol teat dip solution

Product Information				
Product Type	OTC ANIMAL DRUG	Item Code (Source)	NDC:86018-010	
Route of Administration	TOPICAL			

Active Ingredient/Active Moiety				
Ingredient Name	Basis of Strength	Strength		
PHENOL (UNII: 339 NCG44TV) (PHENOL - UNII:339 NCG44TV)	PHENOL	156 g in 1 L		
PHENOLATE SODIUM (UNII: 4NC0T56V35) (PHENOL - UNII:339NCG44TV)	PHENOL	0.06 g in 1 L		

Inactive Ingredients	
Ingredient Name	Strength
SO DIUM HYDRO XIDE (UNII: 55X04QC32I)	
SODIUM BORATE (UNII: 91MBZ8H3QO)	

GLYCERIN (UNII: PDC6 A3C0 O X)	
P-DODECYLBENZENESULFONIC ACID (UNII: OC21S23N1O)	
XANTHAN GUM (UNII: TTV12P4NEE)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
WATER (UNII: 059QF0KO0R)	

Packaging					
#	Item Code	Package Description	Marketing Start Date	Marketing End Date	
1	NDC:86018-010-04	1021.95 L in 1 CONTAINER			
2	NDC:86018-010-01	3.785 L in 1 JUG			
3	NDC:86018-010-02	18.925 L in 1 PAIL			
4	NDC:86018-010-03	208.175 L in 1 DRUM			

Marketing Infor	mation		
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		06/13/2011	

Labeler - CLSolutions (079338118)

Registrant - ABS Corporation (048022727)

Establishment			
Name	Address	ID/FEI	Business Operations
ABS Corporation		048022727	api manufacture, pack, label, manufacture

Revised: 5/2014 CLSolutions