

CHILDRENS TYLENOL- acetaminophen suspension
Kenvue Brands LLC

Children's Tylenol®

Drug Facts

Active ingredient

Active ingredient (in each 5 mL)

Acetaminophen 160 mg

Purpose

Purpose

Pain reliever/fever reducer

Uses

Uses temporarily:

- reduces fever
- relieves minor aches and pains due to:
 - the common cold
 - flu
 - headache
 - sore throat
 - toothache

Warnings

Liver warning

Liver warning: This product contains acetaminophen. Severe liver damage may occur if your child takes

- more than 5 doses in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen

Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: if sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if your child is allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if your child has liver disease

Ask a doctor or pharmacist before use if your child is taking the blood thinning drug warfarin

When using this product do not exceed recommended dose (see overdose warning)

Stop use and ask a doctor if

- pain gets worse or lasts more than 5 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away.

(1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

Directions

■ **this product does not contain directions or complete warnings for adult use**

■ **do not give more than directed (see overdose warning)**

■ **shake well before using**

■ mL = milliliter

■ find right dose on chart below. If possible, use weight to dose; otherwise, use age.

■ remove the child protective cap and squeeze your child's dose into the dosing cup

■ repeat dose every 4 hours while symptoms last

■ do not give more than 5 times in 24 hours

Weight (lb)	Age (yr)	Dose (mL)*
under 24	under 2 years	ask a doctor
24-35 lbs	2-3 years	5 mL
36-47 lbs	4-5 years	7.5 mL

48-59 lbs	6-8 years	10 mL
60-71 lbs	9-10 years	12.5 mL
72-95 lbs	11 years	15 mL

* or as directed by a doctor

Attention: use only enclosed dosing cup specifically designed for use with this product. Do not use any other dosing device.

Other information

Other information

- each 5 mL contains: **potassium 5 mg**
- store between 20-25°C (68-77°F)
- **do not use if carton is opened. Do not use if bottle wrap imprinted with "TYLENOL" is broken or missing**

Inactive ingredients

anhydrous citric acid, flavors, glycerin, microcrystalline cellulose and carboxymethylcellulose sodium, potassium sorbate, purified water, sodium benzoate, sodium citrate, sorbitol solution, sucralose, sucrose, xanthan gum

Questions or comments? call **1-800-458-1635** (toll-free) or **215-273-8755** (collect)

NDC 50580-139-04

NEW

Children's TYLENOL®

Acetaminophen (160 mg per 5 mL)

Oral Suspension

Pain Reliever-Fever Reducer

Pain+Fever

Ages 2-11 Years

DYE-FREE

Free Of:

- Dyes
- Alcohol
- Ibuprofen
- Aspirin
- Parabens
- High Fructose Corn Syrup

• Artificial Flavoring

Natural Apple Flavor with Other Natural Flavors

4 fl oz (120 mL) 160 mg per 5 mL



OPEN HERE →

How can we help?
☎ 1-800-458-1635

NDC 50580-139-04

NEW

Children's TYLENOL®

Acetaminophen (160 mg per 5 mL)
Oral Suspension
Pain Reliever-Fever Reducer
Pain + Fever
Ages 2-11 Years

DYE-FREE

Free Of:

- Dyes
- Alcohol
- Ibuprofen
- Aspirin
- Parabens
- High Fructose Corn Syrup
- Artificial Flavoring

Natural Apple Flavor with Other Natural Flavors



4 fl oz (120 mL) 160 mg per 5 mL

Children's TYLENOL®

👉 **Pediatricians recommend TYLENOL® for pain and fever**
👉 **Relied on by moms for over 60 years**

DIRECTIONS:

- STEP 1.** Shake well before using 
- STEP 2.** Remove child protective cap 
- STEP 3.** Squeeze your child's dose into the dosing cup 

IMPORTANT:

Please see dosing chart or ask a doctor for your child's correct dose

Drug Facts

Active ingredient (in each 5 mL) Purpose
Acetaminophen 160 mg.....Pain reliever/fever reducer.

Uses temporarily:

- reduces fever
- relieves minor aches and pains due to:
 - the common cold
 - flu
 - sore throat
 - headache
 - toothache

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if your child takes more than 5 doses in 24 hours, which is the maximum daily amount

■ with other drugs containing acetaminophen
Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include:
■ skin reddening ■ blisters ■ rash
If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: if sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if your child is allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if your child has liver disease
Ask a doctor or pharmacist before use if your child is taking the blood thinning drug warfarin

When using this product do not exceed recommended dose (see overdose warning)

Stop use and ask a doctor if

- pain gets worse or lasts more than 5 days
 - fever gets worse or lasts more than 3 days
 - new symptoms occur
 - redness or swelling is present
- These could be signs of a serious condition.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Drug Facts (continued)

Directions

- this product does not contain directions or complete warnings for adult use
- do not give more than directed (see overdose warning)
- shake well before using
- mL = milliliter
- find right dose on chart below. If possible, use weight to dose; otherwise, use age.
- remove the child protective cap and squeeze your child's dose into the dosing cup
- repeat dose every 4 hours while symptoms last
- do not give more than 5 times in 24 hours

Weight (lb)	Age (yr)	Dose (mL)*
under 24	under 2 years	ask a doctor
24-35 lbs	2-3 years	5 mL
36-47 lbs	4-5 years	7.5 mL
48-59 lbs	6-8 years	10 mL
60-71 lbs	9-10 years	12.5 mL
72-95 lbs	11 years	15 mL

* or as directed by a doctor

Attention: use only enclosed dosing cup specifically designed for use with this product. Do not use any other dosing device.

Other information

- each 5 mL contains: potassium 5 mg
- store between 20-25°C (68-77°F)
- do not use if carton is opened. Do not use if bottle wrap imprinted with "TYLENOL" is broken or missing

Inactive ingredients

anhydrous citric acid, flavors, glycerin, microcrystalline cellulose and carboxymethylcellulose sodium, potassium sorbate, purified water, sodium benzoate, sodium citrate, sorbitol solution, sucralose, sucrose, xanthan gum

Questions or comments? call 1-800-458-1635 (toll-free) or 215-273-8755 (collect)

Distributed by:
Kenvue Brands LLC
Summit, NJ 07901
© Kenvue Brands LLC 2025
Pat. www.kenvuepats.com
Visit us at www.tylenol.com

30058492



3 00450 14604 5

acetaminophen suspension

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:50580-139
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	160 mg in 5 mL

Inactive Ingredients

Ingredient Name	Strength
SODIUM CITRATE (UNII: 1Q73Q2JULR)	
SUCROSE (UNII: C151H8M554)	
XANTHAN GUM (UNII: TTV12P4NEE)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POTASSIUM SORBATE (UNII: 1VPU26JZZ4)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
CARBOXYMETHYLCELLULOSE SODIUM, UNSPECIFIED (UNII: K679OBS311)	
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
GLYCERIN (UNII: PDC6A3C0OX)	
SORBITOL SOLUTION (UNII: 8KW3E207O2)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
WATER (UNII: 059QF0KO0R)	

Product Characteristics

Color	white (White to off-white)	Score	
Shape		Size	
Flavor	APPLE	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:50580-139-04	1 in 1 CARTON	04/28/2025	
1		120 mL in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:50580-139-08	1 in 1 CARTON	06/30/2025	
2		240 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	04/28/2025	

Labeler - Kenvue Brands LLC (118772437)

Revised: 5/2025

Kenvue Brands LLC